

Importance of Molecular Diagnostic Methods for infectious Diseases

Sajida Sboui¹, Ahmed Tabbabi^{2*}

¹Faculty of Medicine of Monastir, Monastir University, Monastir, Tunisia

²Department of Hygiene and Environmental Protection, Ministry of Public Health, Tunis, Tunisia

***Corresponding Author:** Ahmed Tabbabi, Department of Hygiene and Environmental Protection, Ministry of Public Health, Tunis, Tunisia

SHORT COMMUNICATION

Molecular methods are no longer available only too specialized centers and now play an important role in the diagnosis of common infections seen in ambulatory practice. It is especially for the detection of pathogens that are difficult to cultivate, especially viruses that PCR has become the reference method. While it has replaced traditional methods for some indications, PCR is not applicable in all cases, and it is not foolproof. It is therefore important to know when to use molecular methods, what are their strengths and weaknesses, in order to be able to prescribe them rationally. This article reviews the characteristics of molecular tests and their main applications in outpatient practice.

The direct detection of the microbe by microscopy or by characterization of its molecular structures (proteins or nucleic acids) and the measurement of humoral immune response (specific antibodies) are the main tools of the laboratory diagnosis of infectious diseases. The molecular diagnostics methods consist to detect and analyze the genome of an organism even if antibodies and antigens are molecular structures. The first methods of DNA analysis already existed in the 1970s (1) but the development of Polymerase Chain Reaction (PCR) appeared in the mid-1980s (2, 3). Right after, the possibility of amplifying DNA and RNA (ribonucleic acid) was used for the detection of pathogens and PCR was considered as a central method of the microbiology laboratory thirty years later.

Molecular methods allow diagnosis in the acute phase of a disease, before the appearance of antibodies exactly like antigen detection. However, few days or even a few weeks are necessary to detect the presence of antibodies in the serum (4). In the acute phase, a positive PCR

demonstrates most often an infection, whereas a negative serology must be repeated within a suitable time to demonstrate sero conversion (4, 5).

By allowing the detection of pathogens not or difficult to cultivate, molecular methods have revolutionized medical microbiology, mainly for the diagnosis of viral diseases. Indeed, virus culture requires immortalized human or animal cell lines and may take several weeks. It is less and less used and has been largely supplanted by PCR, which has also replaced serology for many indications, such as acute herpes or varicella zoster infection or shingles (5, 6). Contrary to viruses, molecular techniques have not replaced cultures for bacteria or fungi. However, the detection of certain slow-growing bacteria requiring very specific culture conditions (mycobacteria, Chlamydia) has also been improved and simplified by the development of specific PCRs. PCR detection is available for many parasitic infections such as toxoplasmosis, leishmaniasis, amebiasis or malaria (7). For these diseases, traditional methods (microscopy or serology) are still widely used.

In addition to simple detection, modern molecular methods can also provide quantitative indications, for example to monitor the evolution of HIV viral load or hepatitis C under therapy, and thus detect the emergence of resistant strains. This quantification is also useful for the diagnosis of latent virus reactivations such as polyoma virus BK or CMV (cytomegalovirus) in immune suppressed patients (8).

Although DNA sequencing was developed before PCR, all modern methods are based on it. Its most common application in microbiology is the detection and identification of bacteria by sequencing ribosomal DNA. The detection of

Importance of Molecular Diagnostic Methods for infectious Diseases

mutations resulting from resistance to therapies is another application of sequencing, used, for example, for the characterization of HIV.

REFERENCES

- [1] Southern EM. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* 1975; 98:503-17.
- [2] Saiki RK, Scharf S, Faloona F, et al. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 1985; 230:1350-4.
- [3] Mullis K, Faloona F, Scharf S, et al. Specific enzymatic amplification of DNA in vitro : The polymerase chain reaction. *Cold Spring Harb Symp Quant Biol* 1986; 51: 263-73.
- [4] Péter O. A quoi sert la sérologie en pratique ambulatoire ? *Med Hyg* 1999;57:747-53.
- [5] Lienhard R. Pièges en sérologie infectieuse. *Rev Med Suisse* 2011;7:1964-7.
- [6] Meylan P. Infections à virus de l'herpès simplex : mise à jour pour le praticien. *Rev Med Suisse* 2011;7: 886-93.
- [7] L Dormond K, Jatou-Ogay S, de Valliere Multiplex real-time PCR for the diagnosis of malaria: Correlation with microscopy. *Clin Microbiol Infect.* 2011 Mar;17(3):469-75.
- [8] Meylan P. Infections à virus de l'herpès simplex : mise à jour pour le praticien. *Rev Med Suisse* 2011;7: 886-93.

Citation: *Sajida Sboui " Importance of Molecular Diagnostic Methods for infectious Diseasesm Annals of Microbiology and Infectious Diseases 2(1), 2019, pp.67-68.*

Copyright: © 2019 Sajida Sboui, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.