

SHORT COMMUNICATION

A New Proof of Zero Active Mass Equation a Work on Non-Accelerating Universes

Fernando Salmon Iza

Physicist. Complutense University of Madrid. Independent Researcher, Spain.

Received: 14 July 2026 Accepted: 27 July 2026 Published: 03 August 2026

Corresponding Author: Fernando Salmon Iza, Physicist. Complutense University of Madrid. Independent Researcher, Spain.
fernandosalmoniza@gmail.com

Abstract

For decades, the prevailing view in the scientific community has been that our universe is expanding at an accelerating rate. However, a recent experimental study (Junhjuk Son et al., 2025) seems to confirm that our universe shows no signs of acceleration; that is, we might be living in a non-accelerating universe. In this work, we present evidence that this result agrees with the predictions of relativistic theory. To this end, we present a new proof of the zero active mass equation ($\rho + 3p = 0$) for a universe with FLRW metric and zero spatial curvature ($k = 0$). We study the expansion of the universe by analyzing two partial models: a classical energy model and a thermodynamic fluid dynamics model. The interrelation between these two models demonstrates that all universes with zero spatial curvature, a Lapse function equal to one, satisfy the zero active mass equation and are necessarily non-accelerating. Their linear expansion is solely a consequence of the pressure of the cosmic fluid and not of any other additional terms in the Friedmann equations.

Keywords: Large-Scale Structure of Universe, Cosmology: Theory, Cosmology: Miscellaneous.

1. Introduction

In 2022, Professor Fulvio Melia published a report in *Astronomische Nachrichten* entitled “Initial Energy of a Spatially Flat Universe: A Key to Its Possible Origin” (Melia, F. 2022). In it, he studies the expansion process of the universe from its beginnings and concludes that, according to the Friedmann equations, the spatial curvature of the universe, k , is proportional to the sum of the kinetic and potential energies involved in the expansion process. Therefore, for zero spatial curvature, as is the case with our universe, the total mechanical energy of the expansion process is zero.

Professor Fulvio Melia is also very optimistic about the results he is obtaining with his $R_h = ct$ universe model (Melia F., 2021), which constitutes a valid alternative to the Λ CDM model according to the most recent experimental data (Hubble Tension, uniformity of the cosmic microwave background radiation, age of the most distant galaxies, etc.), where the Λ CDM

model falls short. The $R_h = ct$ universe is a universe with FLRW metric and zero curvature, based on the study of the R_h gravitational horizon (Melia F, 2021) and two equations: the $R_h = ct$ constraint equation and the zero active mass equation ($\rho + 3p = 0$). The latter conditions the scale factor by setting $a'' = 0$, resulting in a non-accelerating universe.

In this work, we will demonstrate the zero active mass equation using a different method than the one currently employed to demonstrate it (Melia 2026). Here, we use the kinetic and potential energies involved in the expansion process (Melia F, 2022) and a thermodynamic universe, finding similarities between the expansion of the universe and the expansion of a gas in a piston. Therefore, we consider the thermodynamic principles related to the fluid model, applicable to all universes with FLRW metric. At all times, we assume a universe with zero spatial curvature, as deduced from Planck data (Planck, 2018).

Citation: Fernando Salmon Iza, A New Proof of Zero Active Mass Equation a Work on Non-Accelerating Universes. Open Access Journal of Physics. 2026;8(3): 114-117.

©The Author(s) 2026. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

2. Development of the Zero Active Mass Equation

2.1 Friedmann Equations

Given the Friedmann equations, of the FLRW metric:

$$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G \rho}{3c^2} - \frac{kc^2}{a^2}$$

$$\left(\frac{\ddot{a}}{a}\right) = -\frac{4\pi G}{3c^2} (\rho + 3p)$$

ρ represents the total energy density in the cosmos, (ρ = Joules/m³)

2.2 Model 1. Expansion of the Universe According to a Work by Fulvio Melia Referenced Here

Here is a brief summary of this model, according to the content of the cited paper (Melia F, 2022), we consider an isotropic, homogeneous and spatially spherical universe, which responds to the FLRW metric and which therefore expands. This universe has a certain energy density ρ at each instant of time. We are going to refer our calculations to an observer located in the center of it. We call R_h its gravitational horizon (Melia F, 2021) and $M(r)$ the mass, which comes from its energy density ρ , contained in a sphere of radius “ r ” centered at the observer’s point, $r < R_h$

We are going to calculate the kinetic energy and potential energy produced by the expansion of the universe for that observer considering that sphere. The increase in its kinetic energy, “ ΔK ”, during the expansion process is given by:

$$\Delta K = (4\pi r^2 \rho \Delta r) (\Delta r / \Delta t)^2 / 2$$

and the corresponding increase in its potential energy “ ΔU ”, is given by:

$$\Delta U = -(4\pi r^2 \rho \Delta r) GM(r) / r$$

In the expansion process, this potential energy will correspond to the work done by dark energy against gravitational forces.

According to the conclusions of the work of Fulvio Melia (Melia F, 2022), the spatial curvature of the universe k is proportional to the sum of the kinetic K and potential U energies involved in the process of expansion of the universe

$$k \sim K + U$$

Thus, in a universe of zero curvature $k=0$, the following holds true:

$$0 = K + U$$

$$0 = dK + dU$$

$$dK = -dU$$

2.3 Model 2. Expansion of the Universe According to a Gaseous Fluid Model Inside a Piston

We will study thermodynamically the expansion of a gaseous fluid inside a piston, where the external pressure of the piston represents the forces that tend to compress it (i.e., gravitational forces, in relation to Model 1), and the pressure inside the piston represents the forces that tend to expand it (i.e., dark energy forces, in relation to Model 1). For this thermodynamic study of the expansion, we consider the approximation of an expanding ideal gas to be sufficient.

2.4 Similarity of Magnitudes in Both Models

When studying the expansion between the two models, we must align several quantities:

The value of total energy in the Model 1 and internal energy in the Model 2, E_i , will be the same.

In Model 1, the change in potential energy during the expansion process corresponds in Model 2 to the work done by the gas inside the piston during the expansion process.

2.5 Derivation of the Zero Active Mass Equation

Taking into account the similarities made between the two models and the calculations to be performed, we consider the consideration of an ideal gas as a gaseous fluid inside the piston to be valid.

Given the ideal gas law:

$$PV = (n N_A) k_B T$$

where:

P is the pressure of the gas.

V is the volume of the gas.

n is the number of moles.

N_A is Avogadro’s number.

k_B is the Boltzmann constant.

T is the absolute temperature of the gas.

In an ideal gas, the internal energy, E_i , is the same as its kinetic energy K , given by the equation:

$$E_i = K = (3/2) (n N_A) k_B T$$

Substituting into the ideal gas law equation, we have:

$$PV = (2/3) E_i = (2/3) K$$

Differentiating this equation, we get:

$$PdV + VdP = (2/3) dK \quad (1)$$

PdV is the work done by the gas during the adiabatic expansion process inside the piston, Model 2. Based on the similarity we have established between the two models, this work against the piston walls during expansion corresponds in Model 1 to the work done by expansion forces against gravitational forces during the expansion process; that is, the potential energy with the opposite sign involved in Model 1 during the expansion process. Thus:

$$-dU = PdV$$

Substituting in (1):

$$-dU + VdP = (2/3) dK$$

According to Model 1, and in a universe with zero spatial curvature, in the process of expansion of the universe it is true that:

$$0 = dK + dU$$

$$dK = -dU$$

Thus:

$$dK + VdP = (2/3) dK$$

$$VdP = (-1/3) dK$$

$$dP = (-1/3) dK/V$$

Since the energy in ideal gases is solely a function of temperature and the kinetic energy, K , coincides with its internal energy E_i , which we have in turn made coincide with the total energy, it follows that:

$$\int dP = P = \int \left(-\frac{1}{3}\right) dK/V =$$

$$= \left(-\frac{1}{3}\right) \int dE_i/V = (-1/3) \rho$$

$$p = (-1/3) \rho$$

$$(\rho + 3p) = 0$$

Substituting into our Friedman's second equation:

$$a'' = 0$$

3. Discussion

We have obtained a proof of the zero active mass equation based on cosmological and thermodynamic considerations. To do this, we have conducted a comparative study of two models that partially represent the expansion of the universe, interrelating them. The first, Model 1, is an energy model in which the energies involved—kinetic and potential—are analyzed, and in which we have taken into account a new equation that relates them, an equation obtained by Fulvio Melia and published in a work referenced here. (Melia F, 2022) The second model, Model 2, is a simple thermodynamic model, corresponding to the expansion of an ideal gas in a piston. In this model,

the thermodynamic variables, pressure, volume, and temperature are analyzed through the ideal gas law and the work done by the gas in its expansion is related to the potential energy involved in the expansion process of the universe according to Model 1. The application of the equation found by Fulvio Melia in his work relating kinetic and potential energy in the expansion for universes of zero spatial curvature leads immediately to the equation of zero active mass, as we wanted to demonstrate.

4. Conclusion

Taking a different approach than Fulvio Melia's (Melia, F, 2026) to demonstrate the equation, we have obtained a new proof of the zero active mass equation for universes with zero space curvature, $k = 0$. Using reasoning based on cosmology and thermodynamics, we studied the expansion of the universe using two different energy models: one from classical mechanics and the other from thermodynamics. By interrelating these two models, we conclude that, according to the FLRW metric and for zero space curvature, $k = 0$, the zero active mass equation holds. It is worth noting that we used a Lapse function $g_{00} = 1$. Furthermore, during the proof, we used a result from a study by Fulvio Melia (Melia, F, 2022) which states that the space curvature k is proportional to the mechanical energy (kinetic + potential) involved in the expansion. Therefore, zero curvature implies zero mechanical energy, leading us to conclude that, for a universe in which the zero active mass equation holds, no energy input is required for its expansion other than that provided by the cosmic fluid pressure. This means that, in FLRW universes with zero spatial curvature and a Lapse function equal to one, the zero active mass equation holds, and no additional energy is needed to maintain the expansion beyond that provided by the cosmic fluid pressure. In other words, the expansion is always non-accelerated.

Recently (Junhjuk Son et al. 2025), after decades of opinions favoring an accelerating universe, it seems to be confirmed that our universe is not accelerating. To this end, the supernovas data that led to the conclusion that the universe was accelerating have been re-analyzed, along with new discoveries regarding distance and luminosity estimation based on the ages of host galaxies. This seems to confirm that the universe is in a non-accelerating expansion. This would bring the conclusions of our work closer to the most recent experimental data

Data Availability

No new data were generated or analyzed in support of this research.

Conflict of Interest

The author states that there is no conflict of interest

5. References

1. Junhjuk Son, Young-Wook Lee, Chul Chung, Seunghyun Park, Hyejeon Cho, 2025, Strong progenitor age bias in supernova cosmology – II. Alignment with DESI BAO and signs of a non-accelerating universe, Monthly notices of the Astronomical Society, Vol 544, Issue 1, November
2. Melia, Fulvio, 2021, The Cosmic Spacetime. 1th ed. CRC Press. London, New York, pp 153-230.
3. Melia, Fulvio, 2022, Initial energy of a spatially flat universe: A hint of its possible origin, Astronomische Nachrichten, Vol 343, (3).
4. Melia, Fulvio, 2026. The Physics of the Cosmology. Academic Press, Elsevier. London, pp 63-65.
5. Planck 2018 VI. Cosmological parameters. 2021 Astronomy & Astrophysics, August 10.