

A Rational Approach to the Clinical Diagnostics of Neuroleptic Cardiomyopathy

Volkov V. P.

Tver center of judicial examinations, Russia.

patowolf@yandex.ru

**Corresponding Author:* Volkov V. P., Tver center of judicial examinations, Russia.

Abstract

Introduction: The presented review is devoted to generalization of results of own researches of the author on a problem of almost not studied pathology – a neuroleptic (antipsychotic) cardiomyopathy (NCMP).

NCMP belongs to secondary specific metabolic dilated cardiomyopathies. It is caused by side cardiotoxic effect of antipsychotic preparations. Many parties of epidemiology, pathogenesis, morphology, clinic and diagnostics of NCMP remain still insufficiently developed.

Results: As showed our researches the development of NCMP passes 3 stages: 1) latent, 2) developed and 3) terminal. Each stage has the clinical features, electrocardiograph signs and certain morphology.

The lethal outcome in latent and in developed stages either comes from the intercurrent diseases or is the sudden cardiac death of arrhythmogenic genesis. The progressing congestive heart failure serves in a terminal stage as an immediate cause of death.

Apparently from our researches morphologically NCMP is characterized by a moderate hypertrophy of heart, expansion of its ventricles and absence of the expressed coronary atherosclerosis.

At microscopic research of a myocardium at early stages microcirculation violations come to light. Damages of an extracellular matrix (interstitial oedema and myofibrosis) develop then. Dystrophic-degenerative and atrophic changes of cardiomyocytes act into the forefront in a terminal stage. It leads to contractile myocardial dysfunction.

Conclusion: On the basis of synthesis of the data obtained in a number of the own researches clinical and morphological criteria of diagnostics of NCMP are allocated and proved. Considering the described features of NCMP the given pathology is allocated in separate independent nosological unit expediently and quite logical.

Keywords: antipsychotics, cardiotoxicity, clinic, diagnostics, morphology, neuroleptic cardiomyopathy, nosological independence.

INTRODUCTION

Neuroleptic (antipsychotic) cardiomyopathy (NCMP) is one of serious complications of psychotropic therapy caused by side cardiotoxic effect of antipsychotic preparations [1, 2, 3, 4].

NCMP belongs to secondary specific metabolic dilated cardiomyopathies [5, 6, 7]. The disease is characterized by diffusion damage of a myocardium,

sharp decrease in its contractile function and as a result the progressing congestive heart failure (CHF) [2, 5, 6, 8-10].

Many parties of epidemiology, pathogenesis, morphology, clinic and diagnostics of NCMP remain still insufficiently developed.

The goal of the real review is the short generalization of the results of the own researches of this problem.

MATERIAL AND METHODS

Our long-term researches are conducted on different material and various methods including statistical (parametric and nonparametric). At macroscopic studying of heart original own organometric approach is applied [11]. The microscopic condition of a myocardium is estimated by means of a morphometric method.

RESULTS

According to our section data among the died patients with schizophrenia which obligatory component of therapy is reception of antipsychotic preparations the number of the suffering of NCMP (11.4 %) is considerable and statistically significant above than in gross section material on a psychiatric pathological department in general (2.0 %) [8].

As showed our researches the development of NCMP passes 3 stages: 1) latent, it is clinically completely compensated, 2) developed at which cardiac violations distinctly are defined, but without the expressed signs of CHF, and 3) terminal when the clinic of CHF acts into the forefront [9, 12-14].

The lethal outcome in latent and in developed stages either comes from the intercurrent diseases or is the sudden cardiac death (SCD) of arrhythmogenic genesis [5, 12, 14]. The last is observed at 44.2 % of the dead of NCMP according to our data [15, 16]. The progressing CHF [5, 12, 14] serves in a terminal stage as an immediate cause of death.

The disease develops slowly and at the beginning is hardly noticeable. In a latent stage it is practically shown by nothing.

During this period the complaints of patients have uncertain character or are in general absent. Fatigue and short wind at considerable physical activity is most often noted. Thus it must be kept in mind known difficulties of detection of complaints at mental patients connected both with their inadequate behavior and lack of due criticism to their state and with quite often certain medicament load.

In a latent stage of NCMP the findings of the examination are not numerous and aren't specific. The tachycardia serving as almost constant phenomenon at reception of neuroleptics [1] is observed as a rule. De an auscultation in is defined the deafness of cardiac sounds. Poorly the borders of heart are usually changed.

An arterial hypo- and a norm tonicity significantly prevail from the arterial pressure. The insignificant arterial hypertension is only approximately in $\frac{1}{3}$ cases observed [5, 12].

On the electrocardiogram during this period there are most often the following pathological signs: 1) diffusion muscular changes; 2) different types of violation of conductivity, in particular the blockade of the left leg of Gis's bunch; 3) deviation of an electric axis of a heart to the left; 4) overload of the right departments of the heart; 5) hypertrophy of the left ventricle [5, 6, 8-10, 17].

In the developed stage the clinic of NCMP is rather distinctly shown, but the signs of terminal CHF are absent or poorly noticeable. The complaints of patients are more certain: weakness, fatigue, heartbeating, short wind at moderate physical activity, sometimes passing pains in the cardiac region. By the physical examination the deafness of cardiac sounds, some expansion of the borders of a heart, tachycardia, passing breathlessness notes. During this period the steadily normal or labile arterial pressure equally often meets, but there is a tendency to the moderate increase of it [5, 12].

The terminal stage of a current of NCMP is characterized by accession to already listed symptoms of the known manifestations of the increasing CHF: short wind at a rest or the small physical activity, orthopnea posture, the increase of a liver, the peripheral and cavitary oedemata, sometimes an anasarka etc. The heart borders are expanded that is confirmed by the roentgen exploration. The cardiac sounds are deaf. There are almost always a tachycardia and an arrhythmia. The moderate arterial hypertension is observed by a little more than at $\frac{1}{4}$ patients, and the arterial pressure constantly exceeds 150/100 mm of mercury at 13.6 % of patients [5, 12].

On an electrocardiogram in developed and terminal stages of NCMP there are such most dangerous phenomena: 1) violations of conductivity; 2) lengthening of an interval of QT in recalculation on Bazett's formula - a correct QT interval (QTc); 3) overload of the right departments of the heart [5, 6, 8-10, 17]. The special attention is deserved by monitoring of parameters of QTc as highly informative indicator in the conditions of a decompensation of the heart [18].

DISCUSSION

The high frequency of NCMP revealed on autopsy at the patients accepting antipsychotics testifies to what broad application in medical practice of neuroleptics is the reason of development of the given cardiac pathology in patients with schizophrenia.

It is seen from our researches clinical diagnostics of NCMP is difficult as its symptom complex has no peculiar features [19].

On the basis of synthesis of the data obtained in a number of the own researches [2, 6, 8, 10] the clinical criteria of diagnostics of NCMP are allocated and proved [5, 19, 20]. The diagnostic scheme is approximately submitted as follows.

I. Big criteria (absolute)

- [1] The treatment by neuroleptics;
- [2] The absence of other cardiac pathology.

II. Small criteria (relative)

- [1] The cardiac complaints (even minimum);
- [2] The increase in the sizes of a heart (be percussion and the roentgen exploration);
- [3] The findings by the physical examination (tachycardia, arrhythmia, deafness of cardiac sounds);
- [4] The changes of an electrocardiogram (especially the increase in QTS, violations of a rhythm and a conductivity);
- [5] The existence of CHF in the absence of other reasons of its development;
- [6] The SCD of patients (it is important for posthumous diagnostics).

It is necessary to emphasize that each of the listed signs taken separately out of communication with others isn't specific to NCMP. However in the set these criteria can become a reliable basis for clinical verification of this disease. The long reception of antipsychotic preparations is the main condition forcing to think of NCMP.

There is basis to consider that for diagnostics of NCMP it is enough to have available both big signs and at least two small. However in this direction nevertheless further researches and accumulation of practical experience are necessary.

At the same time in our opinion the offered criteria of diagnostics of NCMP and the approximate diagnostic scheme will be able to become useful for the early identification of this serious complication of an antipsychotic therapy in order to its timely corrective treatment.

CONCLUSION

Proceeding from told about NCMP this pathology is allocated in separate independent nosological unit expediently and quite logical. On ICD-10 it can be carried to the heading I42.7 "The cardiomyopathy caused by influence of medicines and other external factors" with the additional code reflecting an external cause of illness (a class of antipsychotic preparations) – Y49.3-Y49.5 [5, 17].

REFERENCES

- [1] Volkov VP. Cardiotoxicity of phenothiazine neuroleptics (review of literature). *Psichiat psychopharmacother* 2010; 12(2): 41-5.
- [2] Volkov VP. Phenothiazine dilated cardiomyopathy: some aspects of clinic and morphology. *Clin Med* 2009; 8: 13-6.
- [3] Buckley NA, Sanders P. Cardiovascular adverse effects of antipsychotic drugs. *Drug Saf* 2000; 23: 215-28.
- [4] Coulter DM, Bate A, Meyboom RHB, Lindquist M, Edwards IR. Antipsychotic drugs and heart muscle disorder in international pharmacovigilance: data mining study. *Br Med J* 2001; 322: 1207-9.
- [5] Volkov VP. Clinical characteristic of an antipsychotic cardiomyopathy. In: Volkov VP, editor. Actual problems of therapeutic clinic: collective scientific monograph. Novosibirsk: SibAC 2013; p. 94-116.
- [6] Volkov VP. Electrocardiographic manifestations of an antipsychotic cardiomyopathy at patients with schizophrenia at stages of its morphogenesis. *Upper Volga Med J* 2012; 10(1): 13-6.
- [7] Tereshchenko SN, Jaiani NA. Dilated cardiomyopathy today. *Heart failure* 2001; 3(2): 58-60.
- [8] Volkov VP. To a question of a secondary phenothiazine cardiomyopathy. *Clin Med* 2011; 5: 30-3.

A Rational Approach to the Clinical Diagnostics of Neuroleptic Cardiomyopathy

- [9] Volkov VP. Features of an electrocardiogram at a phenothiazine cardiomyopathy. Upper Volga Med J 2009; 7(4): 3-7.
- [10] Volkov VP. Features of the electrocardiogram at a phenothiazine cardiomyopathy. Clin Med 2011; 4: 27-30.
- [11] Volkov V.P. New method of an organometry of heart. In: Volkov VP, editor. Paraclinical disciplines: new methods and diagnostic opportunities": collective monograph. Novosibirsk: SibAC 2014; p. 78-100.
- [12] Volkov VP. Dynamics of arterial pressure at an antipsychotic cardiomyopathy. In: Innovations in science: materials of the XIXth international correspondence scientific and practical conference (on April 22, 2013). Novosibirsk: SibAC 2013; p. 130-7.
- [13] Volkov VP. Morphological features of an antipsychotic cardiomyopathy. In: Medicine: calls of today: international correspondence scientific conference (Chelyabinsk, June, 2012). Chelyabinsk 2012; p. 33-6.
- [14] Volkov VP. Morphometric aspects of a morphogenesis of an antipsychotic cardiomyopathy. Ros J Cardiol 2012; 3: 68-73.
- [15] Volkov VP. Sudden death of patients with schizophrenia. Upper Volga Med J 2009; 7(2): 3-7.
- [16] Volkov VP. Sudden cardiac death at schizophrenia. Mental health 2013; 1: 50-4.
- [17] Volkov VP. Antipsychotic cardiomyopathy. In: Volkov VP, Zacharov RI, editors. Pharmacotherapy: new prospects and problems: collective scientific monograph. Novosibirsk: Siberian Association of Consultants 2012; p. 65-84.
- [18] De Luna AB. The guidance on a clinical electrocardiogram. M.: Medicine 1993.
- [19] Volkov VP. Antipsychotic cardiomyopathy: diagnostic approaches and criteria of the diagnosis. Doctor 2013; 9: 69-71.
- [20] Volkov VP. Antipsychotic cardiomyopathy: clinical-morphological criteria of the diagnosis. Curing doctor 2013; 10: 77-80.

Citation: Volkov V. P. *A Rational Approach to the Clinical Diagnostics of Neuroleptic Cardiomyopathy*. Archives of Cardiology and Cardiovascular Diseases. 2018; 1(1): 49-52.

Copyright: © 2018 Volkov V. P. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.