

RESEARCH ARTICLE

Frequency of Hepatitis B, C and HIV Infections among Transfusion-Dependent Beta Thalassemia Patients in Bangladesh Shishu Hospital and Institute

Md. Abdul Wohab¹, Nilufar Akter Banu Choudhury², Umme Nusrat Ara³

¹Associate Professor, Department of Paediatric Hematology & Oncology, BSH&I, Dhaka, Bangladesh

²Assistant Professor, Department of Paediatric Hematology & Oncology, BSH&I, Dhaka, Bangladesh

³Assistant Professor, Department of Paediatric Hematology & Oncology, BSH&I, Dhaka, Bangladesh

Received: 18 April 2025 Accepted: 02 May 2025 Published: 08 May 2025

Corresponding Author: Md. Abdul Wohab, Associate Professor, Department of Paediatric Hematology & Oncology, BSH&I, Dhaka, Bangladesh.

Abstract

Transfusion transmitted infections have remained a major deterrent to public health, particularly among the patients with transfusion-dependent Beta thalassemia in developing countries. Although proper donor selection through adoption of WHO-advised infection panel has lowered the rate of infections, the multi-transfused patients are not free of risk. In this study, we screened 200 transfusion-dependent Beta thalassemia patients to determine the frequency of Hepatitis C Virus (HCV), Hepatitis B Virus (HBV) and Human Immunodeficiency Virus (HIV) using the ELISA method. Among them, infected cases with HCV, HBV and HIV were 8.5%, 1.5% and 0%, respectively. Moreover, 2% of the patients were found to be co-infected with both HBV and HCV. Immunochromatography (ICT)-based rapid test kits are usually used to screen and confirm these infections in the blood of the patients. However, ICT-based tests are not sensitive enough to detect the infections. So, a combination of both Nucleic Acid testing (NAT) and serological testing are suggested to significantly reduce the risk of viral infections during blood transfusion.

Keywords: Thalassemia, Transfusion Transmitted Infection, Immunochromatographic (ICT) Testing, HBV, HCV.

1. Introduction

Beta-thalassemia is a group of inherited blood disorders characterized by reduced or no synthesis of the beta-globin chain, and if the synthesis of both genes is reduced or absent, the person has beta-thalassemia major [1,2]. Patients with beta-thalassemia major are at risk for blood-borne viral infections such as Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Human Immunodeficiency Virus (HIV) due to frequent blood transfusions [3]. Frequent blood transfusions increase the life expectancy of these patients but are associated with an increased risk of

blood-borne diseases. Prevention of viral infections is an important goal in the treatment of patients with beta-thalassemia major [4]. According to reports, before 1990, in Western countries and before 1995 in Iran, donated blood was not screened in blood transfusion organizations for hepatitis C virus. Therefore, patients in beta-thalassemia major were more likely to become infected with viral infections due to the need for frequent blood transfusions [3]. Screening donors with serological tests and through interviews and medical examinations has reduced the risk of post-transfusion infections, especially hepatitis C. Today,

Citation: Md. Abdul Wohab, Nilufar Akter Banu Choudhury, Umme Nusrat Ara. Frequency of Hepatitis B, C and HIV Infections among Transfusion-Dependent Beta Thalassemia Patients in Bangladesh Shishu Hospital and Institute. Archives of Pediatrics and Neonatology 2025;8(1): 01-05

©The Author(s) 2025. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

due to the increasing use of hepatitis B vaccine and screening tests for all donated blood for hepatitis B, the prevalence of HBV in patients with beta-thalassemia major has decreased significantly [5]. These inherited blood disorders generally occur in the countries of thalassemia belt including Mediterranean and portions of West Africa, North Africa, Middle East and South Asian countries including Bangladesh, India, and Sri Lanka [6]. As the patients with thalassemia, especially the patients with beta thalassemia major (BTM) and a group of patients with Hemoglobin E-beta thalassemia (EBT) are transfusion-dependent and these patients are very much prone to transfusion-transmitted viral infections. After heart failure, viral infections are the second most common cause of mortality and the foremost cause of morbidity among the patients with thalassemia followed by patients with bacterial and parasitic infections [7]. Although hepatitis B virus (HBV), hepatitis C virus (HCV), human immunodeficiency virus (HIV), West Nile virus (WNV), human T cell lymphotropic viruses I, II (HTLV-I/II) had been reported most frequently, HCV and HBV had been known as the most prevalent etiological agents of chronic viral hepatitis and hepatocellular carcinoma among the thalassemia patients [8]. In the developed countries, the TTIs risks have been virtually eliminated because of the careful and efficient screening approaches [9]. However, challenges in developing countries span the entire blood safety chain from donor selection to post-transfusion surveillance and these risks are really a burning question for public health development [10]. Bangladesh lies in the global thalassemia belt with around 9176 new thalassaemic patients each year [11]. The patients with transfusion dependent BTM and EBT undergo regular transfusions with varying degrees of frequencies of transfusions in every 7 to 120 days. However, although it is mandatory, World Health Organization (WHO)–recommended standards for donor blood screening are not always followed in developing countries like Bangladesh due to resource limitations and occasional lack of consciousness [12]. The patients with transfusion dependent BTM and EBT in Bangladesh not only suffer from various complications for the disease itself, but also due to the lack of proper screening methods of donor blood, and thus transfusion-transmitted infections are threatening as ‘Silent killers’ [13]. The minimization of the risks of TTIs highly depends on selection of safe blood donors through screening of the blood using sensitive methods for pathogen detection and such precaution reduces the risk of allogenic blood transfusion. However, transmission of diseases still occurs if there

is the inability of the test method to detect the disease in the ‘pre-seroconversion’ or ‘window’ phase of their infections, immunologically variant viruses, non-seroconverting chronic or immune–silent carriers and laboratory testing errors [14]. In the present study, the patients with thalassemia used to visit Bangladesh Shishu Hospital Thalassemia centre for routine blood transfusion. In the hospital, immunochromatographic tastings (ICTs) were usually performed to detect the presence of a number of common infections in the blood of donors before every transfusion. However, these ICT-based rapid diagnostic test (RDT) kits have not been validated using confirmatory approaches, such as ELISA or molecular methods and usually the ICT-based methods have repeatedly demonstrated low sensitivity and specificity to detect the major TTIs [14,15]. The present study investigated the frequency of HBV, HCV and HIV infections among transfusion-dependent beta thalassemia patients of the Bangladesh Shishu Hospital Thalassemia centre using ELISA method with the aim to explore the efficacy of RDT based screening method.

2. Materials and Methods

2.1. Study Population

This prospective observational study was done on 200 patients with transfusion-dependent Beta thalassemia; they were enrolled over a period of 24 months from January 2023 to December 2024. These patients were in the age range of 1–18 years. They came to visit the Bangladesh Shishu Hospital Thalassemia centre for follow-up examination and blood transfusion. Both EBT and BTM patients participated in this study and they were all differentially diagnosed by Hemoglobin capillary electrophoresis. The study also included 30 healthy controls (aged 1–18 years) who did not have any previous history of hemoglobinopathies or related blood disorders. Prior to sample collection, a written informed consent was obtained from the adults as well as the legal guardians of the children (both for patients and controls). About 5.0 mL of blood were collected from each participant in serum tube using the standard venipuncture method and serum was separated from the collected blood. In case of patients, blood was collected before transfusion. The study was approved by the National Ethics Review Committee (NERC) of Bangladesh Shishu Hospital & Institute, Dhaka, Bangladesh.

2.2. Donor Selection

Before transfusion, some regular immunochromatographic testings (ICTs) were done to detect the presence of any infection in the donors' blood like HBV, HCV, HIV, Syphilis and Malaria.

2.3. Viral Infection Detection

As the presence of Hepatitis B surface antigen (HBsAg) in the serum is a good indicator of HBV infection, Bioelisa HBsAg 3.0 kits (Biokit, Barcelona, Spain) were used to detect HBsAg infections. Bioelisa HCV 4.0 (Biokit, Barcelona, Spain) and HIV 1/2/O Antibody EIA test kits (Acon Laboratories, CA, USA) were used for the detection of HCV and HIV, respectively. The tests were performed in accordance with the manufacturer's instruction provided with the kits. The color intensity of the final reaction mixture was measured with an ELISA reader (LabSystems Multiskan, Texas, United States) at 450 nm.

2.4. Statistical Analysis

We performed two-tailed chi-square test (two by two table) using <http://www.openepi.com/> to calculate the frequency of infections among the two transfusion interval groups (≤ 30 days and > 30 days). Estimation of the relative risk ratio (RR) and odds ratio (OR) were done at 95% confidence interval (CI). A P-value of less than 0.05 was considered significant.

3. Results

3.1. Demographic Characteristics of the Patients

In this study, 55% of the participants were male and 45% were female including both BTM and EBT patients. We recorded different demographic data of the patients, such as name, address, gender, height, weight, and BMI as well as transfusion related information like age of first transfusion, transfusion interval and number, splenectomy etc. The baseline information of the patients is given in Table 1.

Table 1. Baseline information of the patients enrolled in this study.

Categories	Patient
Gender	Male, N = 110 (55%); Female, N = 90 (45%)
Thalassemia type	EBT, N = 110 (55%); BTM, N = 90 (45%)
Age (years), mean \pm SD	5 \pm 2.21
BMI (kg/m ²), mean \pm SD	15.22 \pm 2.32
Transfusion interval (days), mean \pm SD	30.12 \pm 7.33
Transfusion number, mean \pm SD	45.12 \pm 12.44
Splenomegaly (cm), mean \pm SD	5.22 \pm 2.45
Splenectomy	N = 25 (12.5%)

(Abbreviations: EBT = E Beta Thalassemia, BTM = Beta Thalassemia Major, SD = Standard Deviation, BMI = Body Mass Index).

3.2. Trend of Infections among the Patients

A total of 25 patients (12.50%) with thalassemia were found to be infected (Male: N = 15, 60% and female: 10, 40%) in which we identified 20 (10%) patients infected with either HBV or HCV and 5 (2.5%) of

them were co-infected with both HBV and HCV., Moreover, the number of patients infected with HCV (N = 17, 8.5%) was higher than that of HBV (N = 3, 1.5%) (Table 2). Furthermore, none of the patients was infected with HIV.

Table 2. Frequency of viral mono-infection and co-infection among the transfusion-dependent E-beta thalassemia (EBT) and Beta thalassemia major (BTM) patients.

Thalassemia Type	Infection Type				No. of Total Infected (%)
	No. of Patients with HBV and HCV (%)	No. of Patients with HBV (%)	No. of Patients with HCV (%)	No. of Patients with HIV (%)	
EBT (N = 110)	2 (1)	2 (1)	10 (5)	0 (0)	14 (7)
BTM (N = 90)	3 (1.5)	1 (0.5)	7 (3.5)	0 (0)	11 (5.5)
Total (N = 200)	5 (2.5)	3 (1.5)	17 (8.5)	0 (0)	25 (12.5)

(Abbreviations: EBT = E Beta Thalassemia, BTM = Beta Thalassemia Major, HBV = Hepatitis B Virus, HCV = Hepatitis C Virus, HIV = Human Immunodeficiency Virus).

4. Discussion

In the Bangladesh Shishu Hospital Thalassemia centre, effective screening processes for infections are done before the blood transfusion for the proper selection of donors. However, we still could identify patients with either HBV or HCV or both. In the case of HBV, there were some asymptomatic donors who were in the 'window period' (i.e., the early infection period when an immunologic test is non-reactive) without any expression of HBsAg. Also, people with occult HBV (absence of HBsAg and the presence of HBV DNA in liver tissues with or without HBV DNA in the serum) may remain undetected by traditional ICT kits [16]. A previous study also showed that the sensitivity of ICT-based rapid tests was not high enough to detect the hepatitis status of a donor and these kits might not detect certain prevalent serotypes of HBV in any particular region [8]. Detection of the HBV DNA by implementation of nucleic acid testing (NAT) assays had been reported to shorten the core window period to reduce the residual risk [17]. In this study, the number of patients infected with HCV was higher than that of HBV. It should be noted here that although there was an HBV vaccine coverage to some extent in Bangladesh, there is still no HCV vaccine available yet [18]. This may explain why there were more thalassemia patients infected with HCV than with HBV. The rate of infection was found to be significantly higher in the patients who had to undergo frequent transfusions which indicates that multi-transfused thalassemia patients are more prone to infections. This is compatible with previous findings which indicate that receiving multiple transfusions for a long period of time might cause immunomodulation resulting in susceptibility to infections [9]. In the Bangladesh Shishu Hospital Thalassemia centre, sometimes ELISA methods are applied as a confirmatory test after the donors are suspected as positives by ICT approach. Nevertheless, a previous study had shown that ELISA only could detect HBsAg, whereas real-time PCR represents the infection status by detecting HBV-DNA [10]. On the other hand, recombinant immunoblot assay (RIBA) has a greater accuracy in detecting HCV [18]. The multi-transfused patients with thalassemia may face increased immune dysfunction in the presence of iron overload following splenectomy, which makes them more susceptible to infections [19]. Moreover, in case of emergency, there is a common practice in Bangladesh of risky blood donations by professional donors without any kind of testing who are mostly

drug abusers [20]. Although, nowadays, an effective awareness of safe blood transfusions and iron chelation therapy have made the morbidity rate of patients with transfusion-dependent Beta thalassemia lower, new complications like hepatocellular carcinoma are taking hold in the patients with thalassemia, which may be due to carcinogenicity of iron overload and chronic infections [21]. Specially, the thalassemia patients who become co-infected with both HBV and HCV are at a greater risk of cirrhosis and hepatocellular carcinoma compared to the mono-infected patients [22]. Thus, there is no alternative to making the blood transfusion process safer by proper selection of voluntary healthy blood donors, along with NAT which reveals viral agents earlier in the 'window period' than other immunoassays. Additionally, screening of HBV-DNA and HCV-RNA needs to be practiced regularly to avoid any kind of risks during blood transfusions [19]. Studies in India, China and Saudi Arabia had reported that NAT-based detection methods for HCV and HBV could tremendously improve the efficacy of screening for protecting blood recipient from TTIs [23,24]. Furthermore, introduction of HBV NAT in the USA, along with the HBV vaccination policy made a substantial contribution to blood transfusion safety and decreased residual risk of HBV infections [25]. In the UK, NAT has reduced the risk of HCV by 95% and that of HIV by 10% [26]. However, in the case of very low levels of viremia, NAT might not be effective enough to detect the infections. Despite these limitations, to ensure safe blood transfusion, combination of both NAT and serological testing may significantly reduce the risk of viral infections during transfusion and thus may improve the quality of life of the thalassemia patients [8].

5. Conclusion

To ensure the frequent transfusion patients a good quality of life from these preventable HBV, HCV and HIV complications, efficient screening methods for proper donor selection should be adopted in Bangladesh. In addition, awareness should be built to ascertain a safe transfusion.

6. References

1. Galanello R., Origa R. Beta-thalassemia. *Orphanet Journal of Rare Diseases*. 2010;5(1):12.
2. Muncie H.L., Jr., Campbell J.S. Alpha and beta thalassemia. *American Family Physician*. 2009;80(4):339–344.
3. Memon M., Memon M. Hepatitis C: an epidemiological

- review. *Journal of Viral Hepatitis*. 2002;9(2):84–100. doi: 10.1046/j.1365-2893.2002.00329.x.
4. Ardalan F.A., Osquei M.R., Toosi M.N., Irvanloo G. Synergic effect of chronic hepatitis C infection and beta thalassemia major with marked hepatic iron overload on liver fibrosis: a retrospective cross-sectional study. *BMC Gastroenterology*. 2004;4(1):17. doi: 10.1186/1471-230X-4-17.
 5. André F. Hepatitis B epidemiology in asia, the Middle East and africa. *Vaccine*. 2000;18: S20–S22.
 6. Tahura S. Thalassaemia and other hemoglobinopathies in Bangladeshi children. *Imp. J. Interdiscip. Res*. 2017; 3:180–184.
 7. Kiani R.A., Anwar M., Waheed U., Asad M.J., Abbasi S., Zaheer H.A. Epidemiology of Transfusion Transmitted Infection among Patients with β -Thalassaemia Major in Pakistan. *J. Blood Transfus*. 2016; 2016:1–5.
 8. Vento S., Cainelli F., Cesario F. Infections and thalassaemia. *Lancet Infect. Dis*. 2006; 6:226–233.
 9. Blanco S., Balangero M.C., Valle M.C., Montini O.L., Carrizo L.H., Gallego S. Usefulness of nucleic acid testing to reduce risk of hepatitis B virus transfusion-transmitted infection in Argentina: High rate of recent infections. *Transfusion*. 2016; 57:816–822.
 10. Pruett C.R., Vermeulen M., Zacharias P., Ingram C., Tagny C.T., Bloch E.M. The Use of Rapid Diagnostic Tests for Transfusion Infectious Screening in Africa: A Literature Review. *Transfus. Med. Rev*. 2015; 29:35–44.
 11. Noor F.A., Sultana N., Bhuyan G.S., Islam T., Hossain M., Sarker S.K., Islam K., Khan W.A., Rahman M., Qadri S.K., et al. Nationwide carrier detection and molecular characterization of β -thalassemia and hemoglobin E variants in Bangladeshi population. *Orphanet J. Rare Dis*. 2020; 15:15.
 12. Awan S.A., Junaid A., Sheikh S. Transfusion Transmissible Infections: Maximizing Donor Surveillance. *Cureus*. 2018;10: e3787.
 13. Saeed M., Hussain S., Rasheed F., Ahmad M., Arif M., Rahmani M.T.H. Silent killers: Transfusion Transmissible Infections-TTI, among asymptomatic population of Pakistan. *J. Pak. Med. Assoc*. 2017; 67:369–374.
 14. Agarwal N., Chatterjee K., Coshic P., Borgohain M. Nucleic acid testing for blood banks: An experience from a tertiary care centre in New Delhi, India. *Transfus. Apher. Sci*. 2013; 49:482–484.
 15. Bloch E.M., Shah A., Kaidarova Z., Laperche S., Lefrere J.-J., Van Hasselt J., Zacharias P., Murphy E.L. Anglophone Africa Transfusion Research Group the Anglophone Africa Transfusion Research Group A pilot external quality assurance study of transfusion screening for HIV, HCV and HBsAG in 12 African countries. *Vox Sang*. 2014; 107:333–342.
 16. Laperche S. Francophone African Group for Research in Blood Transfusion Multinational assessment of blood-borne virus testing and transfusion safety on the African continent. *Transfusion*. 2013; 53:816–826.
 17. Keechilot C.S., Shenoy V., Kumar A., Biswas L., Vijayarajratnam S., Dinesh K., Nair P. Detection of occult hepatitis B and window period infection among blood donors by individual donation nucleic acid testing in a tertiary care center in South India. *Pathog. Glob. Health*. 2016; 110:287–2
 18. Ahmed M.U., Begum H.A., Hossain T., Chakraborty P. Incidence of Common Transfusion Transmitted Diseases among Blood Donors. *J. Armed Forces Med. Coll. Bangladesh*. 2009; 5:4–6.
 19. Choudhry V.P. Hepatitis B and C Infections in Multitransfused Thalassaemic Patients. *Indian J. Pediatr*. 2015; 82:212–214.
 20. Mollah A.H., Nahar N., Siddique A., Anwar K.S., Hassan T., Azam G. Common transfusion-transmitted infectious agents among thalassaemic children in Bangladesh. *J. Health Popul. Nutr*. 2003; 21:67–71.
 21. Moukhadder H.M., Halawi R., Cappellini M.D., Taher A.T. Hepatocellular carcinoma as an emerging morbidity in the thalassemia syndromes: A comprehensive review. *Cancer*. 2017; 123:751–758.
 22. Konstantinou D., Deutsch M. The spectrum of HBV/HCV coinfection: Epidemiology, clin characteristics, viral interactions and management. *Ann. Gastroenterol*. 2015; 28:221–228.
 23. Kumar R., Gupta S., Kaur A., Gupta M. Individual donor-nucleic acid testing for human immunodeficiency virus-1, hepatitis C virus and hepatitis B virus and its role in blood safety. *Asian J. Transfus. Sci*. 2015; 9:199–202.
 24. Dong J., Wu Y., Zhu H., Li G., Lv M., Wu D., Li X., Zhu F., Lv H. A pilot study on screening blood donors with individual-donation nucleic acid testing in China. *Blood Transfus*. 2013; 12:172–179.
 25. Stramer S., Notari E.P., Krysztof D.E., Dodd R.Y. Hepatitis B virus testing by minipool nucleic acid testing: Does it improve blood safety? *Transfusion*. 2013; 53:2449–2458.
 26. Soldan K., Barbara J.A.J., Ramsay M., Hall A.J. Estimation of the risk of hepatitis B virus, hepatitis C virus and human immunodeficiency virus infectious donations entering the blood supply in England, 1993-2001. *Vox Sang*. 2003; 84:274–286.