

Knowledge of Breast Self-Examination Among LHV's and Female Nursing Students of KP

Aurang Zeb^{1*}, Humera Naz², Nadim Khan³, Tariq Aziz⁴, Arifa Siddique⁵

¹Assistant Professor Rehman College of Nursing Peshawar

²student Rehman College of Nursing Peshawar

³student Rehman College of Nursing Peshawar

⁴student Rehman College of Nursing Peshawar

⁵student Rehman College of Nursing Peshawar

aurang.zeb@rmi.edu.pk

**Corresponding Author: Aurang Zeb, Assistant Professor Rehman College of Nursing Peshawar, Pakistan.*

Abstract

Background: BSE is cheap and non-invasive method of breast cancer detection. Breast cancer is most common cause of death in women worldwide. Early diagnosis can lead to effective treatment and improves the quality of life.

Objective: This study was performed with aim to determine the knowledge of LHV's and undergraduate female students of BSN regarding Breast Self-Examination.

Method: This is cross sectional study, was performed on 58 LHV's and undergraduates students of BSN of RCN, PHI and RTI. Data was collected by using a self-adopted questionnaire and was analyzed by SPSS software (version 22).

Results: Among 58 students, (53.4%) participants have good knowledge, 44.8% have excellent knowledge and 1.7% have poor knowledge. (51) 87.9% were practicing BSE, (2)3.4% were not practiced it and the remaining 8.60% were missing. There is no significant association between variables. There is a good breast self-examination awareness in students.

Conclusion: BSE is cheap and noninvasive tool to identify any abnormality in breast and also help to detect early sign of breast cancer. Teaching nursing students regarding BSE helps them to receive gain more knowledge and to disseminate the information in community in future.

INTRODUCTION

Breast cancer starts when cells in the breast begin to divide and go in an abnormal way. Furthermore, the term breast cancer refers to a malignant tumor that is developed from cell in the breast. Breast cancer starts when cells in the breast begin to divide and go in an abnormal way. About 1(12%) in 8 women in the United States are reported to have developed breast cancer over the entire life (1)

Breast cancer is one of the leading causes of death in Pakistan. It is about 100 times frequent among women as compared to men (2). It is the most common invasive cancer in women, and the second main cause

of cancer death in women, after lung cancer (3). Cancer is classified into various stages according to size and spread of tumor to lymph nodes or other parts of the body. The stages are briefly given below:

Stage 0: It is known as ductal carcinoma. The cells are limited to duct.

Stage 1: At this stage, tumor is up to 2cm, and it has no affected surroundings.

Stage 2: The tumor is 2cm across, and it has started affecting the nearby nodes.

Stage 3: The tumor is up to 5cm across, and it may be spread to some lymph nodes.

Stage 4: The cancer is spread to other organs especially liver, brain or lungs.

The exact cause remains unclear, but there are some risk factors which lead to breast cancer. Age, genetics, dense breast tissue, estrogen exposure, alcohol consumption, radiation exposure and hormonal therapy might be one of the causes for cancer(4).

Identification of breast cancer at an early stage is of great importance that help in decreasing morbidity and mortality rate. A 95% survival rate can be achieved when breast cancer is identified with early stage (5).

A self-breast examination is a method that is performed by a female to identify any abnormal changes in the breasts (e.g.) tumor, lump or cyst. It can be performed every month and it helps women to become familiar with the size, shape and texture of her breast.

Women who perform breast self-examination had more perceived benefits than women who do not perform it. Breast Self- examination (BSE) provides a relatively simple, low cost method of early detection that can be performed more frequently than mammography or clinical examination. Monthly BSE has been reported to be effective in detecting the early symptoms of breast cancer which, in turn, greatly reduces mortality from breast cancer (6).

Breast cancer has been considered as a major health problem in females, because of its high incidence in recent years. Due to the role of breast self-examination (BSE) in early diagnosis and prevention of morbidity and mortality rate of breast cancer, promoting student knowledge, capabilities and attitude are required in this regard. Although, American Cancer Society do not recommend SBE as a powerful screening measure for diagnosis of breast cancer (7), it still can increase the awareness about BC and alert the women and the physician about the need to perform more advanced screening measure, especially women with positive family history of breast cancer (8). Study conducted in Korea by about knowledge, attitude and practice of obstetric nurses in relation to breast cancer and self-breast examination resulted that 99% participants thought self-breast examination is important; however, only 58.9% practiced self-breast examination (9). The nurses who had the knowledge of self-breast examination were found superior in attitude and practice to those who had no knowledge.

Clarke and Savage reported that BSE can be taught by a variety of professionals including nurses, physicians, trained peer educators, researches and

graduate students without major differences. Heyman et al. found that a program of instruction improved the abilities of nurses to teach SBE to their patients (10). A study in Southeastern Iran Shows that Only 21.6% of women had good knowledge about BSE. Research on female health care workers in Tehran about the subject matter shows that 63% of the participants know about BSE and in Turkey more than half of the nursing and midwifery students had sufficient information about BSE. In Ilorin Nigeria, 95.6% of respondents were aware of BSE, 66.5% of Sudanese students know about BSE.

In a study conducted on Beliefs and Behavior of Saudi Women in the University of Tabuk towards Self-Breast Examination practice shows that only (7.8%) practiced BSE regularly each month in the past year, and (9%) is intended to perform BSE monthly in future. (11).

A cross-sectional descriptive study was carried out on "Breast Self-Examination Knowledge and Practice among Newly Diagnosed Patients of Carcinoma Breast" at Outpatient department of General surgery in Pakistan Institute of Medical Sciences, Islamabad. One hundred ninety-one patients were included in the study, Result showed significant relationship ($p < 0.05$) between knowledge and practice of BSE with age, marital status, education level, living area and socioeconomic status (12).

The nurses and LHV's have a role for health promotion and nature of the nurse client relationship that provides opportunities for health education. LHV's and Nurses students require much information as possible about women's health behaviors for development of effective health programs such as self-breast examination.

If nursing professionals were aware of importance of doing BSE then they may give information or education in Primary Health Care services. This will allow women to know the normal and abnormal changes of breasts and help in early detection of breast cancer and will decrease the mortality rate of women due to breast cancer and there will also decrease in disease burden on government.

METHODOLOGY

Study Design

Cross sectional study was conducted to assess the knowledge of BSE among LHV's and female nursing students.

Knowledge of Breast Self-Examination Among LHVs and Female Nursing Students of KP

Study Setting

The study was conducted in three academic settings of Peshawar KPK: Public Health Institute (PHI), Regional Training Institute (RTI) and Rehman College of Nursing (RCN). These institutes provide education to LHVs and nursing students.

Study Population

Study population included LHVs and female nursing students.

Sample Technique and Size

Convenient sampling method was applied to obtain target sample from selected institutes. The sample size was 60.

Inclusion and Exclusion Criteria

Only female students present in the classes during data collection and want to participate in the study were included.

Data Collection Tool

A questionnaire was adopted from study about Knowledge and beliefs of breast self-examination and

breast cancer among market women in Ibadan (13). It included questions based on knowledge of BSE. The data collection process was carried by female students of nursing during their research training course of academic phase.


Ethical Consideration

Approval for data collection was obtained from the director/ principles of selected institutes. Written consent was also obtained from students as mentioned above, anonymity and confidentiality is assured.


RESULTS

Total 58 subjects were participated in the study. Among which (15) 25.9 % were married, (38) 65.5% were unmarried while (5) 8.6% did not mention their marital status.

The age of the participants' was 19-53 years. The mean was 29.21 with S.D 11.364. According to academic education 22.40% participants were matriculated, 43.10% were intermediated and 25.90% were bachelors as shown in the paragraph below:




Regarding professional education 56.90% were LHV students and 34.50% were undergraduate students of BSN.



In motherhood (14) 42.1% have their children and (39)67.2% have no children.

Knowledge of Breast Self-Examination Among LHV's and Female Nursing Students of KP

Among all the participants, 87.9% were performing BSE, 3.4% were not performing. The participants were categorized into poor, good and excellent categories based on their knowledge regarding SBE.



Those who have their children are 24.10% and those who do not have children is 67.20%. The mean score in the knowledge part of the questionnaire is 1.701 ± 10.02 .

ANOVA, T test and regression were applied to identify any significant association between variables but no significant association was noted.

DISCUSSION

The study was aimed to identify the knowledge of BSE among LHV's and female nursing students. The knowledge score of this study shows that (53.4%) participants have good knowledge, 44.8% have excellent knowledge and 1.7% have poor knowledge. A study in Nigeria shows that 45.7% of the population have good knowledge (13). Study on female students in Erbil Medical Technical Institute in Kurdistan Region -Iraq found that only 4.3% of the participants had a good level of knowledge, and only 6% had a good level of practicing BSE (14).

The practice of Self Breast examination in this study shows that (51) 87.9% were performing BSE and those who were not performing BSE were 3.4%. and the remaining 8.60% were missing and most of the participants reported that they had heard about BSE, and nearly half of them have good knowledge about BSE. This result is similar to another study conducted in the University of Tabuk at Saudi Arabia where (95.5%) reported that they have heard about BSE from different sources like TV, radio and newspaper etc (15). A research carried out among female undergraduate students in a higher teachers training college in Cameroon regarding awareness of breast cancer and breast self-examination shows that 47% of the subjects who knew about breast cancer had

heard about breast self-examination (BSE). Majority ($n = 47$, 32.9%) did not know how often BSE should be performed while (16).

Current study result regarding practicing of BSE is also similar to another study conducted on obstetric nurses in Korea in 2011 by Kim and Park. They suggested that 58.9% practiced self-breast examination. Further more a study was carried out in Jordanian university female students and were assessed used regression analysis which showed positive relationship between confidence and BSE performance.

CONCLUSION

BSE is a cheap and noninvasive tool to identify any abnormality in breast and also help to detect early sign of breast cancer. Teaching nursing students regarding BSE helps them to receive gain more knowledge and to disseminate the information in community in future.

REFERENCES

- [1] Stratton MR. Pathology of familial breast cancer: differences between breast cancers in carriers of BRCA1 or BRCA2 mutations and sporadic cases. *The Lancet*. 1997 May 24;349(9064):1505-10.
- [2] Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA: a cancer journal for clinicians*. 2011 Mar;61(2):69-90
- [3] Allen TL, Van Groningen BJ, Barksdale DJ, McCarthy R. The breast self-examination controversy: what providers and patients should know. *The Journal for Nurse Practitioners*. 2010 Jun 1;6(6):444-51.
- [4] Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation

- of relapse and survival with amplification of the HER-2/neu oncogene. *science*. 1987 Jan 9;235(4785):177-82.
- [5] Saadat, M. and Ansari-Lari, M., 2009. Polymorphism of XRCC1 (at codon 399) and susceptibility to breast cancer, a meta-analysis of the literatures. *Breast cancer research and treatment*, 115(1), pp.137-144.
- [6] Thomas DB, Gao DL, Ray RM, Wang WW, Allison CJ, Chen FL, Porter P, Hu YW, Zhao GL, Pan LD, Li W. Randomized trial of breast self-examination in Shanghai: final results. *Journal of the national Cancer Institute*. 2002 Oct 2;94(19):1445-57.
- [7] Lauby-Secretan B, Scoccianti C, Loomis D, Benbrahim-Tallaa L, Bouvard V, Bianchini F, Straif K. Breast-cancer screening—viewpoint of the IARC Working Group. *New England journal of medicine*. 2015 Jun 11;372(24):2353-8.
- [8] Taha H, Al-Qutob R, Nyström L, Wahlström R, Berggren V. "Voices of Fear and Safety" Women's ambivalence towards breast cancer and breast health: a qualitative study from Jordan. *BMC women's health*. 2012 Dec;12(1):21.
- [9] Im Kim J, Oh KO, Li CY, Min HS, Chang ES, Song R. Breast cancer screening practice and health-promoting behavior among Chinese women. *Asian nursing research*. 2011 Sep 1;5(3):157-63.
- [10] Savage SA, Clarke VA. Factors associated with screening mammography and breast self-examination intentions. *Health Education Research*. 1996 Dec 1;11(4):409-21.
- [11] Abolfotouh MA, Ala'a AB, Mahfouz AA, Al-Assiri MH, Al-Juhani AF, Alaskar AS. Using the health belief model to predict breast self examination among Saudi women. *BMC Public Health*. 2015 Dec;15(1):1163.
- [12] Gilani SI, Khurram M, Mazhar T, Mir ST, Ali S, Tariq S, Malik AZ. Knowledge, attitude and practice of a Pakistani female cohort towards breast cancer. *JPMA. The Journal of the Pakistan Medical Association*. 2010 Mar 1;60(3):205.
- [13] Balogun MO, Owoaje ET. Knowledge and practice of breast self-examination among female traders in Ibadan, Nigeria. *Annals of Ibadan Postgraduate Medicine*. 2005;3(2):52-6.
- [14] Aminisani N, Fattahpour R, Dastgiri S, Asghari-Jafarabadi M, Allahverdipour H. Determinants of breast cancer screening uptake in Kurdish women of Iran. *Health promotion perspectives*. 2016;6(1):42.
- [15] Gonzales A, Alzaatreh M, Mari M, Saleh AA, Alloubani A. Beliefs and Behavior of Saudi Women in the University of Tabuk Toward Breast Self Examination Practice. *Asian Pacific journal of cancer prevention: APJCP*. 2018;19(1):121.
- [16] Sama CB, Dzekem B, Kehbila J, Ekabe CJ, Vofo B, Abua NL, Dingana TN, Angwafo III F. Awareness of breast cancer and breast self-examination among female undergraduate students in a higher teachers training college in Cameroon. *Pan African Medical Journal*. 2017;28(1):164.

Citation: Aurang Zeb, Humera Naz, Nadim Khan, Tariq Aziz, Arifa Siddique. *Knowledge of Breast Self-Examination Among LHV's and Female Nursing Students of KP. Open Access Journal of Nursing*. 2019; 2(1): 11-13.

Copyright: © 2019 Aurang Zeb, Humera Naz, Nadim Khan, Tariq Aziz, Arifa Siddique. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.