

# Therapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage

Gamea RMR, El-Bhoty SB, El-Ebiary MT, El-Gharib MN

Department of Obstetrics and Gynecology, Faculty of Medicine, Tanta University, Tanta, Egypt.  
mohgharib2@hotmail.com

*\*Corresponding Author: El-Gharib MN, Department of Obstetrics and Gynecology, Faculty of Medicine, Tanta University, Tanta, Egypt.*

## Abstract

*The present cohort study was planned to investigate the clinical effect of sildenafil citrate (SC) tablets as well as nitric oxide (NO) patch for treatment of first trimester pregnant women with a history of unexplained recurrent spontaneous abortion.*

*The survey was carried on 60 first trimester pregnant women with a history of recurrent miscarriage. The patient was divided into two equal groups. The first patient group received SC 20 mg orally, 3 times/day and the second patient group received NO patch. We judged the effectiveness of this treatment on the patients.*

*We establish that both oral SC and NO patch decreased the incidence of spontaneous recurrent first trimester abortion.*

*In ending, we report evidence that both oral SC and NO patch can offer an optional therapy for cases of spontaneous recurrent first trimester abortion.*

**Keywords:** *Sildenafil citrate; Nitric oxide patch; Unexplained recurrent spontaneous miscarriage; Abortion.*

## INTRODUCTION

Recurrent miscarriage (RM) is a common pregnancy complication. It is defined as three or more consecutive pregnancy losses that take office before the fetus reaches viability, i.e., prior to 24 gestational weeks. [1] The etiology of RM is variable. [2] The mechanisms of early embryo loss observed are not yet fully realized. [3]

Approximately 15 to 20% of clinically recognized pregnancies end in miscarriage, and approximately 1 to 5% of pregnant women experience the recurrent miscarriage (RM), of which 40 to 55% are caused by unknown causes. [4]

A substantial body of experimental evidence suggests a relationship between nitric oxide (NO) and prostaglandins [5], particularly in pathophysiologic events associated with gestation. It is found that epidermal growth factor and IL-1 enhance PG production by stimulating NO synthase (iNOS) activity. [6]

El-Far et al., showed for the first time, in the preliminary report, that sildenafil citrate might be a novel, interesting, safe anti-abortive option in the treatment of threatened miscarriage in four patients with a history of unexplained recurrent spontaneous miscarriage. [7]

The aim of this study was to evaluate the effects of nitric oxide patch versus oral sildenafil citrate in treating women with a history of unexplained first trimester recurrent fetal wastage.

## PATIENTS AND METHODS

The present cohort study was a comparative clinical study that included 60 first trimester pregnant women who had been selected from the outpatient high risk pregnancy clinic of the Obstetrics Department of Tanta University Hospital during the period from June 2017 to June 2018.

The inclusion criteria of the selected women were: Age ranged from (20-35) years old, BMI  $\geq$  25 kg/m<sup>2</sup>, First trimester pregnant women with a history

## Therapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage

of two or more unexplained recurrent spontaneous first trimester miscarriage, Singleton pregnancy, gestational age (4-6 weeks) that was calculated from the first day of the last normal menstrual period and confirmed by transvaginal ultrasonography to detect intrauterine gestational sac.

The exclusion criteria were: patients that received any potential causes of abortion as: Congenital anomalies of the uterus, Luteal phase defect, Smokers, History of PCO or endometriosis, Abnormal karyotyping, Any chronic disorders including endocrine disorders, renal, cardiovascular, hepatic infectious disorders or immunological (APS).

A written informed consent from every patient included in this survey. The consent is turned out by the medical ethics committee of Tanta University Hospital.

### METHODS

All cases were subjected to the following:

1. Full history taking with special concern about the details of each abortion.
2. Through clinical general and abdominal exams.
3. Choosing blood pressure at booking and follow up in each visit using puppetry and auscultatory methods while patient in semi-sitting stance, then the average arterial blood pressure was estimated by  $MAP = 1/3 (SBP) + 2/3 (DBP)$  where (MAP is the mean arterial pressure, SBP is the systolic blood pressure, and DBP is the diastolic blood pressure.)
4. Investigations related to recurrent miscarriage including:
  - Routine laboratory investigations for pregnancy as (Rh factor, CBC, blood group and Urine analysis)
  - Immunological investigations (anticardiolipin, antinuclear and antiphospholipids antibodies).
  - Hormonal investigations (Thyroid function tests, fasting blood sugar, postprandial blood sugar, HbA1c and tests for thrombophilipitis)
5. Patients were randomly assigned into two groups:
  - Group A (SC group): Thirty patients received 20

mg sildenafil citrate (Respatio, Pharma Right Group) orally 3 times daily at booking till 14 weeks gestational age.

- Group B(NO group): Thirty patients used (50 mg) Nitroglycerin transdermal patch (Novartis Pharma AG, Stein, Switzerland) on the abdominal skin below the umbilicus, releasing nitroglycerin at a rate of 0.4 mg/h, once daily for 12 hours from the day of booking till 14 weeks gestational age.
- Both groups received folic acid 400 microgram/day.
- Transabdominal pulsed wave color Doppler imaging was performed for each selected woman before treatment, two hours after beginning treatment, and was repeated every 2 weeks, according to the antenatal schedule of high risk pregnancy till 14 weeks gestational age. It was done to evaluate the uterine arteries bilaterally and sub-endometrial blood flow (pulsatility index (PI) and restrictive index (RI).
- Ultrasound examination was performed for all selected women at the end of 14 weeks GA to exclude any fetal congenital anomalies.
- Ethics committee: The study started after medical ethical committee approval. All included cases have been informed about aim and the nature of this study, risk elements, possible side effect of the drug.

### RESULTS

Among the cases of SC group success pregnancy occurred in 65% of cases (13 out of 20 patients). Among cases of NO group a success of pregnancy took place in approximately 55% of cases (11 out of 20 patients). Normal babies were delivered in all cases of the two groups. The other results are summarized in 6 tables.

In SC group 2 patients refused to take the drug and 3 patients complained of severe headache therefore stopped drug administration. Four patients of NO group refused the application of NO patch to abdominal skin and 2 suffered from skin rash and severe headache.

### DISCUSSION

Recurrent early pregnancy loss accounts for almost 20% of clinically recognized pregnancy losses. Apart from the well-known risk factors, the effect of pregnancy

## Therapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage

depends, to a great extent, on the success rate of early events, such as implantation, the establishment of fetomaternal circulation, maintenance of increased blood flow on the implantation site. [8]

The vasoactive drug, sildenafil citrate has proven useful in reducing adverse pregnancy events. An important biological function of its action is an inhibition of cyclic guanosine monophosphate (cGMP) phosphodiesterase type 5 activity, which could have beneficial effects on trophoblast survival. [9]

Nitric oxide is required for trophoblast cell survival and appears to be critical for endovascular invasion necessary for normal placental perfusion and the associated phenotype switching. [10]

The purpose of this work was to assess the effects of nitric oxide patch versus oral sildenafil citrate in first-trimester pregnant women with a history of spontaneous recurrent miscarriage.

In the current study, we found that among the cases of SC group success of pregnancy occurred in 65% of cases. This may be explicated by the fact that sildenafil citrate has been proven to be useful in increasing endometrial thickness and achieving pregnancy in women with varied uterine disorders. It is suspected that this medication causes selective vasodilatation, resulting in improved endometrial development. [11]

The administration of sildenafil enhanced endometrial development in 70% of patients studied. High implantation and ongoing pregnancy rates were accomplished in a cohort with a poor prognosis for success. [12]

Endometrial thickness is one of the most potent predictors of implantation rate and ongoing pregnancy success rate. The endometrial growth is dependent on the uterine blood flow and angiogenesis. Sildenafil citrate leads to smooth muscle relaxation and vasodilation. Because of these biological properties, it is a potential candidate for female infertility, especially in the management of thin endometrium, which leads to low implantation and pregnancy rates. [13]

El-Far and associates in 2009 conclude that the sildenafil citrate tablets used as suppositories might be a safe anti-abortive option in the treatment of threatened miscarriage in patients with a history of unexplained recurrent miscarriage. [7]

In addition, the contemporary study revealed that among cases of NO group a success of pregnancy took place in approximately 55% of cases. This can be ascribed to the reductive effects of NO on the contractile action of the human uterine cervix in early pregnancy. [14]

The present study depicted a decrease in the average arterial blood pressure (MAP) of both studied groups 2 hours after the administration of the drug (table 2). Nevertheless, there was a substantial divergence between the two groups in MAP during the 2 hours after medication and 14 weeks pregnant after drug usage with more decrease among the SC group. This can be excused by the fact that sildenafil citrate induces vasodilatation through inhibition of type 5 phosphodiesterase (PDE5). [15]

**Table1.** *The clinical data of the two study groups*

Groups	Age (years)		BMI (kg/m <sup>2</sup> )		Gestational age (Weeks)		Number of previous abortions		HB% (gm/dl)	
	SC (n=25)	NO patch (n=24)	SC (n=25)	NO patch (n=24)	SC (n=25)	NO patch (n=24)	SC (n=25)	NO Patch (n=24)	SC (n=25)	NO patch (n=24)
<b>Range</b>	22 - 35	20 - 33	25 - 30	22-33	4-6	4-6	2-4	2-4	10.5 - 12.5	11-12.4
<b>Mean ± SD</b>	27.32 ±4.298	26.88± 3.745	28.40± 1.291	27.46± 2.859	7.12± .833	6.92±0.83	2.48± .653	2.67 ±0.637	11.556 ±0.5276	11.471 ±0.3581
<b>P. Value</b>	<b>0.701</b>		<b>0.141</b>		<b>0.396</b>		<b>0.317</b>		<b>0.514</b>	

SC= Sildenafil citrate

No = Nitroglycerine

The Doppler study of this work indicated a strong reduction in sub-endometrial arteries RI and both uterine arteries RI in the two groups in the 2 hours after medication and at 14 weeks gestation (tables 3-6). EL-Far and associates set up similar effects.[7]

Trapani and co-workers in 2015 found that the use of sildenafil citrate or transdermal nitroglycerin is associated with a substantial decrease in both uterine artery Doppler and uterine artery Pulsatility index, as well as maternal arterial blood pressure. [16]

## Therapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage

**Table2:** The mean arterial blood pressure (MAP)  $((d*2) +s)/3$ , of the two study groups.

	Mean arterial blood pressure $((d*2) +s) /3$ (mm Hg) with sildenafil (n=25)						Mean arterial blood pressure $((d*2) +s)/3$ mmHg with nitrodermal patch (n=24)					
	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)
<b>Range (mm Hg)</b>	87-102	80 -92	79-91	76-90	73-90	70 - 90	85-105	81- 97	80-95	78-92	76-90	75-95
<b>Mean ± SD</b>	94.8± 4.664	79.52± 4.363	78.21± 4.253	77.42± 4.458	77.52 ±4.524	77.2± 4.635	96.17± 4.975	88.58± 4.742	87.92± 4.453	86.52± ±4.668	85.38± 4.763	84.92± m4.80
<b>P-value</b>	<0.001*						<0.001*					
	0.1410											

GA = Gestational age

**Table3.** The resistive index of the meanuterine artery of the two study groups.

	Mean uterine artery resistive index (RI) With sildenafil (n=25)						Mean uterine artery resistive index (RI) With nitrodermal patch (n=24)					
	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)
<b>Mean ± SD</b>	0.601 ±0.0435	0.488 ±0.0275	0.472 ±0.024	0.47 ±0.023	0.451 ±0.022	0.434 ±0.019	0.597 ±0.0406	0.453 ±0.027	0.434 ±0.026	0.429 ±0.025	0.411 ±0.02	0.4079 ±0.01793
<b>P-value</b>	<0.001*						<0.001*					
	0.024398											

GA = Gestational age

**Table4.** The pulsatility index (PI) of the mean uterine artery of the two study groups.

	Mean uterine artery (PI) With sildenafil (n=25)						Mean uterine artery (PI) With nitrodermal patch (n=24)					
	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)
<b>Mean ± SD</b>	1.98 ±0.14	1.5 ±0.09	1.32 ±0.08	1.24 ±0.062	1.15 ±0.059	1.1 ±0.055	1.85 ±0.124	1.35 ±0.0852	1.26 ±0.074	1.21 ±0.0598	1.15 ±0.053	0.92 ±0.0492
<b>P-value</b>	<0.001*						<0.001*					
	0.20678											

GA = Gestational age

## Therapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage

**Table5.** Sub-endothelial arteries resistive index (RI) of the two study groups.

	Sub-endothelial artery (RI) With sildenafil (n=25)						Sub-endothelial artery (RI) With nitrodermal patch (n=24)					
	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)
<b>Mean ± SD</b>	0.872 ±0.0456	0.7648 ±0.035	0.744 ±0.0332	0.724 ±0.03125	0.708 ±0.02856	0.698 ±0.0245	0.925 ±0.052	0.829 ±0.0437	0.814 ±0.0421	0.78 ±0.0392	0.758 ±0.0375	0.727 ±0.0325
<b>P-value</b>	<0.001*						<0.001*					
	0.185223											

GA = Gestational age

**Table6.** Sub-endothelial arteries pulsatility index (PI) of the two study groups.

	Sub-endothelial a. (PI) With sildenafil (n=25)						Sub-endothelial a. (PI) With nitrodermal patch (n=24)					
	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)	Before	2 hours after	8 weeks GA	10 weeks GA	12 weeks GA	14 weeks GA (n=23)
<b>Mean ± SD</b>	2.88 ±0.62	1.98 ±0.58	1.89 ±0.54	1.85 ±0.51	1.83 ±0.49	1.79 ±0.472	2.92 ±0.65	2.15 ±0.612	1.92 ±0.584	1.872 ±0.542	1.852 ±0.529	1.819 ±0.518
<b>P-value</b>	<0.001*						<0.001*					
	0.784688											

GA = Gestational age

Recently, nitric oxide synthase (eNOS) has been reported to be over-expressed in the glandular and luminal epithelium of the endometrium in women with recurrent miscarriage and unexplained infertility, suggesting a detrimental effect of excess NO on endometrial receptivity and implantation. [17] Lastly, we close that both SC and NO patch are additional choices for treatment of unexplained recurrent spontaneous abortion.

### DISCLOSURE

We announce that we did not get any fund from whatever individual or establishment. If the patient refused to finish the field, she was taken out and replaced by another one from who are satisfying the inclusion criteria of the survey. We did not classify the patients according to their religious belief or civilization or race or any other unrelated points.

### REFERENCES

[1] Wang L, Tang H, Xiong Y, Tang L. Differential expression profile of long noncoding RNAs

in human chorionic villi of early recurrent miscarriage. Clin Chim Acta 2017;464:17e23.

- [2] Rull K, Tomberg K, Kořks S, Mañnik J, Mořls M, Sirotkina M, et al. Increased placental expression and maternal serum levels of apoptosis-inducing TRAIL in recurrent miscarriage. Placenta 2013;34:141e8.2.
- [3] Haddad EK, Duclos AJ, and Baines MG. Early Embryo Loss Is Associated with Local Production of Nitric Oxide by Decidual Mononuclear Cells. J. Exp. Med. 1995; 182: 1143-1152.
- [4] Yamada H, Kato EH, Kobashi G, Ebina Y, et al. Repeated pregnancy loss: etiology of thrombophilia. Semin. Thromb. Hemost. 2001; 27: 121-129.
- [5] Salvemini D, Masferrer JL. Methods Enzymol. 1996; 269:12–25.13. & Mollace V, Muscoli C, Masini E, Cuzzocrea S, Salvemini A. Pharmacol Rev. 2005; 57:217–252.

## Therapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage

- [6] Ribeiro ML, Perez Marti´nez S, Farina M, Ogando D, Gimeno M, Franchi AM. Prostaglandins Leukot Essent Fatty Acids.1999; 61:353–358. & Farina M, Ribeiro ML, Ogando D, Gimeno M, Franchi AM. Prostaglandins Leukot Essent Fatty Acids. 2000; 62:243–247.
- [7] El-Far, M, El-Motwally G, Hashem IA, and Bakry N. Biochemical role of intravaginal sildenafil citrate as a novel antiabortive agent in unexplained recurrent spontaneous miscarriage: first clinical study of four case reports from Egypt. Clin Chem Lab Med. 2009; 47(11): 1433-8.
- [8] Suryanarayana V, Rao I, Kanakavalli M, Padmalatha V, Deenadayal M and Singh L. Recurrent early pregnancy loss and endothelial nitric oxide synthase gene polymorphisms. Arch Gynecol Obstet. 2006; 274: 119–124.
- [9] Jay M. Bolnick, Michael P. Diamond, Brian A. Kilburn, Alan D. Bolnick, Michael Hertz, Manvinder Singh, D. Randall Armant, and Jing Dai. Sildenafil Prevents Apoptosis of Human First-Trimester Trophoblast Cells Exposed to Oxidative Stress: Possible Role for Nitric Oxide Activation of 3-cyclic Guanosine Monophosphate Signaling. Reproductive Sciences. 2015; 22(6): 718-724.
- [10] Huang LT, Hsieh CS, Chang KA, Tain YL. Roles of nitric oxide and asymmetric dimethylarginine in pregnancy and fetal programming. Int J Mol Sci. 2012;13(11):14606-14622.
- [11] Zinger M, Liu JH, and Thomas MA. Successful use of vaginal sildenafil citrate in two infertility patients with Asherman’s syndrome. J Womens Health (Larchmt). 2006;15(4):442-4.
- [12] Sher G and Fisch JD. The Effect of vaginal sildenafil on teh outcome of in vitro fertilization (IVF) after multiple IVF failures attributed to poor endometrial development. Fertil Steril. 2002; 78(5):1073-6.)
- [13] Jyoti M. Benni, Paragouda A. Patil. An overview on sildenafil and female infertility. Indian Journal of Health Sciences. 2016; 9 (2): 131-136.
- [14] Ekerhovd E, Brannstrom M, Delbro D and Norstrom A. Nitric oxide mediated inhibition of contractile activity in human uterine cervix. Molecular Human Reproduction. 1999; 4(9): 915-922.
- [15] Wareing, M., J.E. Myers, M. O’Hara, and P.N. Baker. Sildenafil citrate (Viagra) enhances vasodilatation in fetal growth restriction. J Clin Endocrinol Metab. 2005; 90(5): 2550-5.
- [16] Trapani A Jr, Gonçaves LF, Trapani TF, Franco MJ, Galluzzo RN, Pires MM. Ultrasound Obstet Gynecol. 2016; 48 (1): 61-5.
- [17] Najafi T, Novin MG, Ghazi R and Khorram O (2012). Altered endometrial expression of endothelial nitric oxide synthase in women with unexplained recurrent miscarriage and infertility. Reprod. Biomed. Online. 2012; 25: 408-414.

**Citation:** Gamea RMR, El-Bhoty SB, El-Ebiary MT, El-Gharib MN. *RTherapeutic Effects of Oral Sildenafil Citrate Versus Nitric Oxide Patch on Pregnant Women with a History of Recurrent Miscarriage. Archives of Reproductive Medicine and Sexual Health . 2018; 1(2): 30-35.*

**Copyright:** © 2018 Gamea RMR, El-Bhoty SB, El-Ebiary MT, El-Gharib MN. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*