

## RESEARCH ARTICLE

# Three-Center One-Electron Molecular Integrals over Dirac Wave Functions for Solving the Molecular Matrix Dirac Equation

**Kazuhiro Ishida**

Matsugasaki 219-48, Kashiwa City, Chiba 277-0835, Japan.

\*A part of this research was presented at the 19th annual meeting of Japan Society for Molecular Science held at Hiroshima during September 9-12, 2025.

Received: 03 November 2025 Accepted: 17 November 2025 Published: 12 December 2025

Corresponding Author: Kazuhiro Ishida. Matsugasaki 219-48, Kashiwa City, Chiba 277-0835, Japan.

**Abstract**

New Gaussian-transform formulas can be derived for special derivatives of Dirac wave function. Using the transform formulas, the title molecular integrals over Dirac wave functions can be derived. All molecular integral formulas can be derived for the first time. We should add the Dirac wave function to our basis set for solving the molecular matrix Dirac equation.

**Keywords:** Molecular Integrals, Molecular Dirac Equation, Relativistic Calculation, Dirac Wave Function, NMR Spectra.

## 1. Introduction

Generally speaking, the fundamental equation of the physics must be gauge invariant. The Dirac equation is the fundamental one of the relativistic quantum mechanics. Recently, Yoshizawa [1] derived the gauge invariant matrix Dirac equation with using the restricted magnetic balance [RMB] [2], as given by

$$\begin{pmatrix} \vec{V} & \vec{T}_m \\ \vec{T}_m & \vec{W}_m - \vec{T}_m \end{pmatrix} \begin{pmatrix} \vec{C}_-^L & \vec{C}_+^L \\ \vec{C}_-^S & \vec{C}_+^S \end{pmatrix} = \begin{pmatrix} \vec{S} & \vec{0} \\ \vec{0} & \frac{1}{2m_e c^2} \vec{T}_m \end{pmatrix} \begin{pmatrix} \vec{C}_-^L & \vec{C}_+^L \\ \vec{C}_-^S & \vec{C}_+^S \end{pmatrix} \begin{pmatrix} \vec{\epsilon}_- & \vec{0} \\ \vec{0} & \vec{\epsilon}_+ \end{pmatrix} \quad (1.1)$$

where  $\vec{C}_-^L$  is the coefficient matrix of the large component spinor for the energy matrix  $\vec{\epsilon}_-$ ,  $\vec{C}_+^L$ , is that for  $\vec{\epsilon}_+$ ,  $\vec{C}_-^S$ , and  $\vec{C}_+^S$  are those for the small component spinor,  $\vec{0}$  is the zero matrix,

$$V_{\mu\nu} = \langle \chi_\mu | V | \chi_\nu \rangle \quad (1.2)$$

$$(T_m)_{\mu\nu} = \frac{1}{2m_e} \langle \chi_\mu | \vec{\sigma} \cdot (\vec{p} + \vec{A}) \vec{\sigma} \cdot (\vec{p} + \vec{A}) | \chi_\nu \rangle \quad (1.3)$$

$$(W_m)_{\mu\nu} = \frac{1}{4m_e^2 c^2} \langle \chi_\mu | \vec{\sigma} \cdot (\vec{p} + \vec{A}) V \vec{\sigma} \cdot (\vec{p} + \vec{A}) | \chi_\nu \rangle \quad (1.4)$$

and

$$S_{\mu\nu} = \langle \chi_\mu | \chi_\nu \rangle \quad (1.5)$$

in which  $m_e$  is the electron rest mass,  $c$  is the speed of light,  $\vec{\sigma}$  is the Pauli spin matrices,  $\vec{p} = -i\hbar\vec{\nabla}$  is the momentum, and  $\vec{A}$  is the vector potential due to the nuclear spin. The vector potential must be included for the invariance of the Dirac equation, as shown by Sun et al. [3]. We choose it as the Gauss-type charge density

**Citation:** Kazuhiro Ishida. Three-Center One-Electron Molecular Integrals over Dirac Wave Functions for Solving the Molecular Matrix Dirac Equation. Open Access Journal of Chemistry 2025; 7(2): 62-88.

©The Author(s) 2025. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

distribution [GCDD] model as given by

$$\vec{A} = \frac{Z_M e}{c^2} \frac{4}{\sqrt{\pi} r_0^3} F_1 \left( \frac{r_M^2}{r_0^2} \right) \vec{\mu}_M \times \vec{r}_M \quad (1.6)$$

where  $Z_M e$  is the nuclear charge of the M-th nucleus in the case that the Dirac equation is extended to the molecule,  $\vec{\mu}_M = (\mu_{Mx}, \mu_{My}, \mu_{Mz})$  is the nuclear magnetic moment,  $\vec{r}_M = \vec{r} - \vec{M} = (x_M, y_M, z_M)$  is the coordinate of the electron,  $r_0$  is the scale parameter for the finite nucleus of GCDD model, and  $F_m(x) = \int_0^1 dt t^{2m} \exp(-xt^2)$  is the molecular incomplete gamma function. We use the operator notation for all integrals. Thus  $\int_0^1 dt$  is the integral operator, which integrates the integrand followed to it. We use the atomic units throughout the present article ( $m_e = 1$ ,  $e = 1$ ,  $\hbar = 1$ ,  $4\pi\epsilon_0 = 1$ ,  $c = 137.035999139$ ). However, we describe  $m_e$ ,  $e$ , and  $\hbar$  explicitly, for the readers convenience when one converts the units to the natural units. Some experiment shows that the real nucleus is not the point-like one but a finite-sized [4]. However, the charge distribution in the finite nucleus is not determined. We choose the GCDD model in the present article. In the statistics, the Gauss-type distribution is called as the normal one. The gauge invariant Dirac equation has no rigorous solution. To solve it, we use a proper basis set,  $\{\chi_\mu\}$ .

Many researchers extend the matrix Dirac equation to the molecule [1,2,5-18]. Especially, many are for relativistic calculations of NMR spectra [1,2,14-18]. We may call the extended Dirac equation as the molecular matrix Dirac one. It is natural to add the atomic Dirac wave function to our basis set. However, all researchers use the Gaussian-type orbitals (GTOs) for their basis set, because there is no molecular integral-formula for the Dirac wave function. In previous three articles [19-21], the author derived the Gaussian-transform formulas for the Dirac wave function [19] and for its first derivative [20]. Using the transform formulas, the author derived several molecular integral-formulas over Dirac wave functions as follows: (a) He derived integral-formulas

for the fundamental properties [20], as the overlap integral,  $S_{\mu\nu}$ , the kinetic energy one,  $\frac{1}{2m_e} \langle \chi_\mu | p^2 | \chi_\nu \rangle$ , the nuclear attraction one,  $V_{\mu\nu}$  of the point-like nucleus,  $V = -\frac{Z_M e^2}{r_M}$ , and of the GCDD model,

$$V = -Z_M e^2 \frac{2}{\sqrt{\pi} r_0} F_0 \left( \frac{r_M^2}{r_0^2} \right), \quad (1.7)$$

the electron-repulsion one,  $\langle \chi_\mu \chi_\kappa | V | \chi_\nu \chi_\lambda \rangle$ , of the usual one,  $V = \frac{e^2}{r_{12}}$ , and of the finite-sized electron,  $V = \frac{2e^2}{\sqrt{\pi} r_e} F_0 \left( \frac{r_{12}^2}{r_e^2} \right)$  ( $r_e$  is the classical radius of the electron), where  $\chi_\mu = r_A^{-\epsilon_A} \exp(-\zeta_A r_A)$  is the atomic Dirac wave function centered at A. Note that the Dirac wave function is singular at the position of the nucleus located at A. Therefore, the Dirac wave function cannot be written as a linear combination of GTOs.

(b) Next, he did those for relativistic kinetic energy terms given by Eq. (1.3) [21].

(c) He did that for the physical quantity [19],  $\langle \chi_\mu | i\vec{\sigma} \cdot (\vec{p} \times V \vec{A} + \vec{A} \times V \vec{p}) | \chi_\nu \rangle$ , of the homogeneous charge density [HCDD] model,

$$V = \begin{cases} -\frac{3Z_M e^2}{2r_{0H}} \left( 1 - \frac{1}{3} \frac{r_M^2}{r_{0H}^2} \right) & (0 \leq r_M \leq r_{0H}) \\ -\frac{Z_M e^2}{r_M} & (r_M > r_{0H}) \end{cases}, \quad (1.8)$$

where  $r_{0H}$  is the radius of the finite-sized nucleus in the HCDD model, and of the GCDD model,  $V$  is given by Eq. (1.7). Further, the author showed that the GTO cannot describe this quantity correctly for the case

of hydrogen atom [22]. In the present article, we derive the Gaussian-transform formula for a special first derivative and a special second derivative of the Dirac wave function in the next section. Using the transform formulas, we derive the quantity given by Eq. (1.4), for the GCDD model in the third section.

## 2. New Gaussian-Transform Formulas

First, we derive a new Gaussian-transform for the special first-derivative of the Dirac wave function. The first-derivative of the Dirac wave function can be written as

$$\nabla r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) = -\vec{r}_A \left( \frac{\varepsilon_A}{r_A^{2+\varepsilon_A}} + \frac{\zeta_A}{r_A^{1+\varepsilon_A}} \right) \exp(-\zeta_A r_A). \quad (2.1)$$

The special first-derivative can be written as

$$\vec{r}_A \cdot \nabla r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) = -(\varepsilon_A r_A^{-\varepsilon_A} + \zeta_A r_A^{1-\varepsilon_A}) \exp(-\zeta_A r_A). \quad (2.2)$$

The author derived the Gaussian-transform of the first term in the right-hand side of Eq. (2.2) in a previous article [19] as given by

$$-\varepsilon_A r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) = \frac{-\varepsilon_A \zeta_A^{1+\varepsilon_A}}{2\sqrt{\pi} \Gamma(1+\varepsilon_A)} \int_0^\infty dS S^{-3/2} \exp(-S r_A^2) \left[ \frac{\zeta_A^2}{2S} \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{4+\varepsilon_A}} - \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{2+\varepsilon_A}} \right] \exp\left[-\frac{\zeta_A^2}{4St^2}\right] \quad (2.3)$$

The Gaussian-transform of the second term in the right-hand side of Eq. (2.2) can be derived as follows: We have

$$\zeta_A r_A^{1-\varepsilon_A} \exp(-\zeta_A r_A) = \frac{\zeta_A r_A^2}{r_A^{1+\varepsilon_A}} \exp(-\zeta_A r_A). \quad (2.5)$$

We know the identity described as the formula number 3.471.3 in the Gradshteyn and Ryzhik [23] given by

$$\frac{\exp(-\beta)}{\beta^\nu} = \frac{1}{\Gamma(\nu)} \int_0^1 dt \frac{(1-t)^{\nu-1}}{t^{\nu+1}} \exp\left(-\frac{\beta}{t}\right). \quad (2.6)$$

The author derived the Gaussian-transform of the ns-STO in a previous article [24] given by,

$$r_A^{n_A} \exp(-\zeta_A r_A) = \frac{\zeta_A^{n_A}}{2^{n_A} \sqrt{\pi}} \sum_{i_A=0}^{\infty} (-1)^{i_A} (2i_A - 1)!! \binom{n_A}{2i_A} \left(\frac{2}{\zeta_A^2}\right)^{i_A} \int_0^\infty dS S^{-n_A+i_A-1/2} \exp\left[-\frac{\zeta_A^2}{4S} - Sr_A^2\right] \quad (2.7)$$

Using Eq. (2.6) with  $\beta = \zeta_A r_A$  and  $\nu = 1 + \varepsilon_A$  and doing Eq. (2.7) with  $\zeta_A = \frac{\zeta_A}{t}$  and  $n_A = 2$ , we have

$$\zeta_A r_A^{1-\varepsilon_A} \exp(-\zeta_A r_A) = \frac{\zeta_A^{3+\varepsilon_A}}{4\sqrt{\pi} \Gamma(1+\varepsilon_A)} \int_0^\infty dS S^{-5/2} \exp(-Sr_A^2) \left[ \frac{\zeta_A^2}{2S} \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{5+\varepsilon_A}} - 3 \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{3+\varepsilon_A}} \right] \exp\left[-\frac{\zeta_A^2}{4St^2}\right] \quad (2.8)$$

Substituting Eq. (2.5) and (2.8) into Eq. (2.2), we have the final formula for the special first-derivative given by

$$\begin{aligned} -\vec{r}_A \cdot \nabla r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) &= \frac{\zeta_A^{1+\varepsilon_A}}{2\sqrt{\pi} \Gamma(1+\varepsilon_A)} \int_0^\infty dS \exp(-Sr_A^2) \left[ \frac{\zeta_A^2}{2S} \left( \frac{\zeta_A^2}{2S} \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{5+\varepsilon_A}} - 3 \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{3+\varepsilon_A}} \right) \right. \\ &\quad \left. + \varepsilon_A \left( \frac{\zeta_A^2}{2S} \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{4+\varepsilon_A}} - \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{2+\varepsilon_A}} \right) \right] \exp\left[-\frac{\zeta_A^2}{4St^2}\right] \end{aligned} \quad (2.9)$$

Next, we derive the Gaussian-transform for the special second-derivative of the Dirac wave function given by

$$\nabla^2 r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) = \left[ \frac{\varepsilon_A(\varepsilon_A-1)}{r_A^{2+\varepsilon_A}} + \frac{2\varepsilon_A(\varepsilon_A-1)}{r_A^{1+\varepsilon_A}} + \frac{\zeta_A^2}{r_A^{\varepsilon_A}} \right] \exp(-\zeta_A r_A) \quad (2.10)$$

Using a similar derivation to that from Eq. (2.2) to (2.9), we have the final formula for the special second-derivative given by

$$\nabla^2 r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) = \frac{\zeta_A^{3+\varepsilon_A}}{2\sqrt{\pi} \Gamma(1+\varepsilon_A)} \int_0^\infty dS S^{-3/2} \exp(-Sr_A^2) \left[ \frac{\zeta_A^2}{2S} \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{4+\varepsilon_A}} - \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{2+\varepsilon_A}} \right]$$

$$-\frac{1-\varepsilon_A}{1+\varepsilon_A} \left( \varepsilon_A \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{4+\varepsilon_A}} + (2+\varepsilon_A) \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{2+\varepsilon_A}} \right) \exp \left[ -\frac{\zeta_A^2}{4S_1 t^2} \right] \quad (2.11)$$

Note that, the GTO,  $\exp(-S\mathbf{r}_A^2)$ , can be converted to the object function by the integral-transform as in Eq. (2.9) and (2.11).

### 3. Molecular Integrals

We derive molecular integral-formulas for physical quantities given by Eq. (1.4) as follows: Using the Dirac identity [25], we have

$$(W_m)_{\mu\nu} = \frac{1}{4m_e^2 c^2} \langle \chi_\mu | \vec{\sigma} \cdot (\vec{p} + \vec{A}) V \vec{\sigma} \cdot (\vec{p} + \vec{A}) | \chi_\nu \rangle = \frac{1}{4m_e^2 c^2} \langle \chi_\mu | (\vec{p} + \vec{A}) \cdot V (\vec{p} + \vec{A}) + i \vec{\sigma} \cdot [(\vec{p} + \vec{A}) \times V (\vec{p} + \vec{A})] | \chi_\nu \rangle \quad (3.1)$$

The first physical quantity in the right-hand side of Eq. (3.1) can be written as three terms as given by

$$(\vec{p} + \vec{A}) \cdot V (\vec{p} + \vec{A}) = \vec{p} \cdot V \vec{p} + [\vec{p} \cdot V \vec{A} + \vec{A} \cdot V \vec{p}] + \vec{A} \cdot V \vec{A} \quad (3.2)$$

The second physical quantity can be written as two terms as given by

$$i \vec{\sigma} \cdot [(\vec{p} + \vec{A}) \times V (\vec{p} + \vec{A})] = i \vec{\sigma} \cdot (\vec{p} \times V \vec{p}) + i \vec{\sigma} \cdot (\vec{p} \times V \vec{A} + \vec{A} \times V \vec{p}) \quad (3.3)$$

The author already derived the molecular integral-formula over Dirac wave functions for the last term in the right-hand side of Eq. (3.3) [19]. Thus, we derive molecular integral-formulas for remaining four terms for the finite nucleus of the GCDD model in the next subsections individually.

#### 3.1 The Term $\vec{p} \cdot V \vec{p}$

We derive molecular integral-formula for the term  $\vec{p} \cdot V \vec{p}$  with the GCDD model of  $V$ , which is given by Eq. (1.7), as follows: We have

$$\langle \chi_{\mu A} | \vec{p} \cdot V \vec{p} | \chi_{\nu B} \rangle = \int d\vec{r} r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) \vec{p} \cdot V \vec{p} r_B^{-\varepsilon_B} \exp(-\zeta_B r_B). \quad (3.1.1)$$

The latter part of the integrand in Eq. (3.1.1) can be written as

$$\begin{aligned} \vec{p} \cdot V \vec{p} r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) &= Z_M e^2 \hbar^2 \nabla \frac{2}{\sqrt{\pi} r_0} F_0 \left( \frac{r_M^2}{r_0^2} \right) \nabla r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) \\ &= Z_M e^2 \hbar^2 \left[ \frac{4}{\sqrt{\pi} r_0^3} F_1 \left( \frac{r_M^2}{r_0^2} \right) (-\vec{r}_M) \cdot \nabla + \frac{2}{\sqrt{\pi} r_0} F_0 \left( \frac{r_M^2}{r_0^2} \right) \nabla^2 \right] r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) \\ &= Z_M e^2 \hbar^2 \left[ \frac{4}{\sqrt{\pi} r_0^3} F_1 \left( \frac{r_M^2}{r_0^2} \right) (-\vec{r}_B) \cdot \nabla - \frac{4}{\sqrt{\pi} r_0^3} F_1 \left( \frac{r_M^2}{r_0^2} \right) \vec{M} \cdot \nabla + \frac{2}{\sqrt{\pi} r_0} F_0 \left( \frac{r_M^2}{r_0^2} \right) \nabla^2 \right] r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) \end{aligned} \quad (3.1.2)$$

where  $F_m$  is centered at  $\vec{M} = (0,0,0)$ . Substituting Eq. (3.1.2) into (3.1.1) and using the Gaussian-transform formulas Eq. (2.9) and (2.11) for the resulting integral, we have

$$I = \langle \chi_{\mu A} | \vec{p} \cdot V \vec{p} | \chi_{\nu B} \rangle = Z_M e^2 \hbar^2 \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{4\pi \Gamma(1+\varepsilon_A) \Gamma(1+\varepsilon_B)} \int_0^\infty dS_1 \int_0^\infty dS_2 (S_1 S_2)^{-3/2} f_0 (f_1 I_1 + f_2 I_2 + f_3 I_3) \quad (3.1.3)$$

where

$$f_0 = \left[ \frac{\zeta_A^2}{2S_1} \int_0^1 dt_1 \frac{(1-t_1)^{\varepsilon_A}}{t_1^{4+\varepsilon_A}} - \int_0^1 dt \frac{(1-t)^{\varepsilon_A}}{t^{2+\varepsilon_A}} \right] \exp \left[ -\frac{\zeta_A^2}{4S_1 t_1^2} \right] \quad (3.1.4)$$

$$f_1 = \left[ \frac{\zeta_B^2}{2S_2} \left( \frac{\zeta_B^2}{2S_2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{5+\varepsilon_B}} - 3 \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{3+\varepsilon_B}} \right) + \varepsilon_B \left( \frac{\zeta_B^2}{2S_2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \right] \exp \left[ -\frac{\zeta_B^2}{4S_2 t_2^2} \right] \quad (3.1.5)$$

$$f_2 = \left( \varepsilon_B \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} + \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \exp \left[ -\frac{\zeta_B^2}{4S_2 t_2^2} \right] \quad (3.1.6)$$

$$f_3 = \left[ \frac{\zeta_B^2}{2S_2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} - \frac{1-\varepsilon_B}{1+\varepsilon_B} \left( \varepsilon_B \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} + (2+\varepsilon_B) \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \right] \exp \left[ -\frac{\zeta_B^2}{4S_2 t_2^2} \right] \quad (3.1.7)$$

$$I_1 = \int d\vec{r} \frac{4}{\sqrt{\pi} r_0^3} F_1 \left( \frac{r_M^2}{r_0^2} \right) \exp(-S_1 r_A^2 - S_2 r_B^2), \quad (3.1.8)$$

$$I_2 = \int d\vec{r} \frac{4\overrightarrow{MB} \cdot \overrightarrow{r_B}}{\sqrt{\pi} r_0^3} F_1 \left( \frac{r_M^2}{r_0^2} \right) \exp(-S_1 r_A^2 - S_2 r_B^2), \quad (3.1.9)$$

and

$$I_3 = \int d\vec{r} \frac{2\zeta_B^2}{\sqrt{\pi} r_0} F_0 \left( \frac{r_M^2}{r_0^2} \right) \exp(-S_1 r_A^2 - S_2 r_B^2). \quad (3.1.10)$$

We first evaluate  $I_1$ . We use the Gaussian product rule given by

$$\exp(-S_1 r_A^2 - S_2 r_B^2) = \exp \left[ -\frac{S_1 S_2}{S_{12}} \overline{AB}^2 - S_{12} r_P^2 \right] \quad (3.1.11)$$

where  $S_{12} = S_1 + S_2$  and  $\vec{P} = \frac{S_1}{S_{12}} \vec{A} + \frac{S_2}{S_{12}} \vec{B}$ . Next, we use the Sack's formula given by [26]

$$\exp(-S_{12} r_P^2) = 4\pi \exp(-S_{12} r_M^2 - S_{12} \overline{MP}^2) \sum_{l=0} i_l(2S_{12} \overline{MP} r_M) \sum_{m=-l}^l Y_l^m(\widehat{MP}) Y_l^m(\widehat{r_M})^*, \quad (3.1.12)$$

where  $i_l(x)$  is the modified spherical Bessel function of the first kind and  $Y_l^m(\widehat{MP})$  is the spherical harmonics. We know that

$$i_l(x) = \frac{x^l}{(2l+1)!!} \sum_{j=0} \frac{(x^2/4)^j}{j!(l+3/2)_j} \quad (3.1.13)$$

where  $(a)_j = a(a+1) \cdots (a+j-1)$  is the Pochhammer symbol. We use the Gaussian product rule again as given by

$$\exp \left[ -\frac{S_1 S_2}{S_{12}} \overline{AB}^2 \right] \exp(-S_{12} \overline{MP}^2) = \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2). \quad (3.1.14)$$

Using Eq. (3.1.11), (3.1.12), and (3.1.14) for Eq. (3.1.8), we have

$$I_1 = 4\pi \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) I_{1a} \quad (3.1.15)$$

where

$$I_{1a} = \frac{4}{\sqrt{\pi} r_0^3} \int_0^\infty dr_M F_1 \exp(-S_{12} r_M^2) \sum_{l=0} i_l(2S_{12} \overline{MP} r_M) \int \widehat{r_M} \sum_{m=-l}^l Y_l^m(\widehat{MP}) Y_l^m(\widehat{r_M})^*. \quad (3.1.16)$$

The angular part can be evaluated as in a previous article [24] given by

$$\int \widehat{r_M} \sum_{m=-l}^l Y_l^m(\widehat{MP}) Y_l^m(\widehat{r_M})^* = \delta_{l0}. \quad (3.1.17)$$

Thus, we have

$$I_{1a} = \frac{4}{\sqrt{\pi} r_0^3} \int_0^\infty dr_M F_1 \exp(-S_{12} r_M^2) i_0(2S_{12} \overline{MP} r_M) = I_{1a}^{in} + I_{1a}^{out} \quad (3.1.18)$$

where

$$I_{1a}^{in} = \frac{4}{\sqrt{\pi} r_0^3} \int_0^{R_0} dr_M F_1 \exp(-S_{12} r_M^2) i_0(2S_{12} \overline{MP} r_M), \quad (3.1.19)$$

and

$$I_{1a}^{out} = \frac{4}{\sqrt{\pi} r_0^3} \int_{R_0}^\infty dr_M F_1 \exp(-S_{12} r_M^2) i_0(2S_{12} \overline{MP} r_M), \quad (3.1.20)$$

in which  $R_0 = br_0$  ( $b = 7$ ) separates the inner and outer part of the finite-sized nucleus of the GCDD model. We choose  $b = 7$  by the reason described in a previous article [19].

In the inner part, we use the power series of the  $F_m(x)$  as given by

$$F_m(x) = \frac{1}{2m+1} {}_1F_1 \left( m + \frac{1}{2}; m + \frac{3}{2}; -x \right) = \frac{1}{2m+1} \sum \frac{(-x)^n (m+1/2)_n}{n! (m+3/2)_n}, \quad (3.1.21)$$

where  ${}_1F_1(a_1; c_1; -x)$  is the confluent hypergeometric function (CHF). In the outer part, we use the asymptotic expansion of the CHF given by

$$F_m(x) = \frac{\Gamma(m+1/2)}{2x^{m+1/2}}. \quad (3.1.22)$$

Using Eq. (3.1.21) for (3.1.19), we have

$$I_{1a}^{in} = \frac{4}{3\sqrt{\pi}r_0^3} \sum_{n=0}^{\infty} \frac{(3/2)_n(-1/r_0^2)^n}{n!(5/2)_n} \sum_{j=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(3/2)_j} \int_0^{R_0} dr_M r_M^{2n+2j+2} \exp(-S_{12}r_M^2), \quad (3.1.23)$$

The integral in Eq. (3.1.23) can be evaluated as given by

$$\begin{aligned} \int_0^{R_0} dr_M r_M^{2n+2j+2} \exp(-S_{12}r_M^2) &= \frac{1}{2} \int_0^{R_0^2} dx x^{n+j+1/2} \exp(-S_{12}x) = \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{n+j+\frac{3}{2}} \gamma\left(n+j+\frac{3}{2}; S_{12}R_0^2\right) \\ &= \frac{1}{2} R_0^{2n+2j+3} \frac{\Gamma(n+j+3/2)}{\Gamma(n+j+5/2)} {}_1F_1\left(n+j+\frac{3}{2}; n+j+\frac{5}{2}; -S_{12}R_0^2\right) \end{aligned} \quad (3.1.24)$$

where  $\gamma(\alpha; x)$  is the incomplete gamma function of the first kind, which is appeared as the formula number 8.354.1 in the. Gradshteyn and Ryzhik [23] as given by

$$\gamma(\alpha, x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{\alpha+n}}{n!(\alpha+n)} = x^{\alpha} \frac{\Gamma(\alpha)}{\Gamma(\alpha+1)} {}_1F_1(\alpha; \alpha+1; -x). \quad (3.1.25)$$

Substituting Eq. (3.1.24) into (3.1.23), we have

$$\begin{aligned} I_{1a}^{in} &= \frac{2b^3}{3\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-b^2)^n (3/2)_n}{n!(5/2)_n} \sum_{j=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2 R_0^2)^j}{j!(3/2)_j} \left[ \frac{\Gamma(n+j+3/2)}{\Gamma(n+j+5/2)} - S_{12} R_0^2 \frac{\Gamma(n+j+5/2)}{\Gamma(n+j+7/2)} + O(R_0^4) \right] \\ &= \frac{2b^3}{3\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-b^2)^n (3/2)_n}{n!(5/2)_n} \left[ \frac{\Gamma(n+3/2)}{\Gamma(n+5/2)} + \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) R_0^2 \frac{\Gamma(n+5/2)}{\Gamma(n+7/2)} + O(R_0^4) \right] \\ &= \frac{2b^3}{3\sqrt{\pi}} \left[ \frac{\Gamma(3/2)}{\Gamma(5/2)} {}_2F_2\left(\frac{3}{2}, \frac{3}{2}; \frac{5}{2}, \frac{5}{2}; -b^2\right) + \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) R_0^2 \frac{\Gamma(5/2)}{\Gamma(7/2)} {}_1F_1\left(\frac{3}{2}; \frac{7}{2}; -b^2\right) \right] + O(R_0^4) \end{aligned} \quad (3.1.26)$$

The error term  $O(R_0^4)$  is in the order of  $R_0^4 = 0.53179747(-15)$  for the case of hydrogen, which is very small.

It is easy to derive that the generalized hypergeometric function  ${}_2F_2\left(\frac{3}{2}, \frac{3}{2}; \frac{5}{2}, \frac{5}{2}; -b^2\right)$  can be expressed as the integral representation given by

$${}_2F_2\left(\frac{3}{2}, \frac{3}{2}; \frac{5}{2}, \frac{5}{2}; -b^2\right) = \frac{\Gamma(5/2)\Gamma(5/2)}{\Gamma(3/2)\Gamma(3/2)} \int_0^1 du \int_0^1 dv \sqrt{uv} \exp(-b^2uv) \quad (3.1.27)$$

The integral in Eq. (3.1.27) has a constant value, which can be evaluated by the Gauss-Legendre quadrature (GLQ). The constant value is 0.9961218237(-2), obtained from the 4096-point GLQ. The value of the  ${}_1F_1\left(\frac{3}{2}; \frac{7}{2}; -b^2\right)$  is also a constant. It can be evaluated by the asymptotic expansion of it from the formula number 13.5.1 of the Abramowitz and Stegun [27] as given by

$${}_1F_1\left(\frac{3}{2}; \frac{7}{2}; -b^2\right) = \frac{\Gamma(7/2)}{b^3} \left(1 - \frac{3}{2b^2}\right) \quad (3.1.28)$$

Thus, we have

$$I_{1a}^{in} = \frac{b^3}{\sqrt{\pi}} \int_0^1 du \int_0^1 dv \sqrt{uv} \exp(-b^2uv) + \frac{1}{2} \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) \left( R_0^2 - \frac{3}{2} r_0^2 \right) + O(R_0^4). \quad (3.1.29)$$

Next, we evaluate  $I_{1a}^{out}$ . Using Eq. (3.1.22), we have

$$I_{1a}^{out} = \int_{R_0}^{\infty} dr_M \frac{1}{r_M} \exp(-S_{12}r_M^2) i_0(2S_{12} \overline{MP} r_M) = \sum_{j=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(3/2)_j} \int_{R_0}^{\infty} dr_M r_M^{2j-1} \exp(-S_{12}r_M^2)$$

$$\begin{aligned}
&= \frac{1}{2} \sum_{j=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2)^j}{j! (3/2)_j} \int_{R_0^2}^{\infty} dx x^{j-1} \exp(-S_{12}x) = \frac{1}{2} \int_{R_0^2}^{\infty} dx \frac{1}{x} \exp(-S_{12}x) + \frac{1}{3} S_{12}^2 \overline{MP}^2 \sum_{j'=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2)^{j'}}{j'! (5/2)_{j'}} \int_{R_0^2}^{\infty} dx x^{j'} \exp(-S_{12}x) \\
&= \frac{1}{2} \Gamma(0, S_{12}R_0^2) + \frac{1}{3} S_{12}^2 \overline{MP}^2 \sum_{j'=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2)^{j'}}{j'! (5/2)_{j'}} \left(\frac{1}{S_{12}}\right)^{j'+1} \Gamma(j'+1, S_{12}R_0^2),
\end{aligned} \tag{3.1.30}$$

where  $\Gamma(\alpha, x)$  is the incomplete gamma function of the second kind. We know that

$$\Gamma(0, S_{12}R_0^2) = -E_i(-S_{12}R_0^2) = -\gamma - \ln(S_{12}R_0^2) - S_{12}R_0^2 + O(R_0^4) \tag{3.1.31}$$

where  $\gamma = 0.5772156649$  is the Euler constant and  $E_i(-x)$  is the exponential integral, and it is easy to derive

$$\Gamma(j'+1, S_{12}R_0^2) = \Gamma(j'+1) - \frac{(S_{12}R_0^2)^{j'+1}}{j'+1} + O(R_0^4) \tag{3.1.32}$$

Thus, we have

$$I_{1a}^{out} = -\frac{1}{2}\gamma - \frac{1}{2}\ln(S_{12}R_0^2) + \frac{1}{3}S_{12}\overline{MP}^2 {}_2F_2\left(1, 1; 2, \frac{5}{2}; S_{12}\overline{MP}^2\right) - \frac{1}{3}S_{12}^2\overline{MP}^2R_0^2 + \frac{1}{2}S_{12}R_0^2 + O(R_0^4) \tag{3.1.33}$$

Substituting Eq. (3.1.29) and (3.1.33) into (3.1.18), we have

$$I_{1a} = -\frac{1}{2}\ln(S_{12}R_0^2) + C_0 + \frac{1}{3}S_{12}\overline{MP}^2 {}_2F_2\left(1, 1; 2, \frac{5}{2}; S_{12}\overline{MP}^2\right) - \frac{1}{2}S_{12}^2\overline{MP}^2r_0^2 + \frac{3}{4}S_{12}r_0^2 + O(R_0^4) \tag{3.1.34}$$

where

$$C_0 = \frac{b^3}{\sqrt{\pi}} \int_0^1 du \int_0^1 dv \sqrt{uv} \exp(-b^2uv) - \frac{1}{2}\gamma = 1.639057508 \tag{3.1.35}$$

Next, we derive  $I_2$  and  $I_3$ . Using a similar derivation from Eq. (3.1.11) to (3.1.34) for  $I_1$ , we have

$$I_2 = 4\pi \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) I_{2a} \tag{3.1.36}$$

and

$$I_3 = 4\pi \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) I_{3a} \tag{3.1.37}$$

where

$$I_{2a} = \frac{\zeta_B^2}{3} \overrightarrow{MB} \cdot \overrightarrow{MP} {}_1F_1\left(1; \frac{5}{2}; S_{12}\overline{MP}^2\right) - \frac{\zeta_B^2}{2} \overrightarrow{MB} \cdot \overrightarrow{MP} S_{12}r_0^2 - \zeta_B^2 \overrightarrow{MB}^2 I_{1a} + O(R_0^4) \tag{3.1.38}$$

and

$$I_{3a} = \frac{1}{2} \frac{\zeta_B^2}{S_{12}} {}_1F_1\left(1; \frac{3}{2}; S_{12}\overline{MP}^2\right) - \frac{1}{4} \zeta_B^2 r_0^2 + O(R_0^4) \tag{3.1.39}$$

Substituting Eq. (3.1.34) into (3.1.15), doing (3.1.38) into (3.1.36), doing (3.1.39) into (3.1.37), and doing these resulting equations into (3.1.3), we have

$$\begin{aligned}
I &= Z_M e \hbar^2 \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{\Gamma(1+\varepsilon_A) \Gamma(1+\varepsilon_B)} \int_0^{\infty} dS_1 \int_0^{\infty} dS_2 (S_1 S_2)^{-3/2} \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) f_0 \\
&\quad \{(f_1 - f_2 \zeta_B^2 \overline{MB}^2) I_{1a} + f_2 I_{2a} + f_3 I_{3a}\} + O(R_0^4)
\end{aligned} \tag{3.1.40}$$

We evaluate the remaining integrals by the numerical integration. To do this, we first change integral variables

as follows: We set  $S_{12} = z$  and  $\frac{S_1}{S_{12}} = w$ . The Jacobian is given by

$$\frac{\partial(S_1 S_2)}{\partial(z, w)} = z \tag{3.1.41}$$

Then, we have

$$I = Z_M e \hbar^2 \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{\Gamma(1+\varepsilon_A) \Gamma(1+\varepsilon_B)} \int_0^1 dw [w(1-w)]^{-3/2} \int_0^{\infty} dz z^{-2} \exp[-wz \overline{MA}^2 - (1-w)z \overline{MB}^2] f_0$$

$$\{(f_1 - f_2 \zeta_B^2 \overline{MB}^2) I_{1a} + f_2 I_{2a} + f_3 I_{3a}\} + O(R_0^4) \quad (3.1.42)$$

Further, we separate integral over  $z$  as follows:  $\int_0^\infty dz = \int_0^{a^2} dz + \int_{a^2}^\infty dz$  where  $a^2$  can be chosen arbitrary. We choose  $a^2 = 4$  here. In the first integral in the right-hand side of the above, we change integral variable from  $z$  to  $u = a^2 z$  and do that from  $z$  to  $u = z/a^2$  in the second integral. Then we have

$$\int_0^\infty dz = a^2 \int_0^1 du + a^2 \int_0^1 du \frac{1}{u^2} \quad (3.1.43)$$

Thus, all integrals are for the interval  $[0, 1]$ , which can be evaluated numerically by using the GLQ. Using Eq. (3.1.43), we have the final formula given by

$$I = Z_M e \hbar^2 \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{a^2 \Gamma(1+\varepsilon_A) \Gamma(1+\varepsilon_B)} \int_0^1 dw [w(1-w)]^{-3/2} \left\{ \int_0^1 du \exp[-wua^2 \overline{MA}^2 - (1-w)ua^2 \overline{MB}^2] f_0^{(1)} \right. \\ \left. [f_1^{(1)} I_{1a}^{(1)} + f_2^{(1)} I_{2a}^{(1)} + f_3^{(1)} I_{3a}^{(1)}] + \int_0^1 du \exp\left[-\frac{w}{u} a^2 \overline{MA}^2 - \frac{1-w}{u} a^2 \overline{MB}^2\right] f_0^{(2)} [f_1^{(2)} I_{1a}^{(2)} + f_2^{(2)} I_{2a}^{(2)} + f_3^{(2)} I_{3a}^{(2)}] \right\} + O(R_0^4), \quad (3.1.44)$$

where

$$f_0^{(1)} = \left[ \frac{\zeta_A^2}{2wua^2} \int_0^1 dt_1 \frac{(1-t_1)^{\varepsilon_A}}{t_1^{4+\varepsilon_A}} - \int_0^1 dt_1 \frac{(1-t_1)^{\varepsilon_A}}{t_1^{2+\varepsilon_A}} \right] \exp\left[-\frac{\zeta_A^2}{4wua^2 t_1^2}\right], \quad (3.1.45)$$

$$f_1^{(1)} = \left[ \frac{\zeta_B^2}{2(1-w)ua^2} \left( \frac{\zeta_B^2}{2(1-w)ua^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{5+\varepsilon_B}} - 3 \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{3+\varepsilon_B}} \right) \right. \\ \left. + \varepsilon_B \left( \frac{\zeta_B^2}{2(1-w)ua^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \right] \exp\left[-\frac{\zeta_B^2}{4(1-w)ua^2 t_2^2}\right], \quad (3.1.46)$$

$$f_2^{(1)} = \left( \varepsilon_B \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} + \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \exp\left[-\frac{\zeta_B^2}{4(1-w)ua^2 t_2^2}\right], \quad (3.1.47)$$

$$f_3^{(1)} = \left[ \frac{\zeta_B^2}{2(1-w)ua^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right. \\ \left. - \frac{1-\varepsilon_B}{1+\varepsilon_B} \left( \varepsilon_B \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} + (2+\varepsilon_B) \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{3+\varepsilon_B}} \right) \right] \exp\left[-\frac{\zeta_B^2}{4(1-w)ua^2 t_2^2}\right], \quad (3.1.48)$$

$$I_{1a}^{(1)} = -\frac{1}{2u^2} \ln(ua^2 R_0^2) + \frac{C_0}{u^2} + \frac{1}{3u} a^2 x_0 {}_2F_2\left(1, 1; 2, \frac{5}{2}; ua^2 x_0\right) - \frac{1}{2} a^4 x_0^2 r_0^2 + \frac{3a^2}{4u} r_0^2, \quad (3.1.49)$$

$$I_{2a}^{(1)} = \frac{\zeta_B^2 y_0}{3u^2} {}_1F_1\left(1; \frac{5}{2}; ua^2 x_0\right) - \frac{\zeta_B^2 a^2 y_0}{2u} r_0^2 - \zeta_B^2 \overline{MB}^2 I_{1a}^{(1)}, \quad (3.1.50)$$

$$I_{3a}^{(1)} = \frac{1}{2} \frac{\zeta_B^2}{u^3 a^2} {}_1F_1\left(1; \frac{3}{2}; ua^2 x_0\right) - \frac{1}{4u^2} \zeta_B^2 r_0^2, \quad (3.1.51)$$

$$f_0^{(2)} = \left[ \frac{u \zeta_A^2}{2w a^2} \int_0^1 dt_1 \frac{(1-t_1)^{\varepsilon_A}}{t_1^{4+\varepsilon_A}} - \int_0^1 dt_1 \frac{(1-t_1)^{\varepsilon_A}}{t_1^{2+\varepsilon_A}} \right] \exp\left[-\frac{u \zeta_A^2}{4w a^2 t_1^2}\right], \quad (3.1.52)$$

$$f_1^{(2)} = \left[ \frac{u \zeta_B^2}{2(1-w)a^2} \left( \frac{\zeta_B^2}{2(1-w)a^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{5+\varepsilon_B}} - 3 \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{3+\varepsilon_B}} \right) \right. \\ \left. + \varepsilon_B \left( \frac{u \zeta_B^2}{2(1-w)a^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \right] \exp\left[-\frac{u \zeta_B^2}{4(1-w)a^2 t_2^2}\right], \quad (3.1.53)$$

$$f_2^{(2)} = \left( \varepsilon_B \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} + \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \exp\left[-\frac{u \zeta_B^2}{4(1-w)a^2 t_2^2}\right], \quad (3.1.54)$$

$$f_3^{(2)} = \left[ \frac{u\zeta_B^2}{2(1-w)a^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right. \\ \left. - \frac{1-\varepsilon_B}{1+\varepsilon_B} \left( \varepsilon_B \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} + (2+\varepsilon_B) \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right) \right] \exp \left[ -\frac{u\zeta_B^2}{4(1-w)a^2 t_2^2} \right], \quad (3.1.55)$$

$$I_{1a}^{(2)} = -\frac{1}{2} \ln \left( \frac{a^2}{u} R_0^2 \right) + C_0 + \frac{1}{3u} a^2 x_0 {}_2F_2 \left( 1, 1; 2, \frac{5}{2}; \frac{a^2}{u} x_0 \right) - \frac{1}{2u^2} a^4 x_0^2 r_0^2 + \frac{3a^2}{4u} r_0^2, \quad (3.1.56)$$

$$I_{2a}^{(2)} = \frac{\zeta_B^2}{3} y_0 {}_1F_1 \left( 1; \frac{5}{2}; \frac{a^2}{u} x_0 \right) - \frac{\zeta_B^2}{2} \frac{a^2 y_0}{u} r_0^2 - \zeta_B^2 \overrightarrow{MA}^2 I_{1a}^{(2)}, \quad (3.1.57)$$

$$I_{3a}^{(2)} = \frac{u \zeta_B^2}{2 a^2} {}_1F_1 \left( 1; \frac{3}{2}; \frac{a^2}{u} x_0 \right) - \frac{1}{4} \zeta_B^2 r_0^2, \quad (3.1.58)$$

$$x_0 = w^2 \overrightarrow{MA}^2 + (1-w)^2 \overrightarrow{MB}^2 + 2w(1-w) \overrightarrow{MA} \cdot \overrightarrow{MB}, \quad (3.1.59)$$

and

$$y_0 = w \overrightarrow{MA} \cdot \overrightarrow{MB} + (1-w) \overrightarrow{MB}^2. \quad (3.1.60)$$

Using the 128-point GLQ, we have the value of  $I$  with 8 significant-figure precision. We obtain

$$I = 0.21580602 e\hbar^2 \text{ for the case of three hydrogen atoms located at } \vec{M} = (0, 0, 0), \vec{A} = \left( -\frac{\sqrt{8}}{3}, -\sqrt{\frac{8}{3}}, \frac{2}{3} \right), \text{ and}$$

$$\vec{B} = \left( -\frac{\sqrt{8}}{3}, \sqrt{\frac{8}{3}}, \frac{2}{3} \right).$$

### 3.2 The Term $\vec{A} \cdot \nabla \vec{A}$

We evaluate the molecular integral-formula over Dirac wave functions for the term  $\vec{A} \cdot \nabla \vec{A}$  with potentials of the finite nucleus for the GCDD model as follows: The  $\vec{A}$  is given by Eq. (1.6). We have

$$\langle \chi_{\mu A} | \vec{A} \cdot V \vec{A} | \chi_{\nu B} \rangle = -\frac{Z_M^2 e^4}{c^4} \sum_{\xi, \eta} \mu_M \xi \mu_M \eta I_{\xi \eta} \quad [\xi, \eta \in \epsilon(x, y, z)] \quad (3.2.1)$$

where

$$I_{\xi \eta} = \frac{32}{\pi^{3/2} r_0^7} \int d\vec{r} F_0 F_1 F_1 (\delta_{\xi \eta} r_M^2 - \xi_M \eta_M) r_A^{-\varepsilon_A} r_B^{-\varepsilon_B} \exp(-\zeta_A r_A - \zeta_B r_B) \quad (3.2.2)$$

We first derive

$$I_{zz} = \frac{32}{\pi^{3/2} r_0^7} \int d\vec{r} F_0 F_1 F_1 (r_M^2 - z_M^2) r_A^{-\varepsilon_A} r_B^{-\varepsilon_B} \exp(-\zeta_A r_A - \zeta_B r_B) \quad (3.2.3)$$

We use the Gaussian-transform for the Dirac wave function centered at A, as given by

$$r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) = \frac{\zeta_A^{1+\varepsilon_A}}{2\sqrt{\pi} \Gamma(1+\varepsilon_A)} \int_0^\infty dS_1 S_1^{-3/2} \exp(-S_1 r_A^2) f_0 \quad (3.2.4)$$

where  $f_0$  is given by Eq. (3.1.4). We use also that centered at B and have

$$I_{zz} = \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{4\pi \Gamma(1+\varepsilon_A) \Gamma(1+\varepsilon_B)} \int_0^\infty dS_1 \int_0^\infty dS_2 (S_1 S_2)^{-3/2} f_0 f_{0B} I_1^{zz} \quad (3.2.5)$$

where

$$f_{0B} = \left[ \frac{\zeta_B^2}{2S_2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right] \exp \left[ -\frac{\zeta_B^2}{4S_2 t_2^2} \right] \quad (3.2.6)$$

and

$$I_1^{zz} = \frac{32}{\pi^{3/2} r_0^7} \int d\vec{r} \ F_0 F_1 F_1 (r_M^2 - z_M^2) \exp(-S_1 r_A^2 - S_2 r_B^2) \quad (3.2.7)$$

We know that  $r_M^2 - z_M^2 = \frac{2}{3} r_M^2 - \frac{2}{3} S_{20}(\vec{r}_M)$ . Then, we have

$$I_1^{zz} = I_2^{zz} + I_3^{zz} \quad (3.2.8)$$

where

$$I_2^{zz} = \frac{64}{3\pi^{3/2} r_0^7} \int d\vec{r} \ r_M^2 F_0 F_1 F_1 \exp(-S_1 r_A^2 - S_2 r_B^2), \quad (3.2.9)$$

and

$$I_3^{zz} = \frac{-64}{3\pi^{3/2} r_0^7} \int d\vec{r} \ F_0 F_1 F_1 \exp(-S_1 r_A^2 - S_2 r_B^2) S_{20}(\vec{r}_M). \quad (3.2.10)$$

Using the Gaussian-product rule, Eq. (3.1.11), the Sack's formula, Eq. (3.1.12), and the Gaussian-product rule again, Eq. (3.1.14), we have

$$I_2^{zz} = 4\pi \ \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) I_{2a}^{zz}, \quad (3.2.11)$$

and

$$I_3^{zz} = 4\pi \ \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) I_{3a}^{zz}, \quad (3.2.12)$$

where

$$I_{2a}^{zz} = \frac{64}{3\pi^{3/2} r_0^7} \int_0^\infty dr_M \ r_M^4 F_0 F_1 F_1 \exp(-S_{12} r_M^2) \sum_{l=0} i_l (2S_{12} \overline{MP} r_M) \ \int \widehat{r}_M \sum_{m=-l}^l Y_l^m(\widehat{MP}) Y_l^m(\widehat{r}_M)^*, \quad (3.2.13)$$

and

$$I_{3a}^{zz} = \frac{-64}{3\pi^{3/2} r_0^7} \int_0^\infty dr_M \ r_M^2 F_0 F_1 F_1 \exp(-S_{12} r_M^2) \sum_{l=0} i_l (2S_{12} \overline{MP} r_M) \ \int \widehat{r}_M \sum_{m=-l}^l Y_l^m(\widehat{MP}) Y_l^m(\widehat{r}_M)^* S_{20}(\vec{r}_M). \quad (3.2.14)$$

The angular part can be evaluated by using Eq. (3.1.17) and as in a previous article [24] as given by

$$\int \widehat{r}_M \sum_{m=-l}^l Y_l^m(\widehat{MP}) Y_l^m(\widehat{r}_M)^* S_{20}(\vec{r}_M) = r_M^2 \ \delta_{l2} \frac{S_{20}(\overline{MP})}{\overline{MP}^2}. \quad (3.2.15)$$

Thus, we have

$$I_{2a}^{zz} = \frac{64}{3\pi^{3/2} r_0^7} \int_0^\infty dr_M r_M^4 F_0 F_1 F_1 \exp(-S_{12} r_M^2) i_0 (2S_{12} \overline{MP} r_M), \quad (3.2.16)$$

and

$$I_{3a}^{zz} = \frac{-64}{3\pi^{3/2} r_0^7} \frac{S_{20}(\overline{MP})}{\overline{MP}^2} \int_0^\infty dr_M r_M^4 F_0 F_1 F_1 \exp(-S_{12} r_M^2) i_2 (2S_{12} \overline{MP} r_M). \quad (3.2.17)$$

We evaluate  $I_{2a}^{zz}$  as follows: We have

$$I_{2a}^{zz} = I_{2a}^{zzin} + I_{2a}^{zzout} \quad (3.2.18)$$

where

$$I_{2a}^{zzin} = \frac{64}{3\pi^{3/2} r_0^7} \int_0^{R_0} dr_M r_M^4 F_0 F_1 F_1 \exp(-S_{12} r_M^2) i_0 (2S_{12} \overline{MP} r_M), \quad (3.2.19)$$

and

$$I_{2a}^{zzout} = \frac{64}{3\pi^{3/2} r_0^7} \int_{R_0}^\infty dr_M r_M^4 F_0 F_1 F_1 \exp(-S_{12} r_M^2) i_0 (2S_{12} \overline{MP} r_M). \quad (3.2.20)$$

In order to evaluate  $I_{2a}^{zzin}$ , we use Eq. (3.1.13) and (3.1.21) and have

$$I_{2a}^{zzin} = \frac{64}{27\pi^{3/2}r_0^7} \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(3/2)_j} \sum_{n_1 n_2 n_3=0} \frac{(-1/r_0^2)^{n_1+n_2+n_3} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} I_{2b}^{zzin} \quad (3.2.21)$$

where

$$\begin{aligned} I_{2b}^{zzin} &= \int_0^{R_0} dr_M \ r_M^{2(j+n_{123})+4} \exp(-S_{12}r_M^2) = \frac{1}{2} \int_0^{R_0^2} dx \ x^{j+n_{123}+3/2} \exp(-S_{12}x) = \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+n_{123}} \gamma(j+n_{123}; S_{12}R_0^2) \\ &= \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+n_{123}+5/2} (S_{12}R_0^2)^{j+n_{123}+5/2} \frac{\Gamma(j+n_{123}+5/2)}{\Gamma(j+n_{123}+7/2)} {}_1F_1 \left( j+n_{123} + \frac{5}{2}; j+n_{123} + \frac{7}{2}; -S_{12}R_0^2 \right) \\ &= \frac{1}{2} (R_0^2)^{j+n_{123}+5/2} \frac{\Gamma(j+n_{123}+5/2)}{\Gamma(j+n_{123}+7/2)} {}_1F_1 \left( j+n_{123} + \frac{5}{2}; j+n_{123} + \frac{7}{2}; -S_{12}R_0^2 \right) \end{aligned} \quad (3.2.22)$$

where  $n_{123} = n_1 + n_2 + n_3$ . Substituting Eq. (3.2.22) into (3.2.21), we have

$$\begin{aligned} I_{2a}^{zzin} &= \frac{64}{27\pi^{3/2}r_0^2} \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(3/2)_j} \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} \\ &\quad \frac{\Gamma(j+n_{123}+5/2)}{\Gamma(j+n_{123}+7/2)} {}_1F_1 \left( j+n_{123} + \frac{5}{2}; j+n_{123} + \frac{7}{2}; -S_{12}R_0^2 \right) \\ &= \frac{32}{27\pi^{3/2}r_0^2} \left[ S_{A0} + \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) R_0^2 S_{A1} + \left( \frac{4}{15} S_{12}^4 \overline{MP}^4 - \frac{2}{3} S_{12}^3 \overline{MP}^2 + \frac{1}{2} S_{12}^2 \right) R_0^4 S_{A2} + O(R_0^6) \right] \end{aligned} \quad (3.2.23)$$

where

$$S_{A0} = \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} \frac{\Gamma(n_{123}+5/2)}{\Gamma(n_{123}+7/2)} \quad (3.2.24)$$

$$S_{A1} = \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} \frac{\Gamma(n_{123}+7/2)}{\Gamma(n_{123}+9/2)} \quad (3.2.25)$$

and

$$S_{A2} = \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} \frac{\Gamma(n_{123}+9/2)}{\Gamma(n_{123}+11/2)} \quad (3.2.26)$$

Summations, Eq. (3.2.24), (3.2.25) and (3.2.26), can be calculated in Appendix A.

Substituting Eq. (a.33), (a.35), and (a.41), into (3.2.23), we have

$$\begin{aligned} I_{2a}^{zzin} &= \frac{2}{3r_0^2} - \frac{1}{3R_0^2} - \frac{4}{3\pi r_0^2} - \frac{1}{\pi r_0^2} (C_{A0a} - C_{A0b} - C_{A0c} + C_{A0d}) \\ &+ \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) \left[ \frac{b^3}{3\sqrt{\pi}} C_{A1a} - \frac{8}{3\sqrt{\pi}} C_{A1b} - \frac{5}{2\pi} (C_{A1c} - C_{A1d}) \right] + \left( \frac{4}{15} S_{12}^4 \overline{MP}^4 - \frac{2}{3} S_{12}^3 \overline{MP}^2 + \frac{1}{2} S_{12}^2 \right) \\ &\left[ \frac{R_0^2}{3} - \frac{r_0^2}{2} - \frac{2r_0^2}{3\pi} - \frac{35r_0^2}{4\pi} (C_{A2c} - C_{A2d} - C_{A2e} + C_{A2f}) \right] + O(R_0^4) \end{aligned} \quad (3.2.27)$$

Next, we evaluate  $I_{2a}^{zzout}$ . For the outer part of the finite nucleus, we use the asymptotic expansion of the molecular incomplete gamma function given by Eq. (3.1.22). Substituting Eq. (3.1.22) into (3.2.20), we have

$$I_{2a}^{zzout} = \frac{2}{3} \int_{R_0}^{\infty} dr_M \frac{1}{r_M^2} \exp(-S_{12}r_M^2) i_0(2S_{12} \overline{MP} r_M) = \frac{2}{3} \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(3/2)_j} I_{2b}^{zzout} \quad (3.2.28)$$

where

$$I_{2b}^{zzout} = \int_{R_0}^{\infty} dr_M \ r_M^{2j-3} \exp(-S_{12}r_M^2) = \frac{1}{2} \int_{R_0^2}^{\infty} dx \ x^{j-2} \exp(-S_{12}x)$$

$$= \frac{\delta_{j0}}{2} \int_{R_0^2}^{\infty} dx \frac{1}{x^2} \exp(-S_{12}x) + \frac{\delta_{j1}}{2} \int_{R_0^2}^{\infty} dx \frac{1}{x} \exp(-S_{12}x) + \frac{1-\delta_{j0}-\delta_{j1}}{2} \int_{R_0^2}^{\infty} dx x^{j-2} \exp(-S_{12}x) \quad (3.2.29)$$

Substituting Eq. (3.2.26) into (3.2.25), we have

$$I_{2a}^{zzout} = \frac{S_{12}}{3} \Gamma(-1, S_{12}R_0^2) + \frac{2}{9} S_{12}^2 \overline{MP}^2 \Gamma(0, S_{12}R_0^2) + \frac{4}{45} S_{12}^4 \overline{MP}^4 \sum_{j'=0}^{\infty} \frac{(S_{12}^2 \overline{MP}^2)^{j'}}{(3)_{j'}(7/2)_{j'}} \left(\frac{1}{S_{12}}\right)^{j'+1} \Gamma(j'+1, S_{12}R_0^2) \quad (3.2.30)$$

We use the formula number 8.352.8 of Gradshteyn and Ryzhik [23] as given by

$$\Gamma(-n+k, x) = \frac{(-1)^{n-k}}{(n-k)!} \left[ \Gamma(0, x) - \exp(-x) \sum_{m=0}^{n-k-1} (-1)^m \frac{m!}{x^{m+1}} \right] \quad (3.2.31)$$

Thus, we have

$$\Gamma(-1, S_{12}R_0^2) = E_i(-S_{12}R_0^2) + \frac{\exp(-S_{12}R_0^2)}{S_{12}R_0^2} = \gamma + \ln(S_{12}R_0^2) - S_{12}R_0^2 + \frac{1}{S_{12}R_0^2} - 1 + \frac{1}{2}S_{12}R_0^2 + O(R_0^4) \quad (3.2.32)$$

Substituting Eq. (3.2.32), (3.1.31), and (3.1.32) into (3.2.30), we have

$$I_{2a}^{zzout} = \frac{1}{3R_0^2} - \frac{1}{3} \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) [\gamma + \ln(S_{12}R_0^2)] - \frac{S_{12}}{3} + \frac{4}{45} S_{12}^3 \overline{MP}^4 {}_2F_2 \left( 1, 1; 3, \frac{7}{2}; S_{12} \overline{MP}^2 \right) - \frac{R_0^2}{3} \left( \frac{4}{15} S_{12}^4 \overline{MP}^4 - \frac{2}{3} S_{12}^3 \overline{MP}^2 + \frac{1}{2} S_{12}^2 \right) + O(R_0^4) \quad (3.2.33)$$

Substituting Eq. (3.2.27) and (3.2.33) into (3.2.18), we have

$$I_{2a}^{zz} = \frac{C_{A0}}{r_0^2} + \frac{1}{3} \left( \frac{2}{3} S_{12}^2 \overline{MP}^2 - S_{12} \right) \left[ C_{A1} + \ln(S_{12}R_0^2) - \frac{S_{12}}{3} + \frac{4}{45} S_{12}^3 \overline{MP}^4 {}_2F_2 \left( 1, 1; 3, \frac{7}{2}; S_{12} \overline{MP}^2 \right) \right] + \left( \frac{4}{15} S_{12}^4 \overline{MP}^4 - \frac{2}{3} S_{12}^3 \overline{MP}^2 + \frac{1}{2} S_{12}^2 \right) r_0^2 C_{A2} + O(R_0^4) \quad (3.2.34)$$

where

$$C_{A0} = \frac{2}{3} - \frac{4}{3\pi} - \frac{1}{\pi} (C_{A0a} - C_{A0b} - C_{A0c} + C_{A0d}) = 0.2222222222 \quad (3.2.35)$$

$$C_{A1} = \gamma - \frac{2b^2}{\sqrt{\pi}} C_{A1a} + \frac{8}{\sqrt{\pi}} C_{A1b} - \frac{15}{2\pi} (C_{A1c} - C_{A1d}) = -2.287894210 \quad (3.2.36)$$

and

$$C_{A2} = -\frac{1}{2} - \frac{2}{3\pi} - \frac{35}{4\pi} (C_{A2c} - C_{A2d} - C_{A2e} + C_{A2f}) = -0.7360218971 \quad (3.2.37)$$

Next, we evaluate  $I_{3a}^{zz}$  as follows: We have

$$I_{3a}^{zz} = I_{3a}^{zzin} + I_{3a}^{zzout} \quad (3.2.38)$$

where

$$I_{3a}^{zzin} = \frac{-64}{3\pi^{2/2} r_0^7} \frac{S_{20}(\overline{MP})}{\overline{MP}^2} \int_0^{R_0} dr_M r_M^4 F_0 F_1 F_1 \exp(-S_{12}r_M^2) i_2(2S_{12} \overline{MP} r_M), \quad (3.2.39)$$

and

$$I_{3a}^{zzout} = \frac{-64}{3\pi^{2/2} r_0^7} \frac{S_{20}(\overline{MP})}{\overline{MP}^2} \int_{R_0}^{\infty} dr_M r_M^4 F_0 F_1 F_1 \exp(-S_{12}r_M^2) i_2(2S_{12} \overline{MP} r_M). \quad (3.2.40)$$

With a similar derivation to that for  $I_{2a}^{zzin}$ , we have

$$I_{3a}^{zzin} = S_{12}^2 S_{20}(\overline{MP}) \left\{ -\frac{8b^3}{45\sqrt{\pi}} \int_0^1 du \int_0^1 dv \sqrt{uv} \exp(-b^2 uv) + \frac{32}{45\sqrt{\pi}} \int_0^1 du \int_0^1 dv \frac{\sqrt{uv}}{(1+uv)^3} + \frac{2}{3\pi} \int_0^1 ds \int_0^1 du \int_0^1 dv \frac{\sqrt{uv}}{(1+s+uv)^{7/2}} \right.$$

$$\begin{aligned}
& -\frac{2}{3\pi} \int_0^1 ds \int_0^1 du \int_0^1 dv \frac{s^2 \sqrt{uv}}{(1+s+su)^{7/2}} + \left( \frac{2}{7} S_{12}^2 \overline{MP}^2 - S_{12} \right) \left[ -\frac{4R_0^2}{45} + \frac{2r_0^2}{15} + \frac{8r_0^2}{45\pi} \right. \\
& + \frac{7r_0^2}{3\pi} \int_0^1 ds \int_0^1 du \frac{\sqrt{su}}{(1+s+u)^{9/2}} - \frac{7r_0^2}{3\pi} \int_0^1 ds \int_0^1 du \frac{\sqrt{su}^{3/2}}{(1+s+u)^{9/2}} \\
& \left. - \frac{7r_0^2}{3\pi} \int_0^1 ds \int_0^1 du \frac{s^3 \sqrt{u}}{(1+s+su)^{9/2}} + \frac{7r_0^2}{3\pi} \int_0^1 ds \int_0^1 du \frac{s^3 u^{3/2}}{(1+s+su)^{9/2}} \right] \} + O(R_0^4) \quad (3.2.41)
\end{aligned}$$

All integrals in the above Eq. (3.2.41) are constants as given by Eq. (a.36), (a.37), (a.46), (a.47), (a.48), and (a.49). Substituting these constants into Eq. (3.2.41), we have

$$\begin{aligned}
I_{3a}^{zzin} &= S_{12}^2 S_{20}(\overline{MP}) \left\{ \frac{-8b^3}{45\sqrt{\pi}} C_{A1a} + \frac{32}{45\sqrt{\pi}} C_{A1b} + \frac{2}{3\pi} (C_{A1c} - C_{A1d}) \right. \\
& + \left. \left( \frac{2}{7} S_{12}^2 \overline{MP}^2 - S_{12} \right) \left[ -\frac{4R_0^2}{45} + \frac{2r_0^2}{15} + \frac{8r_0^2}{45\pi} + \frac{7r_0^2}{3\pi} (C_{A2c} - C_{A2d} - C_{A2e} + C_{A2f}) \right] \right\} + O(R_0^4) \quad (3.2.42)
\end{aligned}$$

For the term  $I_{3a}^{zzout}$ , with a similar derivation to that for  $I_{2a}^{zzout}$ , we have

$$\begin{aligned}
I_{3a}^{zzout} &= S_{12}^2 S_{20}(\overline{MP}) \left\{ \frac{4}{45} [\gamma + \ln(S_{12} R_0^2)] - \frac{8}{315} S_{12} \overline{MP}^2 {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; S_{12} \overline{MP}^2 \right) \right. \\
& \left. + \frac{4R_0^2}{45} \left( \frac{2}{7} S_{12}^2 \overline{MP}^2 - S_{12} \right) + O(R_0^4) \right\} \quad (3.2.43)
\end{aligned}$$

Substituting Eq. (3.2.42) and (3.2.43) into (3.2.38), we have

$$\begin{aligned}
I_{3a}^{zz} &= S_{12}^2 S_{20}(\overline{MP}) \left\{ \frac{4}{45} [\ln(S_{12} R_0^2) + C_{A1}] - \frac{8}{315} S_{12} \overline{MP}^2 {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; S_{12} \overline{MP}^2 \right) \right. \\
& \left. + \left( \frac{2}{7} S_{12}^2 \overline{MP}^2 - S_{12} \right) r_0^2 C_{A3} \right\} + O(R_0^4) \quad (3.2.44)
\end{aligned}$$

where

$$C_{A3} = \frac{2}{15} + \frac{8}{45\pi} + \frac{7}{3\pi} (C_{A2c} - C_{A2d} - C_{A2e} + C_{A2f}) = 0.1962725059 \quad (3.2.45)$$

Substituting Eq. (3.2.34) into (3.2.11), doing (3.2.44) into (3.2.12), and doing these resulting equations into (3.2.8), we have

$$\begin{aligned}
I_1^{zz} &= 4\pi \left\{ \frac{C_{A0}}{r_0^2} + \left( \frac{4}{45} S_{12}^2 S_{20}(\overline{MP}) + \frac{2}{9} S_{12}^2 \overline{MP}^2 - \frac{1}{3} S_{12} \right) [\ln(S_{12} R_0^2) + C_{A1}] \right. \\
& - \frac{S_{12}}{3} + \frac{4}{45} S_{12}^3 \overline{MP}^4 {}_2F_2 \left( 1, 1; 3, \frac{7}{2}; S_{12} \overline{MP}^2 \right) - \frac{8}{315} S_{12}^3 \overline{MP}^2 S_{20}(\overline{MP}) {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; S_{12} \overline{MP}^2 \right) \\
& \left. + \left( \frac{4}{15} S_{12}^4 \overline{MP}^4 - \frac{2}{3} S_{12}^3 \overline{MP}^2 + \frac{1}{2} S_{12}^2 \right) r_0^2 C_{A2} + \left( \frac{2}{7} S_{12}^2 \overline{MP}^2 - S_{12} \right) r_0^2 C_{A3} \right\} + O(R_0^4) \quad (3.2.46)
\end{aligned}$$

Substituting Eq. (3.2.46) into (3.2.5), we have

$$\begin{aligned}
I_{zz} &= \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{\Gamma(1+\varepsilon_A) \Gamma(1+\varepsilon_B)} \int_0^\infty dS_1 \int_0^\infty dS_2 (S_1 S_2)^{-3/2} f_0 f_{0B} \\
& \left\{ \frac{C_{A0}}{r_0^2} + \left( \frac{4}{45} S_{12}^2 S_{20}(\overline{MP}) + \frac{2}{9} S_{12}^2 \overline{MP}^2 - \frac{1}{3} S_{12} \right) [\ln(S_{12} R_0^2) + C_{A1}] \right. \\
& - \frac{S_{12}}{3} + \frac{4}{45} S_{12}^3 \overline{MP}^4 {}_2F_2 \left( 1, 1; 3, \frac{7}{2}; S_{12} \overline{MP}^2 \right) - \frac{8}{315} S_{12}^3 \overline{MP}^2 S_{20}(\overline{MP}) {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; S_{12} \overline{MP}^2 \right) \\
& \left. + \left( \frac{4}{15} S_{12}^4 \overline{MP}^4 - \frac{2}{3} S_{12}^3 \overline{MP}^2 + \frac{1}{2} S_{12}^2 \right) r_0^2 C_{A2} + \left( \frac{2}{7} S_{12}^2 \overline{MP}^2 - S_{12} \right) r_0^2 C_{A3} \right\} + O(R_0^4) \quad (3.2.47)
\end{aligned}$$

We evaluate the remaining integrals by the numerical integration. To do this, we first change integral variables

as follows: We set  $S_{12} = z$  and  $\frac{S_1}{S_{12}} = w$ . The Jacobian is given by Eq. (3.1.41). Further, we separate integral over  $z$  as is same as for Eq. (3.1.43). Then, we have the final formula given by

$$\begin{aligned}
 I_{zz} = & \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{\Gamma(1+\varepsilon_A)\Gamma(1+\varepsilon_B)a^2} \int_0^1 dw [w(1-w)]^{-3/2} \left\{ \int_0^1 du \exp[-wua^2 \overline{MA}^2 - (1-w)ua^2 \overline{MB}^2] f_0^{(1)} f_{0B}^{(1)} \right. \\
 & \left[ \frac{C_{A0}}{u^2 r_0^2} + \left( \frac{4}{45} a^4 y_0 - \frac{2}{9} a^4 x_0 + \frac{a^2}{3u} \right) [\ln(ua^2 R_0^2) + C_{A1}] \right. \\
 & - \frac{a^2}{3u} + \frac{4}{45} ua^6 x_0^2 {}_2F_2 \left( 1, 1; 3, \frac{7}{2}; ua^2 x_0 \right) - \frac{8}{315} ua^6 x_0 y_0 {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; ua^2 x_0 \right) \\
 & + \left( \frac{4}{15} u^2 a^8 x_0^2 - \frac{2}{3} ua^6 x_0 + \frac{1}{2} a^4 \right) r_0^2 C_{A2} + \left( \frac{2}{7} u^2 a^8 x_0 y_0 - ua^6 y_0 \right) r_0^2 C_{A3} \left. \right] \\
 & + \int_0^1 du \exp \left[ -\frac{w}{u} a^2 \overline{MA}^2 - \frac{1-w}{u} a^2 \overline{MB}^2 \right] f_0^{(2)} f_{0B}^{(2)} \left[ \frac{C_{A0}}{r_0^2} + \left( \frac{4}{45} \frac{a^4}{u^2} y_0 - \frac{2}{9} \frac{a^4}{u^2} x_0 + \frac{a^2}{3u} \right) \left[ \ln \left( \frac{a^2}{u} R_0^2 \right) + C_{A1} \right] \right. \\
 & - \frac{a^2}{3u} + \frac{4}{45} \frac{a^6}{u^3} x_0^2 {}_2F_2 \left( 1, 1; 3, \frac{7}{2}; ua^2 x_0 \right) - \frac{8}{315} \frac{a^6}{u^3} x_0 y_0 {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; ua^2 x_0 \right) \\
 & + \left( \frac{4}{15} \frac{a^8}{u^4} x_0^2 - \frac{2}{3} \frac{a^6}{u^3} x_0 + \frac{1}{2} \frac{a^4}{u^2} \right) r_0^2 C_{A2} + \left( \frac{2}{7} \frac{a^8}{u^4} x_0 y_0 - \frac{a^6}{u^3} y_0 \right) r_0^2 C_{A3} \left. \right] \left. \right\} + O(R_0^4) \quad (3.2.48)
 \end{aligned}$$

where  $f_0^{(1)}$  is given by Eq. (3.1.45),  $f_0^{(2)}$  is given by (3.1.52),  $x_0$  is given by (3.1.59),  $y_0$  is given by (3.1.60),  $C_{A0}$  is given by (3.2.35),  $C_{A1}$  is given by (3.2.36),  $C_{A2}$  is given by (3.2.37),  $C_{A3}$  is given by (3.2.45), and  $f_{0B}^{(1)}$  and  $f_{0B}^{(2)}$  are given by

$$f_{0B}^{(1)} = \left[ \frac{\zeta_B^2}{2(1-w)ua^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right] \exp \left[ -\frac{\zeta_B^2}{4(1-w)ua^2 t_2^2} \right] \quad (3.2.49)$$

and

$$f_{0B}^{(2)} = \left[ \frac{u\zeta_B^2}{2(1-w)a^2} \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{4+\varepsilon_B}} - \int_0^1 dt_2 \frac{(1-t_2)^{\varepsilon_B}}{t_2^{2+\varepsilon_B}} \right] \exp \left[ -\frac{u\zeta_B^2}{4(1-w)a^2 t_2^2} \right] \quad (3.2.50)$$

For  $I_{yy}$ , we have it by replacing  $y_0$  by  $-\frac{1}{2}(y_0 + \sqrt{3}y_2)$  in  $I_{zz}$ , Eq. (3.2.48), where  $y_m$  is given by

$$y_m = w^2 S_{2m}(\overrightarrow{MA}) + (1-w)^2 S_{2m}(\overrightarrow{MB}) + w(1-w) S_{2m}(\overrightarrow{MA}, \overrightarrow{MB}; 1) \quad (3.2.51)$$

in which  $S_{2m}(\overrightarrow{MA}, \overrightarrow{MB}; 1)$  is the mixed solid harmonics defined in a previous article [28] as given by

$$S_{2m}(\overrightarrow{MA}, \overrightarrow{MB}; 1) = 2MA_z MB_z - (MA_x MB_x + MA_y MB_y) \quad (3.2.52)$$

For  $I_{xx}$ , we have it by replacing  $y_0$  by  $-\frac{1}{2}(y_0 - \sqrt{3}y_2)$  in  $I_{zz}$ , Eq. (3.2.48). For  $I_{xy}$ , a similar derivation to that for

$I_{zz}$ , we have the final formula given by

$$\begin{aligned}
 I_{xy} = & \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{1+\varepsilon_B}}{\Gamma(1+\varepsilon_A)\Gamma(1+\varepsilon_B)a^2} \int_0^1 dw [w(1-w)]^{-3/2} a^2 [wMA_x + (1-w)MB_x][wMA_y + (1-w)MB_y] \\
 & \left\{ \int_0^1 du \exp[-wua^2 \overline{MA}^2 - (1-w)ua^2 \overline{MB}^2] f_0^{(1)} f_{0B}^{(1)} \right. \\
 & \left[ \frac{2}{15} \ln(ua^2 R_0^2) + \frac{2}{15} C_{A1} - \frac{4}{105} ua^2 x_0 {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; ua^2 x_0 \right) + \left( \frac{2}{7} u^2 a^4 x_0 - ua^2 \right) r_0^2 C_{A3} \right. \\
 & + \int_0^1 du \exp \left[ -\frac{w}{u} a^2 \overline{MA}^2 - \frac{1-w}{u} a^2 \overline{MB}^2 \right] f_0^{(2)} f_{0B}^{(2)} \left. \right. \left. \right\} \quad (3.2.48)
 \end{aligned}$$

$$\left[ \frac{2}{15u^2} \ln \left( \frac{a^2}{u} R_0^2 \right) + \frac{2}{15u^2} C_{A1} - \frac{4}{105u^3} x_0 {}_2F_2 \left( 1, 1; 2, \frac{9}{2}; \frac{a^2}{u} x_0 \right) \left( \frac{2a^4}{7u^4} x_0 - \frac{a^2}{u^5} \right) r_0^2 C_{A3} \right] \} + O(R_0^4) \quad (3.2.53)$$

The  $I_{yz}$  can be written as replacing  $[wMA_x + (1-w)MB_x]$  by  $[wMA_z + (1-w)MB_z]$  in  $I_{xy}$ , Eq. (3.2.53).

The  $I_{zx}$  can be done as replacing  $[wMA_y + (1-w)MB_y]$  by  $[wMA_z + (1-w)MB_z]$  in  $I_{xy}$ . Of course,  $I_{\eta\xi} = I_{\xi\eta}$ .

The calculation of  ${}_2F_2$  in Eq. (3.2.48) can be described in Appendix B. Using the 256-point GLQ, we have

the value of  $I_{zz}$ ,  $I_{yy}$ ,  $I_{xx}$ , and  $I_{zx}$  with 7 significant-figure precision and  $I_{xy}$  and  $I_{yz}$  with 5 significant-figure

precision. We obtain  $-\frac{e^4}{c^4} I_{zz} = -0.9808368(-1)$ ,  $-\frac{e^4}{c^4} I_{yy} = -0.9808368(-1)$ ,

$$-\frac{e^4}{c^4} I_{xx} = -0.9808368(-1), \quad -\frac{e^4}{c^4} I_{xy} = -0.32330(-12), \quad -\frac{e^4}{c^4} I_{yz} = 0.22861(-12),$$

and  $-\frac{e^4}{c^4} I_{zx} = -0.1101417(-9)$  for the case of three hydrogen atoms located at  $\vec{M} = (0, 0, 0)$ ,  $\vec{A} = (-\frac{\sqrt{8}}{3}, -\sqrt{\frac{8}{3}}, \frac{2}{3})$ ,

$$\text{and } \vec{B} = (-\frac{\sqrt{8}}{3}, \sqrt{\frac{8}{3}}, \frac{2}{3}).$$

Note that each main term of  $I_{zz}$ ,  $I_{yy}$ , and  $I_{xx}$  is  $C_{A0}/r_0^2$ , which is very large. As the results, each value of  $I_{yy}$  and  $I_{xx}$  is the same as that of  $I_{zz}$ . Also note that each of  $I_{xy}$ ,  $I_{yz}$ , and  $I_{zx}$  has not that main term. As the results, each

value of  $I_{xy}$ ,  $I_{yz}$ , and  $I_{zx}$  is very small comparing with that of  $I_{zz}$ .

### 3.3 The Term $\vec{p} \cdot V\vec{A} + \vec{A} \cdot V\vec{p}$

We evaluate the molecular integral-formula over Dirac wave functions for the term  $\vec{p} \cdot V\vec{A} + \vec{A} \cdot V\vec{p}$  with potentials of the finite nucleus for the GCDD model as follows: We can easily derive the following relation:

$$\vec{p} \cdot V\vec{A} = \vec{A} \cdot V\vec{p} \quad (3.3.1)$$

Thus, we have

$$\langle \chi_{\mu A} | \vec{p} \cdot V\vec{A} + \vec{A} \cdot V\vec{p} | \chi_{\nu B} \rangle = 2 \langle \chi_{\mu A} | \vec{A} \cdot V\vec{p} | \chi_{\nu B} \rangle = \frac{iZ_M^2 e^3 \hbar}{c^2} \sum_{\xi} \mu_{M\xi} I_{\xi} \quad [\xi \in (x, y, z)] \quad (3.3.2)$$

where

$$I_{\xi} = \frac{16}{\pi r_0^4} \int d\vec{r}_M (\vec{B}\vec{M} \times \vec{r}_M)_{\xi} F_0 F_1 r_A^{-\varepsilon_A} \left( \frac{\varepsilon_B}{r_B^{2+\varepsilon_B}} + \frac{\zeta_B}{r_B^{1+\varepsilon_B}} \right) \exp(-\zeta_A r_A - \zeta_B r_B) \quad (3.3.3)$$

First, we evaluate  $I_z$  as given by

$$I_z = \frac{16}{\pi r_0^4} \int d\vec{r}_M (\vec{B}\vec{M} \times \vec{r}_M)_z F_0 F_1 r_A^{-\varepsilon_A} \left( \frac{\varepsilon_B}{r_B^{2+\varepsilon_B}} + \frac{\zeta_B}{r_B^{1+\varepsilon_B}} \right) \exp(-\zeta_A r_A - \zeta_B r_B) \quad (3.3.4)$$

We use Eq. (3.2.4) and the relation derived in a previous article [20] given by

$$\left( \frac{\varepsilon_B}{r_B^{2+\varepsilon_B}} + \frac{\zeta_B}{r_B^{1+\varepsilon_B}} \right) \exp(-\zeta_B r_B) = \frac{\zeta_B^{2+\varepsilon_B}}{2\sqrt{\pi} \Gamma(2+\varepsilon_B)} \int_0^{\infty} dS_2 S_2^{-3/2} \exp(-S_2 r_B^2) f_2 \quad (3.3.5)$$

where  $f_2$  is given by (3.1.6). We use the Gaussian product rule, Eq. (3.1.11), the Sack's formula, Eq. (3.1.12), and Eq. (3.1.14). Then, we have

$$I_z = \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{3+\varepsilon_B}}{\Gamma(1+\varepsilon_A) \Gamma(2+\varepsilon_B)} \int_0^{\infty} dS_1 \int_0^{\infty} dS_2 (S_1 S_2)^{-3/2} \exp[-S_1 \vec{M}\vec{A}^2 - S_2 \vec{M}\vec{B}^2] f_0 f_2 I_1^z \quad (3.3.6)$$

where  $f_0$  is given by Eq. (3.1.4) and

$$I_1^z = \frac{16}{\pi r_0^4} \int_0^{\infty} dr_M r_M^2 F_0 F_1 \exp[-S_{12} r_M^2] \sum_{l=0}^{\infty} i_l (2S_{12} \vec{M}\vec{P} r_M) \int d\vec{r}_M \sum_{m=-l}^l Y_l^m(\vec{M}\vec{P}) Y_l^m(\vec{r}_M)^* (\vec{B}\vec{M} \times \vec{r}_M)_z \quad (3.3.7)$$

The angular part can be evaluated as in a previous article [24] as given by

$$\int d\hat{r}_M \sum_{m=-l}^l Y_l^m(\vec{MP}) Y_l^m(\hat{r}_M)^* (\vec{BM} \times \vec{r}_M)_z = r_M \delta_{l1} \frac{s_1}{s_{12}} \frac{(\vec{BM} \times \vec{MA})_z}{\vec{MP}} = r_M \delta_{l1} \frac{s_1}{s_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \quad (3.3.8)$$

Then we have

$$I_1^z = \frac{16}{\pi r_0^4} \frac{s_1}{s_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \int_0^\infty dr_M r_M^3 F_0 F_1 \exp[-S_{12} r_M^2] i_1(2S_{12} \vec{MP} r_M) = I_1^{zin} + I_1^{zout} \quad (3.3.9)$$

where

$$I_1^{zin} = \frac{16}{\pi r_0^4} \frac{s_1}{s_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \int_0^{R_0} dr_M r_M^3 F_0 F_1 \exp[-S_{12} r_M^2] i_1(2S_{12} \vec{MP} r_M) \quad (3.3.10)$$

and

$$I_1^{zout} = \frac{16}{\pi r_0^4} \frac{s_1}{s_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \int_{R_0}^\infty dr_M r_M^3 F_0 F_1 \exp[-S_{12} r_M^2] i_1(2S_{12} \vec{MP} r_M) \quad (3.3.11)$$

First, we evaluate  $I_1^{zin}$ . We use the power series for the molecular incomplete gamma function, Eq. (3.1.21), and have

$$\begin{aligned} I_1^{zin} &= \frac{16}{3\pi r_0^4} \frac{s_1}{s_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \sum_{n_1, n_2=0} \frac{(-1/r_0^2)^{n_{12}} (3/2)_{n_1} (1/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (3/2)_{n_2}} \int_0^{R_0} dr_M r_M^{2n_{12}+3} \exp[-S_{12} r_M^2] i_1(2S_{12} \vec{MP} r_M) \\ &= \frac{16}{3\pi r_0^4} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \frac{2S_{12} \vec{MP}}{3} \sum_{n_1, n_2=0} \frac{(-1/r_0^2)^{n_{12}} (3/2)_{n_1} (1/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (3/2)_{n_2}} \sum_{j=0} \frac{(S_{12} \vec{MP}^2)^j}{j! (5/2)_j} I_{1a}^{zin} \end{aligned} \quad (3.3.12)$$

where we use Eq. (3.1.13) and

$$\begin{aligned} I_{1a}^{zin} &= \int_0^{R_0} dr_M r_M^{2j+2n_{12}+4} \exp[-S_{12} r_M^2] = \frac{1}{2} \int_0^{R_0^2} dx x^{j+n_{12}+3/2} \exp(-S_{12} x) \\ &= \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+n_{12}+5/2} \gamma \left( j+n_{12} + \frac{3}{2}; S_{12} R_0^2 \right) = \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+n_{12}+5/2} (S_{12} R_0^2)^{j+n_{12}+5/2} \frac{\Gamma(j+n_{12} + 5/2)}{\Gamma(j+n_{12} + 7/2)} \\ &{}_1F_1 \left( j+n_{12} + \frac{5}{2}; j+n_{12} + \frac{7}{2}; -S_{12} R_0^2 \right) = \frac{1}{2} (R_0^2)^{j+n_{12}+5/2} \frac{\Gamma(j+n_{12} + 5/2)}{\Gamma(j+n_{12} + 7/2)} {}_1F_1 \left( j+n_{12} + \frac{5}{2}; j+n_{12} + \frac{7}{2}; -S_{12} R_0^2 \right) \end{aligned} \quad (3.3.13)$$

where we use Eq. (3.1.25). Substituting Eq. (3.3.13) into (3.3.12), we have

$$\begin{aligned} I_1^{zin} &= \frac{16b^5 r_0}{9\pi} S_1 (\vec{MA} \times \vec{MB})_z \sum_{n_1, n_2=0} \frac{(-b^2)^{n_{12}} (3/2)_{n_1} (1/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (3/2)_{n_2}} \\ &\quad \sum_{j=0} \frac{(S_{12} \vec{MP}^2 R_0^2)^j}{j! (5/2)_j} \frac{\Gamma(j+n_{12} + 5/2)}{\Gamma(j+n_{12} + 7/2)} {}_1F_1 \left( j+n_{12} + \frac{5}{2}; j+n_{12} + \frac{7}{2}; -S_{12} R_0^2 \right) \\ &= \frac{16b^5 r_0}{9\pi} S_1 (\vec{MA} \times \vec{MB})_z \left\{ S_{B1} + \left[ \frac{2}{5} S_{12}^2 \vec{MP}^2 - S_{12} \right] R_0^2 S_{B2} + O(R_0^4) \right\} \end{aligned} \quad (3.3.14)$$

where

$$S_{B1} = \sum_{n_1, n_2=0} \frac{(-b^2)^{n_{12}} (3/2)_{n_1} (1/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (3/2)_{n_2}} \frac{\Gamma(n_{12} + 5/2)}{\Gamma(n_{12} + 7/2)} \quad (3.3.15)$$

and

$$S_{B2} = \sum_{n_1, n_2=0} \frac{(-b^2)^{n_{12}} (3/2)_{n_1} (1/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (3/2)_{n_2}} \frac{\Gamma(n_{12} + 7/2)}{\Gamma(n_{12} + 9/2)} \quad (3.3.16)$$

Using a similar derivation to that in Appendix A for these summations, we have

$$S_{B1} = \frac{3\pi}{4b^4} - \frac{9\sqrt{2\pi}}{8b^5} \quad (3.3.17)$$

and

$$S_{B2} = \frac{\pi}{4b^3} - \frac{25\sqrt{\pi}}{16\sqrt{2}b^7} \quad (3.3.18)$$

Thus, we have

$$I_1^{zin} = S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \left\{ \frac{4}{3} R_0 - \sqrt{\frac{2}{\pi}} r_0 + \left[ \frac{2}{5} S_{12}^2 \overline{MP}^2 - S_{12} \right] \left( \frac{4}{9} R_0^3 - \frac{25r_0^3}{9\sqrt{2\pi}} \right) \right\} + O(R_0^5) \quad (3.3.19)$$

Next, we evaluate  $I_1^{zout}$ . For the outer part of the finite nucleus, we use the asymptotic expansion, Eq. (3.1.22), and have

$$I_1^{zout} = 2 \frac{S_1}{S_{12}} \frac{(\overrightarrow{MA} \times \overrightarrow{MB})_z}{\overline{MP}} \int_{R_0}^{\infty} dr_M \frac{1}{r_M} \exp[-S_{12}r_M^2] i_1(2S_{12}\overline{MP}r_M) = 2 \frac{(\overrightarrow{MA} \times \overrightarrow{MB})_z}{\overline{MP}} \frac{2S_1\overline{MP}}{3} \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(5/2)_j} I_{1a}^{zout} \quad (3.3.20)$$

where

$$I_{1a}^{zout} = \int_{R_0}^{\infty} dr_M r_M^{2j} \exp(-S_{12}r_M^2) = \frac{1}{2} \int_{R_0^2}^{\infty} dx x^{j-1/2} \exp(-S_{12}x) = \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+1/2} \Gamma \left( j + \frac{1}{2}; S_{12}R_0^2 \right) \quad (3.3.21)$$

It is easy to derive the following relation

$$\Gamma \left( j + \frac{1}{2}; S_{12}R_0^2 \right) = \Gamma \left( j + \frac{1}{2} \right) - \frac{(S_{12}R_0^2)^{j+1/2}}{j+1/2} + \frac{(S_{12}R_0^2)^{j+3/2}}{j+3/2} + O(R_0^5) \quad (3.3.22)$$

Substituting Eq. (3.3.22) into (3.3.21) and doing the resulting equation into (3.3.20), we have

$$I_1^{zout} = S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \left\{ \frac{2}{3} \frac{\sqrt{\pi}}{\sqrt{S_{12}}} {}_1F_1 \left( \frac{1}{2}; \frac{5}{2}; S_{12}\overline{MP}^2 \right) - \frac{4}{3} R_0 - \left[ \frac{2}{5} S_{12}^2 \overline{MP}^2 - S_{12} \right] \frac{4}{9} R_0^3 \right\} + O(R_0^5) \quad (3.3.23)$$

Substituting Eq. (3.3.19) and (3.3.23) into (3.3.9) and doing the resulting equation into (3.3.5), we have

$$I_z = \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{3+\varepsilon_B}}{\Gamma(1+\varepsilon_A)\Gamma(2+\varepsilon_B)} \int_0^{\infty} dS_1 \int_0^{\infty} dS_2 (S_1 S_2)^{-3/2} \exp[-S_1 \overline{MA}^2 - S_2 \overline{MB}^2] f_0 f_2 S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \left\{ \frac{2}{3} \frac{\sqrt{\pi}}{\sqrt{S_{12}}} {}_1F_1 \left( \frac{1}{2}; \frac{5}{2}; S_{12}\overline{MP}^2 \right) - \frac{\sqrt{2}}{\sqrt{\pi}} r_0 - \left[ \frac{2}{5} S_{12}^2 \overline{MP}^2 - S_{12} \right] \frac{25r_0^3}{9\sqrt{2\pi}} \right\} + O(R_0^5) \quad (3.3.24)$$

We evaluate the remaining integrals by the numerical integration. To do this, we first change integral variables as follows: We set  $S_{12} = z$  and  $\frac{S_1}{S_{12}} = w$ . The Jacobian is given by Eq. (3.1.41). Further, we separate integral over  $z$  as is same as for Eq. (3.1.43). Then, we have the final formula given by

$$I_z = (\overrightarrow{MA} \times \overrightarrow{MB})_z \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{3+\varepsilon_B}}{\Gamma(1+\varepsilon_A)\Gamma(2+\varepsilon_B)} \int_0^1 dw w^{-1/2} (1-w)^{-3/2} \left\{ \int_0^1 du \exp[-wua^2 \overline{MA}^2 - (1-w)ua^2 \overline{MB}^2] f_0^{(1)} f_2^{(1)} \right. \\ \left[ \frac{2}{3} \frac{\sqrt{\pi}}{au^{3/2}} {}_1F_1 \left( \frac{1}{2}; \frac{5}{2}; ua^2 x_0 \right) - \frac{\sqrt{2}r_0}{\sqrt{\pi}u} - \left[ \frac{2}{5} ua^2 x_0 - 1 \right] \frac{25a^2 r_0^3}{9\sqrt{2\pi}} \right] + \int_0^1 du \exp[-\frac{w}{u} a^2 \overline{MA}^2 - \frac{1-w}{u} a^2 \overline{MB}^2] f_0^{(2)} f_2^{(2)} \\ \left. \left[ \frac{2}{3} \frac{\sqrt{\pi}}{au^{1/2}} {}_1F_1 \left( \frac{1}{2}; \frac{5}{2}; \frac{a^2}{u} x_0 \right) - \frac{\sqrt{2}r_0}{\sqrt{\pi}u} - \left[ \frac{2}{5} \frac{a^2}{u} x_0 - 1 \right] \frac{25a^2 r_0^3}{9\sqrt{2\pi}u^2} \right] \right\} + O(R_0^5) \quad (3.3.25)$$

where  $f_0^{(1)}$  is given by Eq. (3.1.45),  $f_2^{(1)}$  is given by (3.1.47),  $x_0$  is given by (3.1.59),  $f_0^{(2)}$  is given by (3.1.52), and  $f_2^{(2)}$  is given by (3.1.54). Replacing  $(\overrightarrow{MA} \times \overrightarrow{MB})_z$  by  $(\overrightarrow{MA} \times \overrightarrow{MB})_y$  in Eq. (3.3.25), we have the final formula of  $I_y$ .

Doing that by  $(\overrightarrow{MA} \times \overrightarrow{MB})_x$  in (3.3.25), we have that of  $I_x$ . Using the 64-point GLQ, we have the value of  $I_z$ ,  $I_y$ , and  $I_x$  as given by  $\frac{ie^2\hbar}{c^2} I_z = -0.97838414(-5)i$ ,  $\frac{ie^2\hbar}{c^2} I_y = 0$ , and  $\frac{ie^2\hbar}{c^2} I_x = -0.69182206(-5)i$  for the case of

three hydrogen atoms located at  $\vec{M} = (0, 0, 0)$ ,  $\vec{A} = (-\frac{\sqrt{8}}{3}, -\sqrt{\frac{8}{3}}, \frac{2}{3})$ , and  $\vec{B} = (-\frac{\sqrt{8}}{3}, \sqrt{\frac{8}{3}}, \frac{2}{3})$ . Note that  $(\vec{MA} \times \vec{MB})_y = MA_z MB_x - MA_x MB_z = 0$ .

### 3.4 The Term $i\vec{\sigma} \cdot (\vec{p} \times \mathbf{V}\vec{p})$

We evaluate the molecular integral-formula over Dirac wave functions for the term  $i\vec{\sigma} \cdot (\vec{p} \times \mathbf{V}\vec{p})$  with the GCDD model of  $\mathbf{V}$  which is given by Eq. (1.7) as follows: It is easy to derive the following relation:

$$\langle \chi_{\mu A} | i\vec{\sigma} \cdot (\vec{p} \times V\vec{p}) | \chi_{\nu B} \rangle = iZ_M e^2 \hbar^2 \sum_{\xi} \sigma_{\xi} I_{\xi}^{\sigma} \quad [\xi \in (x, y, z)] \quad (3.4.1)$$

where

$$I_{\xi}^{\sigma} = \frac{4}{\sqrt{\pi} r_0^3} \int d\vec{r}_M F_1 r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) (\vec{r}_M \times \nabla)_{\xi} r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) \quad (3.4.2)$$

First, we evaluate  $I_z^{\sigma}$  as given by

$$I_z^{\sigma} = \frac{4}{\sqrt{\pi} r_0^3} \int d\vec{r}_M F_1 r_A^{-\varepsilon_A} \exp(-\zeta_A r_A) (\vec{r}_M \times \nabla)_z r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) \quad (3.4.3)$$

We use the Gaussian-transforms for the Dirac wave function, Eq. (3.2.4), and for the derivative of it as in a previous article [20] as given by

$$\nabla r_B^{-\varepsilon_B} \exp(-\zeta_B r_B) = \frac{-\vec{r}_B \zeta_B^{2+\varepsilon_B}}{2\sqrt{\pi} \Gamma(2+\varepsilon_B)} \int_0^{\infty} dS_2 \quad S_2^{-3/2} \exp(-S_2 r_B^2) f_2 \quad (3.4.4)$$

where  $f_2$  is given by Eq. (3.1.6). Then, we have

$$I_z^{\sigma} = \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{2+\varepsilon_B}}{4\pi \Gamma(1+\varepsilon_A) \Gamma(2+\varepsilon_B)} \int_0^{\infty} dS_1 \int_0^{\infty} dS_2 (S_1 S_2)^{-3/2} f_0 f_2 I_{z1}^{\sigma} \quad (3.4.5)$$

where  $f_0$  is given by Eq. (3.1.4) and

$$I_{z1}^{\sigma} = \frac{-4}{\sqrt{\pi} r_0^3} \int d\vec{r}_M (\vec{r}_M \times \vec{r}_B)_z F_1 \exp(-S_1 r_A^2 - S_2 r_B^2) \quad (3.4.6)$$

We know that  $\vec{r}_M \times \vec{r}_B = \vec{r}_M \times (\vec{r}_M + \vec{MB}) = \vec{r}_M \times \vec{MB} = -\vec{r}_M \times \vec{MB}$ . Then, we have

$$I_{z1}^{\sigma} = \frac{4}{\sqrt{\pi} r_0^3} \int d\vec{r}_M (\vec{r}_M \times \vec{MB})_z F_1 \exp(-S_1 r_A^2 - S_2 r_B^2) \quad (3.4.7)$$

Using the Gaussian product rule, Eq. (3.1.11), Sack's formula, (3.1.12), and (3.1.14), we have

$$I_{z1}^{\sigma} = 4\pi \exp(-S_1 \vec{MA}^2 - S_2 \vec{MB}^2) I_{z2}^{\sigma} \quad (3.4.8)$$

where

$$\begin{aligned} I_{z2}^{\sigma} &= \frac{4}{\sqrt{\pi} r_0^3} \int d\vec{r}_M (\vec{r}_M \times \vec{MB})_z F_1 \exp(-S_{12} r_M^2) \sum_{l=0}^l i_l (2S_{12} \vec{MP} r_M) \sum_{m=-l}^l Y_l^m(\vec{MP}) Y_l^m(\vec{r}_M)^* \\ &= \frac{4}{\sqrt{\pi} r_0^3} \int_0^{\infty} dr_M r_M^2 F_1 \exp(-S_{12} r_M^2) \sum_{l=0}^l i_l (2S_{12} \vec{MP} r_M) \int \widehat{r_M} \sum_{m=-l}^l Y_l^m(\vec{MP}) Y_l^m(\widehat{r_M})^* (\vec{r}_M \times \vec{MB})_z \end{aligned} \quad (3.4.9)$$

The angular part can be evaluated as in a previous article [24] as given by

$$\int \widehat{r_M} \sum_{m=-l}^l Y_l^m(\vec{MP}) Y_l^m(\widehat{r_M})^* (\vec{r}_M \times \vec{MB})_z = r_M \delta_{l1} \frac{(\vec{MP} \times \vec{MB})_z}{\vec{MP}} \quad (3.4.10)$$

We know  $\vec{MP} \times \vec{MB} = \frac{S_1}{S_{12}} \vec{MA} \times \vec{MB}$ . Thus, we have

$$I_{z2}^{\sigma} = \frac{4}{\sqrt{\pi} r_0^3} \frac{S_1}{S_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \int_0^{\infty} dr_M r_M^3 F_1 \exp(-S_{12} r_M^2) i_1 (2S_{12} \vec{MP} r_M) = I_{z2}^{\sigma in} + I_{z2}^{\sigma out} \quad (3.4.11)$$

where

$$I_{z2}^{\sigma in} = \frac{4}{\sqrt{\pi} r_0^3} \frac{S_1}{S_{12}} \frac{(\vec{MA} \times \vec{MB})_z}{\vec{MP}} \int_0^{R_0} dr_M r_M^3 F_1 \exp(-S_{12} r_M^2) i_1 (2S_{12} \vec{MP} r_M) \quad (3.4.12)$$

and

$$I_{z2}^{\sigma out} = \frac{4}{\sqrt{\pi} r_0^2} \frac{S_1}{S_{12}} \frac{(\overrightarrow{MA} \times \overrightarrow{MB})_z}{\overline{MP}} \int_{R_0}^{\infty} dr_M r_M^3 F_1 \exp(-S_{12} r_M^2) i_1(2S_{12} \overline{MP} r_M) \quad (3.4.13)$$

First, we evaluate  $I_{z2}^{\sigma in}$ . Using the power series of the molecular incomplete gamma function, Eq. (3.1.21), and doing Eq. (3.1.13), we have

$$I_{z2}^{\sigma in} = \frac{4}{3\sqrt{\pi} r_0^3} \frac{S_1}{S_{12}} \frac{(\overrightarrow{MA} \times \overrightarrow{MB})_z}{\overline{MP}} \frac{2S_{12} \overline{MP}}{3} \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(5/2)_j} \sum_{n=0} \frac{(-1/r_0^2)^n (3/2)_n}{n!(5/2)_n} I_{z2a}^{\sigma in} \quad (3.4.14)$$

where

$$\begin{aligned} I_{z2a}^{\sigma in} &= \int_0^{R_0} dr_M r_M^{2j+2n+4} \exp(-S_{12} r_M^2) = \frac{1}{2} \int_0^{R_0^2} dx x^{j+n+3/2} \exp(-S_{12} x) \\ &= \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+n+5/2} \gamma \left( j+n+\frac{5}{2}; S_{12} R_0^2 \right) \\ &= \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+n+5/2} (S_{12} R_0^2)^{j+n+5/2} \frac{\Gamma(j+n+5/2)}{\Gamma(j+n+7/2)} {}_1F_1 \left( j+n+\frac{5}{2}; j+n+\frac{7}{2}; -S_{12} R_0^2 \right) \end{aligned} \quad (3.4.15)$$

Substituting Eq. (3.4.15) into (3.4.14), we have

$$\begin{aligned} I_{z2}^{\sigma in} &= \frac{4b^5 r_0^2}{9\sqrt{\pi}} S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2 R_0^2)^j}{j!(5/2)_j} \sum_{n=0} \frac{(-b^2)^n (3/2)_n}{n!(5/2)_n} \\ &\quad \frac{\Gamma(j+n+5/2)}{\Gamma(j+n+7/2)} {}_1F_1 \left( j+n+\frac{5}{2}; j+n+\frac{7}{2}; -S_{12} R_0^2 \right) \\ &= \frac{4b^5 r_0^2}{9\sqrt{\pi}} S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \sum_{n=0} \frac{(-b^2)^n (3/2)_n}{n!(5/2)_n} \frac{\Gamma(n+5/2)}{\Gamma(n+7/2)} + O(R_0^4) \\ &= \frac{4b^5 r_0^2}{9\sqrt{\pi}} S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \frac{\Gamma(5/2)}{\Gamma(7/2)} {}_1F_1 \left( \frac{3}{2}; \frac{7}{2}; -b^2 \right) + O(R_0^4) = \frac{4b^5 r_0^2}{9\sqrt{\pi}} S_1 (\overrightarrow{MA} \times \overrightarrow{MB})_z \frac{3\sqrt{\pi}}{4b^3} \left( 1 - \frac{3}{2b^2} \right) + O(R_0^4) \end{aligned} \quad (3.4.16)$$

In the above derivation, we use the asymptotic expansion of the CHF given by Eq. (3.1.28). Next, we evaluate  $I_{z2}^{\sigma out}$ . Using the asymptotic expansion of  $F_1$ , Eq. (3.1.22), we have

$$I_{z2}^{\sigma out} = \frac{S_1}{S_{12}} \frac{(\overrightarrow{MA} \times \overrightarrow{MB})_z}{\overline{MP}} \int_{R_0}^{\infty} dr_M \exp(-S_{12} r_M^2) i_1(2S_{12} \overline{MP} r_M) \quad (3.4.17)$$

Using the power series of  $i_1$ , Eq. (3.1.13), we have

$$I_{z2}^{\sigma out} = \frac{S_1}{S_{12}} \frac{(\overrightarrow{MA} \times \overrightarrow{MB})_z}{\overline{MP}} \frac{2S_{12} \overline{MP}}{3} \sum_{j=0} \frac{(S_{12}^2 \overline{MP}^2)^j}{j!(5/2)_j} I_{z2a}^{\sigma out} \quad (3.4.18)$$

where

$$\begin{aligned} I_{z2a}^{\sigma out} &= \int_{R_0}^{\infty} dr_M r_M^{2j+1} \exp(-S_{12} r_M^2) = \frac{1}{2} \int_{R_0^2}^{\infty} dx x^j \exp(-S_{12} x) \\ &= \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+1} \Gamma(j+1; S_{12} R_0^2) = \frac{1}{2} \left( \frac{1}{S_{12}} \right)^{j+1} \left[ \Gamma(j+1) - \frac{(S_{12} R_0^2)^{j+1}}{j+1} \right] + O(R_0^4) \end{aligned} \quad (3.4.19)$$

In the above derivation, we use Eq. (3.1.32). Substituting Eq. (3.4.19) into (3.4.18), we have

$$I_{z2}^{\sigma out} = \frac{1}{3} (\overrightarrow{MA} \times \overrightarrow{MB})_z \left[ \frac{S_1}{S_{12}} {}_1F_1 \left( 1; \frac{5}{2}; S_{12} \overline{MP}^2 \right) - S_1 R_0^2 \right] + O(R_0^4) \quad (3.4.20)$$

Substituting Eq. (3.4.16) and (3.4.20) into (3.4.11), doing the resulting equation into (3.4.8), and doing it into (3.4.5), we have

$$\begin{aligned} I_z^{\sigma} &= \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{3+\varepsilon_B}}{\Gamma(1+\varepsilon_A) \Gamma(2+\varepsilon_B)} \int_0^{\infty} dS_1 \int_0^{\infty} dS_2 (S_1 S_2)^{-3/2} \exp(-S_1 \overline{MA}^2 - S_2 \overline{MB}^2) f_0 f_2 \\ &\quad (\overrightarrow{MA} \times \overrightarrow{MB})_z \left[ \frac{1}{3} \frac{S_1}{S_{12}} {}_1F_1 \left( 1; \frac{5}{2}; S_{12} \overline{MP}^2 \right) - \frac{1}{2} S_1 R_0^2 \right] + O(R_0^4) \end{aligned} \quad (3.4.21)$$

We evaluate the remaining integrals by the numerical integration. To do this, we first change integral variables as follows: We set  $S_{12} = z$  and  $\frac{S_1}{S_{12}} = w$ . The Jacobian is given by Eq. (3.1.41). Further, we separate integral over  $z$  as is same as for Eq. (3.1.43). Then, we have the final formula given by

$$I_z^\sigma = (\overrightarrow{MA} \times \overrightarrow{MB})_z \frac{\zeta_A^{1+\varepsilon_A} \zeta_B^{3+\varepsilon_B}}{\Gamma(1+\varepsilon_A)\Gamma(2+\varepsilon_B)} \int_0^1 dw w^{-1/2} (1-w)^{-3/2} \left\{ \int_0^1 du \exp[-wua^2 \overrightarrow{MA}^2 - (1-w)ua^2 \overrightarrow{MB}^2] f_0^{(1)} f_2^{(1)} \left[ \frac{1}{3u^2 a^2} {}_1F_1 \left( 1; \frac{5}{2}; ua^2 x_0 \right) - \frac{1}{2u} r_0^2 \right] \right. \\ \left. + \int_0^1 du \exp[-\frac{w}{u} a^2 \overrightarrow{MA}^2 - \frac{1-w}{u} a^2 \overrightarrow{MB}^2] f_0^{(2)} f_2^{(2)} \left[ \frac{1}{3a^2} {}_1F_1 \left( 1; \frac{5}{2}; \frac{a^2}{u} x_0 \right) - \frac{1}{2u} r_0^2 \right] \right\} + O(R_0^4) \quad (3.4.22)$$

where  $f_0^{(1)}$  is given by Eq. (3.1.45),  $f_2^{(1)}$  is given by (3.1.47),  $x_0$  is given by (3.1.59),  $f_0^{(2)}$  is given by (3.1.52), and  $f_2^{(2)}$  is given by (3.1.54). Replacing  $(\overrightarrow{MA} \times \overrightarrow{MB})_z$  by  $(\overrightarrow{MA} \times \overrightarrow{MB})_y$  in Eq. (3.4.22), we have the final formula of  $I_y^\sigma$ . Doing that by  $(\overrightarrow{MA} \times \overrightarrow{MB})_x$  in (3.4.22), we have  $I_x^\sigma$ . Using the 512-point GLQ, we have  $ie^2 \hbar^2 I_z^\sigma = -0.8745759(-1)i$ ,  $ie^2 \hbar^2 I_x^\sigma = -0.6184186(-1)i$ , and  $ie^2 \hbar^2 I_y^\sigma = 0$  for the case of three hydrogen atoms located at  $\vec{M} = (0, 0, 0)$ ,  $\vec{A} = (-\frac{\sqrt{8}}{3}, -\sqrt{\frac{8}{3}}, \frac{2}{3})$ , and  $\vec{B} = (-\frac{\sqrt{8}}{3}, \sqrt{\frac{8}{3}}, \frac{2}{3})$ . Note that  $(\overrightarrow{MA} \times \overrightarrow{MB})_y = MA_z MB_x - MA_x MB_z = 0$ .

## 4. Conclusion

New Gaussian-transform formulas have been derived for special derivatives of the Dirac wave function. Using these, among all necessary molecular integral-formulas for solving the molecular matrix Dirac equation (MMDE), most formulas have been derived together with those in previous articles [19-21]. All integral-formulas have been derived for the first time. We should add the Dirac wave function to our basis set for solving the MMDE.

Necessary integral-formulas to solve the MMDE are still remaining. Such are those for two-electron operators as given by

$$< \chi_\mu \chi_\kappa \left| \vec{\sigma} \cdot (\vec{p} + \vec{A}) \frac{\varepsilon^2}{r_{12}} \vec{\sigma} \cdot (\vec{p} + \vec{A}) \right| \chi_\nu \chi_\lambda > \quad (4.1)$$

Such project is in progress.

## 5. References

1. T. Yoshizawa, “On the development of the exact two-component relativistic method for calculating indirect NMR spin-spin coupling constants”, Chem. Phys. 518, 112-122 (2019).
2. S. Komorovsky, M. Repisky, O.L. Malkina, and V.G. Malkin, “Fully relativistic calculations of NMR shielding tensors using restricted magnetic balanced basis and gauge including atomic orbitals”, J. Chem. Phys. 132, 154101 (2010).
3. W-M. Sun, X-S. Chen, X-F. Liu, and F. Wang, “Gauge-invariant hydrogen-atom Hamiltonian”, Phys. Rev. A82, 012107 (2010)
4. D. Andrae, “Nuclear charge density distribution in quantum chemistry”, in P. Schwerdtfeger, editor, Relativistic Electronic Structure Theory, Part 1, Elsevier, Amsterdam (2002); pp. 203-258.
5. A. Mohanty and E. Clementi, “Dirac-Fock self-consistent field method for closed-shell molecules with kinetic balance and finite nuclear size”, Int. J. Quantum Chem. 39, 487-517 (1991).
6. L. Visscher, O. Visser, P.J.C. Aerts, H. Merrenga, and W.C. Nieuwpoort, “Relativistic quantum chemistry: the MOLFDIR program package”, Comput. Phys. Commun. 81, 120-144 (1994)
7. T. Saue, K. Faegri, T. Helgaker, and O. Gropen, “Principles of direct 4-component relativistic SCF: application to cesium auride”, Mol. Phys. 91, 937-950 (1997).

---

8. K.G. Dyall, “A systematic sequence of relativistic approximations”, *J. Comput. Chem.* 23, 786-793 (2002).
9. J. Seino and M. Hada, “Examination of accuracy of electron-electron Coulomb interactions in two-component relativistic method”, *Chem. Phys. Letters*, 461, 327-331 (2008).
10. D. Peng, N. Middendorf, F. Weigend, and M. Reiher, “An efficient implementation of two-component relativistic exact-decoupling methods for large molecules”, *J. Chem. Phys.* 138, 184105 (2013).
11. W. Liu, “Essentials of relativistic quantum chemistry”, *J. Chem. Phys.* 152, 180901 (2020).
12. S. Knecht, M. Repisky, H.J.A. Jensen, and T. Saue, “Exact two-component Hamiltonians for relativistic quantum chemistry: Two-electron picture-change corrections made simple”, *J. Chem. Phys.* 157, 114106 (2022).
13. A. Sunaga, M. Salmon, and T. Saue, “4-component relativistic Hamiltonian with effective QED potentials for molecular calculations”, *J. Chem. Phys.* 157, 164101 (2022).
14. H. Fukui, T. Baba, Y. Shiraishi, S. Imanishi, K. Kubo, K. Mori, and M. Shimoji, “Calculation of nuclear magnetic shieldings: infinite-order Foldy-Wouthuysen transformation”, *Mol. Phys.* 102, 641-648 (2004).
15. J.I. Melo, M.C. Ruiz de Azua, J.E. Peralta, and G.E. Scuseria, “Relativistic calculation of indirect NMR spin-spin couplings using the Douglas-Kroll-Hess approximation”, *J. Chem. Phys.* 123, 204112 (2005).
16. Y. Xiao, W. Liu, L. Cheng, and D. Peng, “Four-component relativistic theory for nuclear magnetic shielding constants: critical assessments of different approaches”, *J. Chem. Phys.* 126, 214101 (2007).
17. Q. Sun, Y. Xiao, and W. Liu, “Exact two-component relativistic theory for NMR parameters: general formulation and pilot application”, *J. Chem. Phys.* 137, 174105 (2012).
18. L. Cheng, J. Gauss, and J.F. Stanton, “Treatment of scalar-relativistic effects on nuclear magnetic shieldings using a spin-free exact two-component approach”, *J. Chem. Phys.* 139, 054105 (2013).
19. K. Ishida, “Gaussian-transform for the Dirac wave function and its application to the multicenter molecular integral over Dirac wave functions for solving the molecular matrix Dirac equation”, *IgMin Res.* November 04, 2: 897-914 (2024).
20. K. Ishida, “Multicenter molecular integrals over Dirac wave functions for several fundamental properties”, *IgMin Res.* February 17, 3: 076-090 (2025).
21. K. Ishida, “Multicenter molecular integrals over Dirac wave functions for relativistic kinetic energy terms”, *Open Access Journal of Chemistry*, 7: 61-80 (2025).
22. K. Ishida, “A reason why to use the Gaussian-type-orbital is not suitable for the relativistic calculation of the nuclear-magnetic-resonance spectra with using the restricted magnetic balance”, *Comput. Theor. Chem.* 1241, 114804 (2024)
23. I. S. Gradshteyn and I. M. Ryzhik, “Table of Integrals, Series, and Products”, translated from Russian by Scripta Technica Inc., Elsevier, Amsterdam, Seventh Edition, 2007.
24. K. Ishida, “Calculus of several harmonic functions”, *J. Comput. Chem. Jpn. Int. Ed.* 8, 2021-2029 (2022).
25. P. A. M. Dirac, “Principles of Quantum Mechanics”, Oxford University Press, United Kingdom, Fourth Edition, 1958.
26. R. A. Sack, “Generalization of Laplace’s expansion to arbitrary powers and functions of the distance between two points”, *J. Math. Phys.* 5, 245-251 (1964).
27. M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover Publications Inc. New York, 1970.
28. K. Ishida, “Rigorous and rapid calculation of the electron repulsion integral over the uncontracted solid harmonic Gaussian-type orbitals”, *J. Chem. Phys.* 111, 4913-4922 (1999).
29. Japanese edition translated by Y. Muroya with supervised by Y. Ohtsuki from the Russian edition, “Shin Suugaku Koushiki-shu I” (which is meaning “New Collection of Mathematical Formulas I” in English), (Maruzen Inc., Tokyo, 1991)
30. H. Moriguchi, K. Udagawa, and S. Hitotsumatsu, “Suugaku Koushiki I” which is written in Japanese and its meaning is “Mathematical Formulas I” in English, (Iwanami Inc., Tokyo, 1956).

31. E. W. Barnes, "The asymptotic expansion of integral functions defined by generalized hypergeometric series", Proc. London Math. Soc. 5, 59-116 (1907).

### Appendix A. Calculation of summations appearing in Eq. (3.2.22)

We calculate summations in Eq. (3.2.22) as follows: We first calculate the summation given by

$$\begin{aligned} S_{A0} &= \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} \frac{\Gamma(n_{123} + 5/2)}{\Gamma(n_{123} + 7/2)} \\ &= \sum_{n_1 n_2=0} \frac{(-b^2)^{n_{12}} (3/2)_{n_1} (3/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (5/2)_{n_2}} \frac{\Gamma(n_{12} + 5/2)}{\Gamma(n_{12} + 7/2)} {}_2F_2 \left( n_{12} + \frac{5}{2}, \frac{1}{2}; n_{12} + \frac{7}{2}, \frac{3}{2}; -b^2 \right) \end{aligned} \quad (a.1)$$

where  $n_{12} = n_1 + n_2$ . It is easy to derive the recurrence relation:

$${}_2F_2(a_1, a_2; a_1 + 1, a_2 + 1; x) = \frac{a_1}{a_1 - a_2} {}_1F_1(a_2; a_2 + 1; x) - \frac{a_2}{a_1 - a_2} {}_1F_1(a_1; a_1 + 1; x) \quad (a.2)$$

Using Eq. (a.2), we have

$${}_2F_2 \left( n_{12} + \frac{5}{2}, \frac{1}{2}; n_{12} + \frac{7}{2}, \frac{3}{2}; -b^2 \right) = \frac{n_{12} + 5/2}{n_{12} + 2} {}_1F_1 \left( \frac{1}{2}; \frac{3}{2}; -b^2 \right) - \frac{1/2}{n_{12} + 2} {}_1F_1 \left( n_{12} + \frac{5}{2}; n_{12} + \frac{7}{2}; -b^2 \right) \quad (a.3)$$

We know the asymptotic expansion of CHF from the formula number 13.5.1 in the Abramowitz and Stegun [27] as given by

$${}_1F_1 \left( \frac{1}{2}; \frac{3}{2}; -b^2 \right) = \frac{\sqrt{\pi}}{2b} \quad (a.4)$$

We know the integral representation of CHF from the formula number 13.2.1 in the Abramowitz and Stegun [27] as given by

$${}_1F_1 \left( n_{12} + \frac{5}{2}; n_{12} + \frac{7}{2}; -b^2 \right) = \frac{\Gamma(n_{12} + 7/2)}{\Gamma(n_{12} + 5/2)} \int_0^1 dt t^{n_{12} + 3/2} \exp(-b^2 t) \quad (a.5)$$

Substituting Eq. (a.3) and (a.4) into (a.2) and doing the resulting equation into (a.1), we have

$$\begin{aligned} S_{A0} &= \frac{\sqrt{\pi}}{2b} \sum_{n_1 n_2=0} \frac{(-b^2)^{n_{12}} (3/2)_{n_1} (3/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (5/2)_{n_2}} \frac{\Gamma(n_{12} + 2)}{\Gamma(n_{12} + 3)} \\ &\quad - \frac{1}{2} \sum_{n_1 n_2=0} \frac{(-b^2)^{n_{12}} (3/2)_{n_1} (3/2)_{n_2}}{n_1! n_2! (5/2)_{n_1} (5/2)_{n_2}} \frac{\Gamma(n_{12} + 2)}{\Gamma(n_{12} + 3)} \int_0^1 dt t^{n_{12} + 3/2} \exp(-b^2 t) \\ &= \frac{\sqrt{\pi}}{2b} \sum_{n_1=0} \frac{(-b^2)^{n_1} (3/2)_{n_1}}{n_1! (5/2)_{n_1}} \frac{\Gamma(n_1 + 2)}{\Gamma(n_1 + 3)} {}_2F_2 \left( n_1 + 2, \frac{3}{2}; n_1 + 3, \frac{5}{2}; -b^2 \right) - \frac{1}{2} \int_0^1 dt t^{3/2} \exp(-b^2 t) \\ &\quad \sum_{n_1=0} \frac{(-b^2 t)^{n_1} (3/2)_{n_1}}{n_1! (5/2)_{n_1}} \frac{\Gamma(n_1 + 2)}{\Gamma(n_1 + 3)} {}_2F_2 \left( n_1 + 2, \frac{3}{2}; n_1 + 3, \frac{5}{2}; -b^2 t \right) \end{aligned} \quad (a.6)$$

Using Eq. (a.2), we have

$$\begin{aligned} {}_2F_2 \left( n_1 + 2, \frac{3}{2}; n_1 + 3, \frac{5}{2}; -b^2 \right) &= \frac{n_1 + 2}{n_1 + 1/2} {}_1F_1 \left( \frac{3}{2}; \frac{5}{2}; -b^2 \right) - \frac{3/2}{n_1 + 1/2} {}_1F_1(n_1 + 2; n_1 + 3; -b^2) \\ &= \frac{\Gamma(n_1 + 1/2) \Gamma(n_1 + 3)}{\Gamma(n_1 + 3/2) \Gamma(n_1 + 2)} \frac{3\sqrt{\pi}}{4b^3} - \frac{3}{2} \frac{\Gamma(n_1 + 1/2) \Gamma(n_1 + 3)}{\Gamma(n_1 + 3/2) \Gamma(n_1 + 2)} \int_0^1 ds s^{n_1 + 1} \exp(-b^2 s) \end{aligned} \quad (a.7)$$

and

$$\begin{aligned} {}_2F_2 \left( n_1 + 2, \frac{3}{2}; n_1 + 3, \frac{5}{2}; -b^2 t \right) &= \frac{n_1 + 2}{n_1 + 1/2} {}_1F_1 \left( \frac{3}{2}; \frac{5}{2}; -b^2 t \right) - \frac{3/2}{n_1 + 1/2} {}_1F_1(n_1 + 2; n_1 + 3; -b^2 t) \\ &= \frac{\Gamma(n_1 + 1/2) \Gamma(n_1 + 3)}{\Gamma(n_1 + 3/2) \Gamma(n_1 + 2)} \frac{\Gamma(5/2)}{\Gamma(3/2)} \int_0^1 ds \sqrt{s} \exp(-b^2 ts) - \frac{3}{2} \frac{\Gamma(n_1 + 1/2) \Gamma(n_1 + 3)}{\Gamma(n_1 + 3/2) \Gamma(n_1 + 2)} \int_0^1 ds s^{n_1 + 1} \exp(-b^2 s) \end{aligned} \quad (a.8)$$

Substituting Eq. (a.7) and (a.8) into (a.6), we have

$$\begin{aligned}
S_{A0} = & \frac{3\pi}{8b^4} \sum_{n_1=0} \frac{(-b^2)^{n_1} (3/2)_{n_1} \Gamma(n_1 + 1/2)}{n_1! (5/2)_{n_1} \Gamma(n_1 + 3/2)} - \frac{3\sqrt{\pi}}{4b} \sum_{n_1=0} \frac{(-b^2)^{n_1} (3/2)_{n_1} \Gamma(n_1 + 1/2)}{n_1! (5/2)_{n_1} \Gamma(n_1 + 3/2)} \int_0^1 ds s^{n_1+1} \exp(-b^2 s) \\
& - \frac{1}{2} \int_0^1 dt t^{3/2} \exp(-b^2 t) \left[ \sum_{n_1=0} \frac{(-b^2 t)^{n_1} (3/2)_{n_1} \Gamma(n_1 + 1/2)}{n_1! (5/2)_{n_1} \Gamma(n_1 + 3/2)} \int_0^1 ds \sqrt{s} \exp(-b^2 ts) \right. \\
& - \frac{3}{2} \sum_{n_1=0} \frac{(-b^2 t)^{n_1} (3/2)_{n_1} \Gamma(n_1 + 1/2)}{n_1! (5/2)_{n_1} \Gamma(n_1 + 3/2)} \int_0^1 ds s^{n_1+1} \exp(-b^2 s) \left. \right] = \frac{3\pi}{8b^4} \frac{\Gamma(1/2)}{\Gamma(3/2)} {}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2\right) \\
& - \frac{3\sqrt{\pi} \Gamma(1/2)}{4b \Gamma(3/2)} \int_0^1 ds s \exp(-b^2 s) {}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2 s\right) - \frac{3}{4} \int_0^1 dt t^{3/2} \exp(-b^2 t) \\
& \left[ \int_0^1 ds \sqrt{s} \exp(-b^2 ts) \frac{\Gamma(1/2)}{\Gamma(3/2)} {}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2 t\right) \int_0^1 ds s \exp(-b^2 s) \frac{\Gamma(1/2)}{\Gamma(3/2)} {}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2 ts\right) \right] \quad (a.9)
\end{aligned}$$

We use the asymptotic expansion of the CHF from the formula number 13.5.1 in the Abramowitz and Stegun [27] as given by

$${}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2\right) = \frac{\Gamma(5/2)}{b} \left(1 - \frac{1}{2b^2}\right) \quad (a.10)$$

We use the integral representation of the CHF from the formula number 13.2.1 in the Abramowitz and Stegun [27] as given by

$${}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2 s\right) = \frac{3}{2} \int_0^1 du u^{-1/2} (1-u) \exp(-b^2 s u) \quad (a.11)$$

$${}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2 t\right) = \frac{3}{2} \int_0^1 du u^{-1/2} (1-u) \exp(-b^2 t u) \quad (a.12)$$

and

$${}_1F_1\left(\frac{1}{2}; \frac{5}{2}; -b^2 ts\right) = \frac{3}{2} \int_0^1 du u^{-1/2} (1-u) \exp(-b^2 t s u) \quad (a.13)$$

Substituting Eq. (a.10), (a.11), (a.12), and (a.13) into (a.9), we have

$$\begin{aligned}
S_{A0} = & \frac{9\pi^{3/2}}{16b^5} \left(1 - \frac{1}{2b^2}\right) - \frac{9\sqrt{\pi}}{8b} \int_0^1 du u^{-1/2} (1-u) \int_0^1 ds s \exp[-b^2 s(1+u)] \\
& - \frac{9}{8} \int_0^1 ds \sqrt{s} \int_0^1 du u^{-1/2} (1-u) \int_0^1 dt t^{3/2} \exp[-b^2 t(1+s+u)] \\
& + \frac{9}{8} \int_0^1 ds s \int_0^1 du u^{-1/2} (1-u) \int_0^1 dt t^{3/2} \exp[-b^2 t(1+s+su)] \quad (a.14)
\end{aligned}$$

We know the integral representation is nothing but the CHF as given by

$$\int_0^1 ds s \exp[-b^2 s(1+u)] = \frac{\Gamma(2)}{\Gamma(3)} {}_1F_1(2; 3; -b^2(1+u)) = \frac{\Gamma(2)}{\Gamma(3)} \frac{\Gamma(3)}{[b^2(1+u)]^2} \quad (a.15)$$

$$\int_0^1 dt t^{3/2} \exp[-b^2 t(1+s+u)] = \frac{\Gamma(5/2)}{\Gamma(7/2)} {}_1F_1\left(\frac{5}{2}; \frac{7}{2}; -b^2(1+s+u)\right) = \frac{\Gamma(5/2)}{\Gamma(7/2)} \frac{\Gamma(7/2)}{[b^2(1+s+u)]^{5/2}} \quad (a.16)$$

and

$$\int_0^1 dt t^{3/2} \exp[-b^2 t(1+s+su)] = \frac{\Gamma(5/2)}{\Gamma(7/2)} {}_1F_1\left(\frac{5}{2}; \frac{7}{2}; -b^2(1+s+su)\right) = \frac{\Gamma(5/2)}{\Gamma(7/2)} \frac{\Gamma(7/2)}{[b^2(1+s+su)]^{5/2}} \quad (a.17)$$

We use the asymptotic expansion of CHF in Eq. (a.15), (a.16) and (a.17). Substituting Eq. (a.15), (a.16), and (a.17) into (a.14), we have

$$\begin{aligned}
S_{A0} = & \frac{9\pi^{3/2}}{16b^5} - \frac{9\pi^{3/2}}{32b^7} - \frac{9\sqrt{\pi}}{8b^5} \int_0^1 du \frac{u^{-1/2}}{(1+u)^2} + \frac{9\sqrt{\pi}}{8b^5} \int_0^1 du \frac{\sqrt{u}}{(1+u)^2} \\
& - \frac{27\sqrt{\pi}}{32b^5} \int_0^1 ds \int_0^1 du \frac{\sqrt{s} u^{-1/2}}{(1+s+u)^{5/2}} + \frac{27\sqrt{\pi}}{32b^5} \int_0^1 ds \int_0^1 du \frac{\sqrt{s} \sqrt{u}}{(1+s+u)^{5/2}} \\
& + \frac{27\sqrt{\pi}}{32b^5} \int_0^1 ds \int_0^1 du \frac{s u^{-1/2}}{(1+s+su)^{5/2}} - \frac{27\sqrt{\pi}}{32b^5} \int_0^1 ds \int_0^1 du \frac{s \sqrt{u}}{(1+s+su)^{5/2}} \quad (a.18)
\end{aligned}$$

Each value of integrals appearing in Eq. (a.18) is a constant. Each constant value can be evaluated as follows: We use the formula number 2.213.4 and 2.211 of Gradshteyn and Ryzhik [23] given by

$$\int dx \frac{1}{(a+bx)^2 \sqrt{x}} = \frac{\sqrt{x}}{a(a+bx)} + \frac{1}{2a} \int dx \frac{1}{(a+bx)\sqrt{x}} \quad (a.19)$$

and

$$\int dx \frac{1}{(a+bx)\sqrt{x}} = \frac{2}{\sqrt{ab}} \operatorname{arctg} \sqrt{\frac{bx}{a}} \quad (ab > 0) \quad (a.20)$$

where  $\operatorname{arctg} x = \operatorname{arctangent} x$ . Using Eq. (a.19) and (a.20), we have

$$\int_0^1 du \frac{u^{-1/2}}{(1+u)^2} = \frac{1}{2} + \frac{\pi}{4} \quad (a.21)$$

and

$$\int_0^1 du \frac{u^{1/2}}{(1+u)^2} = -\frac{1}{2} + \frac{\pi}{4} \quad (a.22)$$

We use the formula number 2.3.8.1 of the Japanese formula book [29] as given by

$$\int_0^a dx x^{\alpha-1} (a-x)^{\beta-1} (x+z)^{-\rho} \exp(-px) = B(\alpha, \beta) z^{-\rho} a^{\alpha+\beta-1} \Phi_1 \left( \alpha, \rho, \alpha + \beta; -\frac{1}{z}, -p \right) \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, z > 0 \quad (a.23)$$

where  $B(\alpha, \beta) = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$  is the beta function and  $\Phi_1(a_1, a_2, c_1; x_1, x_2)$  is given by

$$\Phi_1(a_1, a_2, c_1; x_1, x_2) = \sum \frac{x_1^{k_1} x_2^{k_2}}{k_1! k_2!} \frac{(a_1)_{k_1+k_2} (a_2)_{k_1}}{(c_1)_{k_1+k_2}} \quad (a.24)$$

Taking the limit as  $p \rightarrow 0$  in Eq. (a.23), we have

$$\int_0^a dx \frac{x^{\alpha-1} (a-x)^{\beta-1}}{(x+z)^\rho} = B(\alpha, \beta) z^{-\rho} a^{\alpha+\beta-1} {}_2F_1 \left( \rho, \alpha; \alpha + \beta; -\frac{1}{z} \right) \quad (a.25)$$

We know the Kummer transformation as given by

$${}_2F_1(a_1, a_2; c_1; x) = (1-x)^{-a_1} {}_2F_1(a_1, c_1 - a_2; c_1; \frac{-x}{1-x}) \quad (a.26)$$

Applying Eq. (a.26) to (a.25), we have

$$\int_0^a dx \frac{x^{\alpha-1} (a-x)^{\beta-1}}{(x+z)^\rho} = \frac{B(\alpha, \beta)}{(1+z)^\rho} a^{\alpha+\beta-1} {}_2F_1 \left( \beta, \rho; \alpha + \beta; \frac{1}{1+z} \right) \quad (a.27)$$

Taking  $a = 1$  and  $z = \frac{a_2}{a_1}$ , we have

$$\int_0^1 dx \frac{x^{\alpha-1} (1-x)^{\beta-1}}{(a_1 x + a_2)^\rho} = \frac{B(\alpha, \beta)}{(a_1 + a_2)^\rho} {}_2F_1 \left( \beta, \rho; \alpha + \beta; \frac{a_1}{a_1 + a_2} \right) \quad [\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, a_1 a_2 > 0] \quad (a.28)$$

Applying Eq. (a.28) with  $\alpha = 1/2$ ,  $\beta = 1$ , and  $\rho = 5/2$  to the integral

$\int_0^1 ds \int_0^1 du \frac{\sqrt{s} u^{-1/2}}{(1+s+u)^{5/2}}$ , taking  $a_1 = 1$  and  $a_2 = 1+s$ , and using the Kummer transformation, we have

$$\int_0^1 ds \int_0^1 du \frac{\sqrt{s} u^{-1/2}}{(1+s+u)^{5/2}} = 2 \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{3/2}(1+s)} + \frac{4}{3} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{3/2}(1+s)^2} = C_{A0a} \quad (a.29)$$

The right-hand side of Eq. (a.29) can be evaluated by the 4096-point GLQ and we have  $C_{A0a} = 0.3132315213$  (which is in 10 significant-figure precision). A similar derivation to the above, we have

$$\int_0^1 ds \int_0^1 du \frac{s u^{-1/2}}{(1+s+s u)^{5/2}} = 2 \int_0^1 ds \frac{s}{(1+2s)^{3/2}(1+s)} + \frac{4}{3} \int_0^1 ds \frac{s^2}{(1+2s)^{3/2}(1+s)^2} = C_{A0b} \quad (a.30)$$

The right-hand side of Eq. (a.30) can be evaluated by the 64-point GLQ and we have  $C_{A0b} = 0.2459321581$ . Also, a similar derivation to the above, we have

$$\int_0^1 ds \int_0^1 du \frac{\sqrt{s} \sqrt{u}}{(1+s+u)^{5/2}} = \frac{2}{3} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{3/2}(1+s)} = C_{A0c} \quad (a.31)$$

Using the 4096-point GLQ, we have  $C_{A0c} = 0.7166865812(-1)$ . Again, a similar derivation to the above, we have

$$\int_0^1 ds \int_0^1 du \frac{s\sqrt{u}}{(1+s+su)^{5/2}} = \frac{2}{3} \int_0^1 ds \frac{s}{(1+2s)^{3/2}(1+s)} = C_{A0d} \quad (a.32)$$

Using the 64-point GLQ, we have  $C_{A0d} = 0.6729936319(-1)$ . Thus, we have the final formula given by

$$S_{A0} = \frac{9\pi^{3/2}}{16b^5} - \frac{9\pi^{3/2}}{32b^7} - \frac{9\sqrt{\pi}}{8b^5} - \frac{27\sqrt{\pi}}{32b^5} (C_{A0a} - C_{A0b} - C_{A0c} + C_{A0d}) \quad (a.33)$$

Next, we calculate the second summation given by

$$S_{A1} = \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3} \Gamma(n_{123}+7/2)}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3} \Gamma(n_{123}+9/2)} \quad (a.34)$$

With a similar derivation to that for  $S_{A0}$ , we have the final formula given by

$$S_{A1} = \frac{9\pi}{16b^4} C_{A1a} - \frac{9\sqrt{\pi}}{4b^7} C_{A1b} - \frac{135\sqrt{\pi}}{64b^7} (C_{A1c} - C_{A1d}) \quad (a.35)$$

where

$$C_{A1a} = \int_0^1 du \int_0^1 dv \sqrt{uv} \exp(-b^2 uv) = 0.9961218237(-2) \quad (a.36)$$

The term  $C_{A1a}$  is appearing in Eq. (3.1.26). For the term  $C_{A1b}$ , using 2048-point GLQ, we have

$$C_{A1b} = \int_0^1 du \int_0^1 dv \frac{\sqrt{uv}}{(1+uv)^3} = 0.2079827971 \quad (a.37)$$

For the term  $C_{A1c}$ , using Eq. (a.28) with  $a_1 = u$ ,  $a_2 = 1+s$ ,  $\alpha = 3/2$ ,  $\beta = 1$ , and  $\rho = 7/2$ , doing the Kummer transformation, doing (a.28) again with  $a_1 = 1$ ,  $a_2 = 1+s$ ,  $\alpha = 3/2$ ,  $\beta = 1$ , and  $\rho = 7/2$  for the resulting equation, doing the Kummer transformation again, and doing the 4096-point GLQ for it, we have its constant value as given by

$$\begin{aligned} C_{A1c} &= \int_0^1 du \int_0^1 dv \int_0^1 ds \frac{\sqrt{uv}}{(1+s+uv)^{7/2}} = \frac{4}{9} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{3/2}(1+s)^2} + \frac{8}{75} \int_0^1 ds \frac{\sqrt{s} {}_2F_1[1.5/2; 7/2; 1/(2+s)]}{(2+s)^{5/2}(1+s)^2} \\ &= 0.3742687401(-1) \end{aligned} \quad (a.38)$$

For the term  $C_{A1d}$ , using Eq. (a.28) with  $a_1 = su$ ,  $a_2 = 1+s$ ,  $\alpha = 3/2$ ,  $\beta = 1$ , and  $\rho = 7/2$ , doing the Kummer transformation, doing (a.28) again with  $a_1 = s$ ,  $a_2 = 1+s$ ,  $\alpha = 3/2$ ,  $\beta = 1$ , and  $\rho = 7/2$  for the resulting equation, doing the Kummer transformation again, and doing the 64-point GLQ for it, we have its constant value as given by

$$\begin{aligned} C_{A1d} &= \int_0^1 du \int_0^1 dv \int_0^1 ds \frac{s^2 \sqrt{uv}}{(1+s+suv)^{7/2}} = \frac{4}{9} \int_0^1 ds \frac{s^2}{(1+2s)^{3/2}(1+s)^2} + \frac{8}{75} \int_0^1 ds \frac{s^2 {}_2F_1[1.5/2; 7/2; s/(1+2s)]}{(1+2s)^{5/2}(1+s)^2} \\ &= 0.1586020540(-1) \end{aligned} \quad (a.39)$$

Further, we calculate the third summation given by

$$S_{A2} = \sum_{n_1 n_2 n_3=0} \frac{(-b^2)^{n_{123}} (3/2)_{n_1} (3/2)_{n_2} (1/2)_{n_3}}{n_1! n_2! n_3! (5/2)_{n_1} (5/2)_{n_2} (3/2)_{n_3}} \frac{\Gamma(n_{123}+9/2)}{\Gamma(n_{123}+11/2)} \quad (a.40)$$

With a similar derivation to that for  $S_{A0}$ , we have the final formula given by

$$S_{A2} = \frac{9\pi^{2/2}}{32b^7} - \frac{27\pi^{2/2}}{64b^9} - \frac{27\sqrt{\pi}}{4b^9} (C_{A2a} - C_{A2b}) - \frac{945\sqrt{\pi}}{128b^9} (C_{A2c} - C_{A2d} - C_{A2e} + C_{A2f}) \quad (a.41)$$

where

$$C_{A2a} = \int_0^1 du \frac{\sqrt{u}}{(1+u)^4} = 2 \int_0^1 dx \frac{x^2}{(1+x^2)^4} \quad (a.42)$$

and

$$C_{A2b} = \int_0^1 du \frac{u^{3/2}}{(1+u)^4} = 2 \int_0^1 dx \frac{x^4}{(1+x^2)^4} \quad (a.43)$$

We take formulas from a Japanese mathematical formula book [30] given by

$$\int dx \frac{x^2}{(x^2 + c)^4} = -\frac{x}{6(x^2 + c)^3} + \frac{x}{24c(x^2 + c)^3} + \frac{x}{16c^2(x^2 + c)} + \frac{1}{16c^{5/2}} \arctan\left(\frac{x}{\sqrt{c}}\right) \quad (c > 0) \quad (a.44)$$

and

$$\int dx \frac{x^4}{(x^2 + c)^4} = \frac{cx}{6(x^2 + c)^3} - \frac{7x}{24(x^2 + c)^3} + \frac{x}{16c(x^2 + c)} + \frac{1}{16c^{5/2}} \arctan\left(\frac{x}{\sqrt{c}}\right) \quad (c > 0) \quad (a.45)$$

where  $\arctan(x) = \arctangent(x)$ . Using Eq. (a.44) with  $c = 1$  and integrating it for the interval  $[0, 1]$ , we have the constant value of  $C_{A2a}$  as given by  $C_{A2a} = \frac{1}{24} + \frac{\pi}{32}$ . Using Eq. (a.45) with  $c = 1$ , we have also the constant value as given by  $C_{A2b} = -\frac{1}{24} + \frac{\pi}{32}$ . For the term  $C_{A2c}$ , using Eq. (a.28) with  $\alpha = \frac{3}{2}$ ,  $\beta = 1$ ,  $\rho = \frac{9}{2}$ ,  $a_1 = 1$ , and  $a_2 = 1 + s$ , doing the Kummer transformation, Eq. (a.26), for the resulting equation, and doing the 4096-point GLQ for it, we have the constant value as given by

$$C_{A2c} = \int_0^1 ds \int_0^1 du \frac{\sqrt{su}}{(1+s+u)^{9/2}} = \frac{2}{3} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{7/2}(1+s)} + \frac{8}{15} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{7/2}(1+s)^2} + \frac{16}{105} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{7/2}(1+s)^3} = 0.1994662002(-1) \quad (a.46)$$

For the term  $C_{A2d}$ , using Eq. (a.28) with  $\alpha = \frac{3}{2}$ ,  $\beta = 1$ ,  $\rho = \frac{9}{2}$ ,  $a_1 = 1$ , and  $a_2 = 1 + u$ , doing the Kummer

transformation for the resulting equation, and doing the 128-point GLQ for it, we have the constant value as given by

$$C_{A2d} = \int_0^1 ds \int_0^1 du \frac{\sqrt{su}^{3/2}}{(1+s+u)^{9/2}} = \frac{2}{3} \int_0^1 du \frac{u^{3/2}}{(2+u)^{7/2}(1+u)} + \frac{8}{15} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{7/2}(1+s)^2} + \frac{16}{105} \int_0^1 ds \frac{\sqrt{s}}{(2+s)^{7/2}(1+s)^3} = 0.8519836140(-2) \quad (a.47)$$

For the term  $C_{A2e}$ , similarly to  $C_{A2c}$  and using the 64-point GLQ, we have its constant value as given by

$$C_{A2e} = \int_0^1 ds \int_0^1 du \frac{s^3 \sqrt{u}}{(1+s+su)^{9/2}} = \frac{2}{3} \int_0^1 ds \frac{s^3}{(1+2s)^{7/2}(1+s)} + \frac{8}{15} \int_0^1 ds \frac{s^4}{(1+2s)^{7/2}(1+s)^2} + \frac{16}{105} \int_0^1 ds \frac{s^5}{(1+2s)^{7/2}(1+s)^3} = 0.5697996217(-2) \quad (a.48)$$

For the term  $C_{A2f}$ , similarly to  $C_{A2c}$  and using the 64-point GLQ, we have its constant value as given by

$$C_{A2f} = \int_0^1 ds \int_0^1 du \frac{s^3 u^{3/2}}{(1+s+su)^{9/2}} = \frac{2}{5} \int_0^1 ds \frac{s^3}{(1+2s)^{7/2}(1+s)} + \frac{4}{35} \int_0^1 ds \frac{s^4}{(1+2s)^{7/2}(1+s)^2} = 0.2821839924(-2) \quad (a.49)$$

## Appendix B. Calculation of ${}_2F_2$ Functions

We calculate  ${}_2F_2\left(1, 1; 2 \frac{9}{2}; x\right)$  and  ${}_2F_2\left(1, 1; 3 \frac{7}{2}; x\right)$  here. Barnes[31] showed the asymptotic expansion formula for the function  ${}_pF_p(\alpha_1, \dots, \alpha_p; \rho_1, \dots, \rho_p; x)$  as given by

$$\prod_{r=1}^p \left\{ \frac{\Gamma(\alpha_r)}{\Gamma(\rho_r)} \right\} {}_pF_p\{x\} = \exp(x) x^{\sum \alpha_r - \sum \rho_r} \left\{ 1 + \sum_{r=1}^p \frac{M_r}{x^r} + \frac{J_R}{x^R} \right\} \quad (b.1)$$

where the  $J_R/x^R$  is the error term. However, he did not show the explicit formula for  $M_r$ , then, after a numerical experiment, we have the asymptotic expansion of the  ${}_2F_2\left(1, 1; 2 \frac{9}{2}; x\right)$  as given by

$${}_2F_2\left(1, 1; 2 \frac{9}{2}; x\right) = \frac{\Gamma(2)\Gamma(9/2)}{\Gamma(1)\Gamma(1)} x^{-9/2} \exp(x) \sum_{n=0}^{40} \frac{(1)_n (7/2)_n}{n! x^n} \quad (\text{for } x \geq 37) \quad (b.2)$$

For  $x \leq 37$ , we calculate it by the power series given by

$${}_2F_2\left(1, 1; 2 \frac{9}{2}; x\right) = \sum_{n=0}^{\infty} \frac{x^n (1)_n (1)_n}{n! (2)_n (9/2)_n} \quad (b.3)$$

After a numerical experiment, we have the asymptotic expansion of the  ${}_2F_2\left(1, 1; 3 \frac{7}{2}; x\right)$  as given by

$${}_2F_2\left(1, 1; 3 \frac{7}{2}; x\right) = \frac{\Gamma(3)\Gamma(7/2)}{\Gamma(1)\Gamma(1)} x^{-9/2} \exp(x) \sum_{n=0}^{40} \frac{(2)_n (5/2)_n}{n! x^n} \quad (\text{for } x \geq 38) \quad (b.4)$$

For  $x \leq 38$ , we calculate it by the power series given by

$${}_2F_2\left(1, 1; 3 \frac{7}{2}; x\right) = \sum_{n=0}^{\infty} \frac{x^n (1)_n (1)_n}{n! (3)_n (7/2)_n} \quad (b.5)$$