

# Clinical Applications of Low-Level Laser Therapy in Reproductive Medicine: A Literature Review

Soheila Borhani, M.D<sup>1\*</sup>, Reza Salman Yazdi, DCLS<sup>2</sup>

<sup>1</sup>NYU Langone Health, New York, USA.

<sup>2</sup>Artin Clinical Laboratory, Tehran, Iran.

Soheila.Borhani@nyumc.org

*\*Corresponding Author: Soheila Borhani, NYU Langone Health, 550 First Ave., New York, NY 10016*

## Abstract

*Infertility affects approximately 15% of couples worldwide, an estimated 30% of which is related to male factor infertility. Application of low level laser therapy (LLLT) to improve fertility status is a rapidly growing discipline in medicine. Laser therapy triggers a variety of biological processes through interaction with primary cellular photoacceptors. The present review aims at evaluating the literature available in the MEDLINE/ PubMed on cellular and molecular mechanisms of photobiomodulation in the domains of reproductive and veterinary medicine. We primarily focused on the clinical application of laser treatment on seminal quality, in particular its role in promoting spermatozoa motility, as well as the role of phototherapy in modulating assisted reproduction (ART). Additionally, we investigated the strength of evidence in support of the positive impact of photobiomodulation on facilitating ART.*

**Keywords:** *low-level laser therapy (LLLT); photobiomodulation; male infertility; sperm motility; assisted reproductive technology (ART)*

## INTRODUCTION

Phototherapy with a low-level laser which is referred to as photobiomodulation is a developing discipline in the medicine. Low-level laser therapy (LLLT) is implicated in a verity of biological processes through interaction with the primary photoacceptors and messengers [1]. The efficacy of LLLT is highly dependent on the irradiated tissues and cells as well as the specific irradiation parameters. There is numerous evidence regarding the “in vitro” and “in vivo” effects of photobiomodulation on different medical conditions. In the dentistry area low-level laser is applied to treat dentine hypersensitivity, periodontitis, and improving oral mucositis [2-4]. Moreover, LLLT is used for management of many dermatologic disorders, including alopecia, and telangiectasia [5, 6]. Additionally, the efficacy of laser phototherapy is approved for the treatment of musculoskeletal and rheumatologic conditions [7, 8]. Of note, the photobiomodulation is implicated in nerve regeneration, promoting of wound healing, and reduction in TNF- $\alpha$  levels as well as a consequent

subsiding of inflammation along with acceleration of cellular proliferation [9-11]. On the other side, LLLT technique is largely considered as an emerging concept in the veterinary medicine.

Clinical applications of the LLLT on improving fertility status is a rapid growing issue. Epidemiological studies estimate about 15% of couples are affected by infertility worldwide [12]. According to Agarwal et al. [13], at least thirty million men are infertile globally with the highest rates in Africa and Eastern Europe. Asthenospermia or attenuation of the motile spermatozoa to less than 50 percent is a significant contributing factor in the male infertility. Asthenospermia is a multifactorial disorder which mandates a diverse therapeutic approach. There is considerable literature on the treatment modalities leading to improving sperm motility, including varicocelelectomy [14], as well as dietary supplementation with some certain vitamins and antioxidants [15]. Furthermore, the potency of such compounds as aromatase inhibitors [16], pentoxifylline [17], thyroxin [18], and exogenous platelet activating

factor [19] have been demonstrated in acceleration of sperm motility. Recent advances in the domain of assisted reproductive techniques provide a novel therapeutic approach to the male factor infertility, and last but not least is the application of laser therapy to promote asthenospermia.

The present review aims to consolidate the database available on the efficacy of low-level laser light in the treatment of infertility. Specifically, we evaluate the cellular and molecular mechanisms of photobiomodulation, and put the focus on the photo-stimulative effect of LLLT on spermatozoa and improving the seminal quality as well as its pivotal role in modulating the Assisted Reproductive Techniques (ART). The resources included the relevant database in the MEDLINE/ PubMed. We did not take the publication date and publication status, nor the manuscript language into consideration. The search strategy consisted of the keyword terms, consisting of low-level laser therapy (LLLT), photobiomodulation, infertility, sperm motility, assisted reproductive technology (ART).

### EFFECT OF PHOTOBIO-MODULATION ON SEMINAL PARAMETERS

Improving male factor infertility using low-level laser therapy is a discernable area in the field of reproductive medicine. Also, the phototherapy has been extensively administered in the veterinary practice and livestock breeding. The underlying pathways of photobiomodulation are not well established. However, several studies addressed the probable mechanisms regarding the interaction of the laser light and spermatozoa. Motility is one of the most prominent characteristics of sperm associated with the fertilizing capability. The mitochondrial apparatus within the midpiece of the spermatozoa provides the required energy for movement of the flagellum or tail [20]. Albuquerque-Pontes et al. [21] indicated that low-level laser irradiation would induce the activity of the cytochrome c oxidase (COX). The COX complex is part of the mitochondrial respiratory chain and plays a critical role in the electron transport cascade. Modulation of this certain cytochrome oxidase activity leads to enhanced oxidative phosphorylation or adenosine triphosphate (ATP) generation. This process would subsequently augment the sperm motility. Likewise, another study evaluating the efficacy of laser on the cryopreserved ram sperm

showed an increase in the COX Vmax values as well as the COX affinity for its substrate. These findings were in parallel with enhanced ATP levels in the irradiated samples and improved sperm motility [22]. Additionally, Passarella et al. [23] reported that some certain NADH-linked reactions occurring in the mitochondria are stimulated and triggered through laser irradiation.

On the molecular level, the LLLT is mediated in the up-regulation of genes coding for a number of mitochondrial enzymes. Specifically, the subunits which are involved in the complexes I and IV of electron transport chain and ATP synthase [24]. Ferraresi et al. [25] evaluated the mitochondrial membrane potential and demonstrated that phototherapy would increase ATP synthesis in the myotubes. On a different note, Tafur et al. [26] declared that low-intensity laser interacts with the endogenous cellular redox mechanisms. This effect is mediated through photoexcitation of cytochrome c oxidase in the mitochondrial electron transport chain. Laser light facilitates electron transferring to the oxygen molecules and production of the reactive oxygen species (ROS). These anions are categorized into three main types, namely superoxide, hydrogen peroxide, and hydroxyl radical. ROS are necessary for the process of spermatozoa maturation or capacitation [27]. Also, low levels of ROS could relatively enhance the sperm acrosome reaction [28]. According to Shahar et al. [29] photobiomodulation resulted in a significant increase in the human sperm motility and capacitation toward activation of protein kinase A and sarcoma protein kinase, as well as the production of reactive oxygen species.

Another aspect of photobiomodulation is the effect of irradiation on the intracellular calcium ion levels, which has a fundamental impact on the sperm motility. Low-level laser therapy increases the calcium influx by means of cellular pumps. In this regard,  $\text{Na}^+/\text{Ca}^{2+}$  exchanger and voltage-gated calcium channels regulate the optimal intracellular calcium concentrations [30]. In addition, Lubart et al. [31] reported that LLLT prevents the calcium uptake by mitochondria of spermatozoa while enhancing the  $\text{Ca}^{2+}$  binding to sperm plasma membrane. On the other hand, laser light at higher doses causes an overload in the intracellular  $\text{Ca}^{2+}$  levels. Such a process leads to hyperactivation of the  $\text{Ca}^{2+}$ -ATPase pump and exhausts

the ATP reservoir of the cells. These particular reactions would ultimately increase the intracellular osmotic pressure and degenerate the spermatozoa [32].

Effect of the laser therapy on the sperm parameters is directly related to the semen sample quality, irradiation methods, applied doses, wavelengths, and time intervals. As mentioned earlier on the role of laser irradiation in the cellular calcium regulation, the spermatozoa react differently in response to the various laser doses. This fact emphasizes the importance of selecting the optimal output power. A number of research studies compared the efficacy of LLLT on male infertility using different irradiation methods, including a recent study conducted by our research group on the impact of 830 nm diode laser on human sperm motility [33]. We evaluated the semen specimens of asthenospermic patients. Each Sample was divided into four equal portions and exposed to aGaAlAs laser beam. Four different doses were administered; no irradiation for the control group, 4 J/cm<sup>2</sup>, 6 J/cm<sup>2</sup>, and 10 J/cm<sup>2</sup>. Sperm motility was assessed by means of computer-aided sperm analysis (CASA) at various time intervals, which included immediately following irradiation, 30, 45 and 60 minutes after the intervention. In order to evaluate the functional capacity of the spermatozooids, the aliquots were undergone the hypo-osmotic swelling (HOS) test. Besides, the sperm DNA fragmentation was assessed through the sperm chromatin dispersion (SCD). The two latter tests were performed only on the control group as well as the ones irradiated by 10 J/cm<sup>2</sup> (the highest irradiation dose in this study). According to the results, LLLT improved the human sperm motility at certain laser density and specific post-exposure time. The semen specimens which received irradiation doses of 4 J/cm<sup>2</sup> and 6 J/cm<sup>2</sup>, revealed a significant increase in the progressive sperm motility at the times of 60 and 45 minutes, respectively. Moreover, the results of the HOS and SCD tests showed any significant difference between the control group and the samples which received 10 J/cm<sup>2</sup> fluency.

In a similar clinical trial on male infertility, Salama et al. [34] studied the effect of light-emitting diode (LED) on improving seminal quality in the patients with and without asthenospermia. 27 cases were involved in

the study. The semen samples were aliquoted into two parts, the ones which exposed to the red LED as well as the control group. The test tubes were irradiated by LED (wavelength; 636.6-nm) for 2, 5 and 10 minutes. The irradiation doses were calculated as 496 mJ/cm<sup>2</sup>, 1.241 J/cm<sup>2</sup> and 2.482 J/cm<sup>2</sup> for the 2, 5 and 10 minutes, respectively. Sperm kinetics analysis, sperm creatine kinase (CK) activity, aniline blue staining (ANBS), and HOS tests were all evaluated. The CK activity test analyzed the rate of adenosine triphosphate synthesis by the spermatozoa, and the aniline blue staining was performed for the assessment of sperm chromatin condensation. The authors indicated a significant increase in the progressive sperm motility among semen specimens irradiated by the red LED. In addition, they reported an augmented sperm creatine kinase activity in the test tubes. However, the aforementioned results were not statistically significant. Furthermore, they found that treatment with LED could not modify the HOS test and ANBS results compared to the control group.

Additionally, low-level laser therapy has a critical role in improving the longevity of spermatozoa in the area of veterinary practice. In this regard, Laffaldano et al. [35], experimented He-Ne laser irradiation on the stored turkey semen. The energy dose of 3.96 J/cm<sup>2</sup> was applied and the effect of the LLLT on sperm preservation for up to 60 hours has been evaluated. Exposure to this specific energy dose has significantly enhanced the viability and semen quality in long-term storage compared to the control group. In another study conducted by the same author [36], the efficacy of photobiomodulation on the rabbit spermatozoa surveillance during liquid storage conditions has been investigated. The semen pools were divided into four aliquots and irradiated with different energy doses (3.96, 6.12, 9 J/cm<sup>2</sup>) of He-Ne laser. The authors found that semen samples which were exposed to the energy dose of 6.12 J/cm<sup>2</sup> better maintained viability during 48 hours of in vitro liquid storage at 15°C. A number of relevant research studies on low-level laser therapy were summarized in the table.1, which specifically evaluate the impact of photobiomodulation on the seminal parameters in both human and veterinary medicine.

## Clinical Applications of Low-Level Laser Therapy in Reproductive Medicine: A Literature Review

**Table 1.** Literature on the effect of photobiomodulation in human and veterinary reproductive medicine.

Author /Ref	Irradiation Source	Studied Species	Interpretation of Results
Firestone et al. [37]	Laser; 905 nm 1.5 J/cm <sup>2</sup>	Human	Increased sperm motility No increase in DNA damage
Siquiera et al. [38]	He-Ne laser; 633 nm 5.57, 10 mW	Bull	Increased sperm motility Increased mitochondrial function
Ban Frangez et al. [39]	LED; 470-850 nm	Human	Increased sperm motility Decreased immotile sperms
Yeste et al. [40]	LED; 660 nm	Boar	Increased sperm motility Increased sperm viability
Cohen et al. [41]	He-Ne laser; 630 nm	Mouse	Increased H <sub>2</sub> O <sub>2</sub> generation Increased intracellular Ca <sup>2+</sup>
Abdel- Salam et al. [42]	Laser; 533 nm 0.076-0.38 J/cm <sup>2</sup>	Bull	Improvement in semen quality
Fernandes et al. [43]	AlGalnP laser; 660 nm 4, 6 J/cm <sup>2</sup>	Bull	Increased sperm motility Increased sperm viability Increased acrosome integrity
Laffalando et al. [22]	He-Ne laser 3.96-9 J/cm <sup>2</sup>	Ram	Increased sperm motility Increased sperm velocity
Laffalando et al. [36]	He-Ne laser; 660 nm 3.96-9 J/cm <sup>2</sup>	Rabbit	Increased sperm motility Increased sperm viability Increased acrosome integrity
Laffalando et al. [35]	He-Ne laser 0.14-10.8 J/cm <sup>2</sup>	Turkey	Increased sperm motility Increased sperm viability
Baques et al. [44]	Laser; 655 nm 4, 6, 10 J/cm <sup>2</sup>	Dog	Increased sperm velocity Increased linear coefficient
Brito et al. [45]	LED; 660 nm 6 J/cm <sup>2</sup>	Dog	No increase in sperm kinetics
Sato et al. [46]	Krypton laser; 647 nm 4, 8, 32 J/cm <sup>2</sup>	Human	Increased sperm motility
Baques et al. [47]	Laser; 655 nm 3.3418 J/cm <sup>2</sup>	Dog	Increased sperm motility
Quero et al. [48]	He-Ne laser; 632 nm 2-16 J/cm <sup>2</sup>	Bull	Increased sperm viability Increased acrosome reaction
Dreyer et al. [49]	He-Ne laser; 633 nm 150-600 J/cm <sup>2</sup>	Bull	Increased acrosome reaction Altered DNA methylation

### EFFECT OF LASER IRRADIATION ON ASSISTED REPRODUCTION

In vitro manipulating of the gamete cells to achieve the fertilized egg is the cornerstone of assisted reproductive technology (ART). The laser beam has been implicated in treating both spermatozoa and oocytes prior to the intracytoplasmic sperm injection (ICSI). Montag et al. [50], indicated that non-contact, diode laser is an effective procedure for immobilization of human spermatozoa and permeabilization of the sperm tail membrane before ICSI. In a similar way, Ebner et al. [51], showed that the spermatozoa samples which were immobilized by laser, required a considerably shorter time for identification, aspiration and injection in comparison to the mechanically immobilized group. Moreover, laser-assisted sperm micromanipulation is a novel approach in the area of artificial insemination. Obruca et al. [52], used the Er:YAG laser for the subzonal insemination (SUZI). During this study, the laser treated population achieved a significantly higher fertilization rates compared to mechanical SUZI. In addition, sperm viability assessment is regarded as a prerequisite for the intracytoplasmic sperm injection. Laser could be implicated in the selection of viable spermatozoa, especially in the individuals with an indefinite HOS test results [53]. The potential role of laser therapy in handling the oocytes during assisted reproduction, also has been noticed in the recent years. Degeneration of the oocytes in the course of ICSI is an inevitable issue, particularly in the case of fragile oolemma. In order to mitigate oocyte degeneration associated with microinjection, Abdelmassih et al. [54], has applied laser beam to produce a microhole on the zona pellucida of the oocyte. This approach provided a less traumatic penetration into the ooplasm than a microneedle and resulted in much lower cellular degeneration. Rienzi et al. [55], has achieved similar results on promoting the oocytes survival rate in the patients with inherent oocyte fragility following ICSI procedure.

Eventually, the impact of LLLT on the oocytes maturation merits consideration. However, there is not sufficient database regarding this concept. The oocytes maturation encompasses a complex of processes resulted in completion of meiosis and subsequent fertilization [56]. Soares et al. [57], investigated the effect of laser therapy on bovine oocyte and particularly the granulosa

cells metabolism. During this study the cumulus-oocytes complexes (COCs) were exposed to the laser irradiation with 633-nm wavelength and 1 J/cm<sup>2</sup> fluency. The COCs were evaluated for cell cycle status, mitochondrial functioning, as well as viability. The number of cells progressing through the cycle and mitochondrial membrane potential enhanced significantly. Besides, cyclin B and cyclin-dependent kinase (CDK4) levels were similarly increased. With regard to the oocytes, there was an escalation in the total mitogen-activated protein kinase along with a decrease in all cell cycle genes transcripts, except the CDK4. In a different experiment, He-Ne laser irradiation at 0.05 and 0.25 J/cm<sup>2</sup> was found to increase the number of unreduced (diploid) oocytes. Concomitantly, oocytes degeneration during the in vitro meiosis was enhanced as well [58]. Likewise, Moreno-Millan et al. [59], evaluated the application of He-Ne irradiation in the “in vitro” fertilization. The authors declared that laser therapy at some certain doses triggered oocyte nuclear damage and suppressing oocyte maturation. The efficacy of LLLT mandates an optimal energy level. Hence, selecting the proper power is a key component in the process of oocyte maturation, and more generally improving the infertility.

### CONCLUSION

A variety of methods have been developed for the treatment of infertility and subfertility, including low level laser therapy (LLLT). Nonetheless, further clinical surveys should be designed to approve the clinical efficacy of LLLT products in the treatment of infertility. Moreover, it is imperative to address the effectiveness of photobiomodulation on improving the pregnancy outcomes in the future studies.

### REFERENCES

- [1] Tiphlova O, Karu T. Stimulation of Escherichia coli division by low-intensity monochromatic visible light. *Photochem Photobiol.* 1988; 48: 467-471.
- [2] Moneib H, Tawfik AA, Youssef SS, Fawzy MM. Randomized split-face controlled study to evaluate 1550-nm fractionated erbium glass laser for treatment of acne vulgaris—an image analysis evaluation. *Dermatol Surg.* 2014; 40: 1191-1200.



- [3] Sculean A, Schwarz F, Becker J. Anti-infective therapy with an Er:YAG laser: influence on peri-implant healing. *Expert Rev. Med. Devices.* 2005; 3: 267–276.
- [4] Gupta A, Keshri GK, Yadav A, Gola S, Chauhan S, Salhan AK, Bala and Singh S. Superpulsed (Ga-As, 904 nm) low-level laser therapy (LLLT) attenuates inflammatory response and enhances healing of burn wounds. *J Biophotonics.* 2015; 8:489-501.
- [5] Rangwala S, Rashid, RM. Alopecia: a review of laser and light therapies. *Dermatol Online J.* 2012; 18: 3.
- [6] Fioramonti P, Fino P, Ponzo I, Ruggieri M, Onesti MG. Intense pulsed light in the treatment of telangiectasias: case report of Behçet's disease with superficial vascular involvement. *J Cosmet Laser Ther.* 2014; 16: 124–128.
- [7] Chen J, Huang Z, Ge M, Gao M. Efficacy of low-level laser therapy in the treatment of TMDs: a meta-analysis of 14 randomized controlled trials. *Journal of Oral Rehabilitation.* 2015; 42: 291–299.
- [8] Denadai AS, de Carvalho Pde T, dos Reis FA, Belchior AC, Pereira DM, Dourado DM, et al. Morphometric and histological analysis of low-power laser influence on bone morphogenetic protein in bone defects repair. *Lasers Med Sci.* 2009; 24:689–95.
- [9] Wang CZ, Chen YJ, Wang YH, Yeh ML, Huang MH, Ho ML, et al. Low-level laser irradiation improves functional recovery and nerve regeneration in sciatic nerve crush rat injury model. *PLOS One.* 2014; 9: e103348.
- [10] Calderhead RG, Kubota J, Trelles MA, Ohshiro T. One mechanism behind LED phototherapy for wound healing and skin rejuvenation: key role of the mast cell. *Laser Ther.* 2008; 17:141-148.
- [11] Goralczyk K, Szymańska J, Szot K, Fisz J, Rosc D. Low-level laser irradiation effect on endothelial cells under conditions of hyperglycemia. *Lasers Med Sci.* 2016; 31:825–831.
- [12] Sharlip ID, Jarow JP, Belker AM, Lipshultz LI, Sigman M, Thomas AJ, et al. Best practice policies for male infertility. *FertilSteril.* 2002; 77: 873–882.
- [13] Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *ReprodBiolEndocrinol.* 2015; 13: doi: 10.1186/s12958-015-0032-1.
- [14] Baker K, McGill J, Sharma R, Agarwal A, Sabanegh E. Pregnancy after varicocelectomy: impact of postoperative motility and DFI. *Urology.* 2013; 81:760-766.
- [15] Kumalic SI, Pinter B. Review of Clinical Trials on Effects of Oral Antioxidants on Basic Semen and Other Parameters in Idiopathic Oligoasthenoteratozoospermia. *Biomed Res Int.* 2014; 2014: 426951.
- [16] Raman JD, Schlegel PN. Aromatase inhibitors for male infertility. *J Urol.* 2002; 167:624–629.
- [17] Stanic P, Sonicki Z, Suchanek E. Effect of pentoxifylline on motility and membrane integrity of cryopreserved human spermatozoa. *Int J Androl.* 2002; 25:186–190.
- [18] Mendeluk GR, Rosales M. Thyroxin Is Useful to Improve Sperm Motility. *Int J FertilSteril.* 2016; 10: 208-214.
- [19] Grassi G, Cappello N, Gheorghe MF, Salton L, Di Bisceglie C, Manieri C, et al. Exogenous platelet activating factor improves the motility of human spermatozoa evaluated with C.A.S.A.: optimal concentration and incubation time. *J Endocrinol Invest.* 2010; 33:684-690.
- [20] Huang YY, Chen AC, Carroll JD, Hamblin MR. Biphasic dose response in low level light therapy. *Dose Response.* 2009; 7:358–383.
- [21] Albuquerque-Pontes GM, Vieira R de P, Tomazoni SS, Caires CO, Nemeth V, Vanin AA, et al. Effect of pre-irradiation with different doses, wavelengths, and application intervals of low-level laser therapy on cytochrome c oxidase activity in intact skeletal muscle of rats. *Lasers Med Sci.* 2015; 30:59–66.
- [22] Iaffaldano N, Paventi G, Pizzuto R, Di Iorio M, Bailey JL, Manchisi A, et al. Helium-neon laser irradiation of cryopreserved ram sperm enhances cytochrome c oxidase activity and ATP levels improving semen quality. *Theriogenology.* 2016; 86: 778-784.

- [23] Passarella S, Marra E, Doonan S, Quagliariello E. Uptake of malate dehydrogenase into mitochondria in vitro. Some characteristics of the process. *Biochem J.* 1983; 210: 207–214.
- [24] Masha RT, Houreld NN, Abrahamse H. Low-intensity laser irradiation at 660 nm stimulates transcription of genes involved in the electron transport chain. *Photomed Laser Surg.* 2013; 31: 47-53.
- [25] Ferraresi C, Kaippert B, Avci P, Huang YY, Pires de Sousa MV, et al. Low-level laser (light) therapy increases mitochondrial membrane potential and ATP synthesis in C2C12 myotubes with a peak response at 3-6 hours. *PhotochemPhotobiol.* 2015; 91: 411–416.
- [26] Tafur J, Mills PJ. Low-Intensity Light Therapy: Exploring the Role of Redox Mechanisms. *Photomed Laser Surg.* 2008; 26: 323–328.
- [27] Griveau JF, Le Lannou D. Influence of oxygen tension on reactive oxygen species production and human sperm function. *Int J Androl.* 1997; 20:195–200.
- [28] Lubart R. A New Method for Stimulating the Fertilizing Capability of Sperm Cells. *Recent Patents on Biomed Eng.* 2009; 2: 193-196.
- [29] Shahar S, Wisner A, Ickowicz D, Lubart R, Shulman A, Breitbart H. Light-mediated activation reveals a key role for protein kinase A and sarcoma protein kinase in the development of sperm hyper-activated motility. *Hum Reprod.* 2011; 26, No.9 pp. 2274–2282.
- [30] Gadella BM, Luna C. Cell biology and functional dynamics of the mammalian sperm surface. *Theriogenology.* 2014; 81:74–84.
- [31] Lubart R, Friedmann H, Sinyakov M, Cohen N, Breitbart H. Changes in calcium transport in mammalian sperm mitochondria and plasma membranes caused by 780 nm irradiation. *Lasers Surg Med.* 1997; 21:493–499.
- [32] Malik Z, Lugaci H. Destruction of erythroleukaemic cells by photoactivation of endogenous porphyrins. *Br J Cancer.* 1987; 56:589–595.
- [33] Salman Yazdi R, Bakhshi S, Jannat Alipoor F, Akhoond MR, Borhani S, Farrahi F, et al. Effect of 830-nm diode laser irradiation on human sperm motility. *Lasers Med Sci.* 2014; 29:97–104.
- [34] Salama N, El-Sawy M. Light-emitting diode exposure enhances sperm motility in men with and without asthenospermia: Preliminary results. *ArchivioItaliano di Urologia e Andrologia.* 2015; 87: 14-19.
- [35] Iaffaldano N, Meluzzi A, Manchisi A, Passarella S. Improvement of stored turkey semen quality as a result of He-Ne laser irradiation. *AnimReprod Sci.* 2005; 85:317-325.
- [36] Iaffaldano N, Rosato MR, Paventib G, Pizzuto R, Gambacorta M, Manchisi A. The irradiation of rabbit sperm cells with He-Ne laser prevents their in vitro liquid storage dependent damage. *AnimReprod Sci.* 2019; 119: 123–129.
- [37] Firestone RS, Esfandiari N, Moskovtsev SI, Burstein E, Videna GT, Librach C. The Effects of Low-Level Laser Light Exposure on Sperm Motion Characteristics and DNA Damage. *J Androl.* 2012; 33: 469-473.
- [38] Siqueira AF, Maria FS, Mendes CM, Hamilton TR, Dalmazzo A, Dreyer TR, et al. Effects of photobiomodulation therapy (PBMT) on bovine sperm function. *Lasers Med Sci.* 2016; 31:1245-1250.
- [39] Ban Frangez H, Frangez I, Verdenik I, Jansa V, Virant Klun I. Photobiomodulation with light-emitting diodes improves sperm motility in men with asthenozoospermia. *Lasers Med Sci.* 2015; 30:235-240.
- [40] Yeste M, Codony F, Estrada E, Leonart M, Balasch S, Pena A, et al. Specific LED-based red light photo-stimulation procedures improve overall sperm function and reproductive performance of boar ejaculates. *Sci. Rep.* 2016; doi: 10.1038/srep22569.
- [41] Cohen N, Lubart R, Rubinstein S, Breitbart H. Light irradiation of mouse spermatozoa: stimulation of in vitro fertilization and calcium signals. *PhotochemPhotobiol.* 1998; 68:407-413.
- [42] Abdel-Salam Z, Dessouki SH, Abdel-Salam SA, Ibrahim MA, Harith MA. Green laser irradiation effects on buffalo semen. *Theriogenology.* 2011; 75:988-994.
- [43] Fernandes GH, Camillo de Carvalho PT, Serra AJ, Crespilho AM, Schatzman Peron JP, Rossato C, et al. The Effect of Low-Level Laser Irradiation

- on Sperm Motility, and Integrity of the Plasma Membrane and Acrosome in Cryopreserved Bovine Sperm. PLOS ONE. 2015; doi: 10.1371/journal.pone.0121487.
- [44] Corral-Baques MI, Rigau T, Rivera M, Rodriguez-Gil JE, Rigau J. Effect of 655-nm diode laser on dog sperm motility. *Lasers Med Sci.* 2005; 20:28-34.
- [45] Brito MF, Carvalho TT, Valle GR, Ferreira LFL, Mambrini JVM, Henry RJM, et al. Effect of low intensity red light on the canine semen cryopreservation. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia.* 2015; 67: 62-70.
- [46] Sato MH, Landthaler D, Hania D, Schill W-B. The Effects of Laser Light on Sperm Motility and Velocity in vitro. *Andrologia.* 1984; 16: 23-25.
- [47] Corral-Baques MI, Rigau T, Rivera M, Rodriguez-Gil JE, Rigau J. The effect of low-level laser irradiation on dog spermatozoa motility is dependent on laser output power. *Lasers Med Sci.* 2008; doi: 10.1007/s10103-008-0606-7.
- [48] Ocana-Quero JM, Gomez-Villamandos R, Moreno-Millan M, Santisteban-Valenzuela JM. Biological effects of helium-neon (He-Ne) laser irradiation on acrosome reaction in bull sperm cells. *J Photochem Photobiol B.* 199; 40:294-298.
- [49] Dreyera TR, Perez Siqueirab AF, Magrinia TD, Nichib M, Assumpcaob M, Milazzottoa MP, et al. Low-level laser irradiation-induced changes in bovine spermatozoa. 2014; arXiv: 1406.5234.
- [50] Montag M, Rink K, Delacretaz G, van der Ven H. Laser-induced immobilization and plasma membrane permeabilization in human spermatozoa. *Hum Reprod.* 2000; 15: 846-852.
- [51] Ebner T, Yaman C, Moser M, Sommergruber M, Hartl J, Tews G. Laser assisted immobilization of spermatozoa prior to intracytoplasmic sperm injection in humans. *Hum Reprod.* 2001; 16: 2628-2631.
- [52] Obruca A, Strohmer H, Sakkas D, Menezes Y, Kogosowski A, Barak Y, et al. Use of lasers in assisted fertilization and hatching. *Hum Reprod.* 1994; 9:1723-1726.
- [53] Gerber PA, Kruse R, Hirchenhain J, Krussel JS, Neumann NJ. Pregnancy after laser-assisted selection of viable spermatozoa before intracytoplasmic sperm injection in a couple with male primary cilia dyskinesia. *FertilSteril.* 2008; doi: 10.1016/j.fertnstert.2007.06.012.
- [54] Abdelmassih S, Cardoso J, Abdelmassih V, Dias JA, Abdelmassih R, Nagy ZP. Laser-assisted ICSI: a novel approach to obtain higher oocyte survival and embryo quality rates. *Hum Reprod.* 2002; 17: 2694-2699.
- [55] Rienzi L, Greco E, Ubaldi F, Iacobelli M, Martinez F, Tesarik J. Laser-assisted intracytoplasmic sperm injection. *FertilSteril.* 2001; 76:1045-1047.
- [56] Albertini DF, Carabatsos MJ. Comparative aspects of meiotic cell cycle control in mammals. *J Mol Med (Berl).* 1998; 76:795-799.
- [57] Soares CA, Annes K, Dreyer TR, Magrini T, Sonoda MT, da Silva Martinho H et al. Photobiological effect of low-level laser irradiation in bovine embryo production system. *J Biomed Opt.* 2014; 19: 35006.
- [58] Ocana-Quero JM, Gomez-Villamandos R, Moreno-Millan M, Santisteban-Valenzuela JM. Helium-Neon (He-Ne) laser irradiation increases the incidence of unreduced bovine oocytes during the first meiotic division in vitro. *Lasers Med Sci.* 1998; 13: 260-264.
- [59] Millan H, Ocana-Quero JH. Preliminary results of the evaluation of the use of clinical laser He-Ne radiation in the process of bovine "in vitro fertilization" *Bulletin UASVM, Veterinary Medicine.* 2009; 66:495.

**Citation:** Soheila Borhani, Reza Salman Yazdi. *Clinical Applications of Low-Level Laser Therapy in Reproductive Medicine: A Literature Review. Archives of Reproductive Medicine and Sexual Health.* 2018; 1(1): 14-21.

**Copyright:** © 2018 Soheila Borhani, Reza Salman Yazdi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.