

Hominins' Hurried Departure from Africa during the Out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

Michael Iannicelli *

* Independent Researcher, Brooklyn College (C.U.N.Y.), Brooklyn, N.Y.

*Corresponding Author: Michael Iannicelli *, postal address: 7109 21 Ave., Brooklyn, N.Y. 11204, michiann@optonline.net.

ABSTRACT

An established hypothesis explaining “The Out of Africa Event1” says that either a paleo-periglacial climate or a past increase in arid conditions was most probably the reason for *H. erectus*’ migration out of Africa and into Eurasia, 1.8 – 1.6 mya, but a study is offered here which blames frequent, lightning-strikes and accidental misuse of fire by hominins as the agents that led to uncontrolled, widespread wildfires. Paleo-winds advanced the wildfires according to wind direction, thus intensifying the widespread continental threat. A Pleistocene, temporary, land bridge, enabled some hominins to escape the threat of widespread wildfires by allowing them to cross over from east Africa to Eurasia. One anomaly noted is that either the wildfires or some vast impediment prevented equatorial *H. erectus* in Africa from climbing latitudinal lines up to north Africa at that time, but that delay to the mid-latitudes may have ended coincidentally with Eurasian travel on the eastern coastline of the juxtaposed Red Sea. The NW-SE oriented sea would have acted as a natural fire-retardant allowing hominins to finally move up to the mid-latitudes. The northbound migration would have also forced tropically-derived hominins to quickly adapt to low temperatures and environmental change while insight is offered here concerning how they may have successfully tackled their newest challenges.

Keywords: accidental misuse of fire; escape from wildfires; adaptation of *H. erectus* emigrants; directional paleo-winds influencing wildfires.

INTRODUCTION

Reasonable hypotheses in the literature attempting to explain why a species of hominins (*H. erectus*) initially migrated out of Africa, ca. 1.9 mya [1], are at a minimum, and so, the object of this study is to present a novel reason that emphasizes ‘widespread mobile wildfires’ as the driving force in chasing *H. erectus* out of the Africa. It is commonly thought that ‘The Out of Africa Event 1’, inter-continental movement of *H. erectus* hominins from out of Africa and into Eurasia was accomplished within a relatively fast period of time according to investigators such as Lockwood [2, p. 73]. A popular, literary hypothesis for ‘The Out of Africa Event 1’ departure by the hominins is generally attributed to paleo-climate change [3, p. 329] such that very arid conditions or cold, periglacial conditions may have reduced food provenances, in motivating hominins to migrate out of Africa. The paleo-environment of *H. erectus* hominins at the time between ca. 1.8 –

1.6 mya, was the arid grasslands [4] proximal to the equator, proven by hominal, fossilized bones found in east Africa. Agustí and Lordkipanidze [5] scrutinized and ultimately disproved secondary hypotheses but they also advocated their own hypothesis, whereby similar tools found in both Africa and at a hominal fossil site approx. 40° further, latitudinal north in Eurasia (Dmanisi, Georgia), somehow accounts for *H. erectus*’ flexibility of being able in adapting to low temperatures and environmental changes of the mid-latitudes. Nevertheless, the threat of widespread wildfires in Africa, as an influential factor for the initial flight of hominins from Africa is developed in this study within a perspective versus the primary hypothesis of paleo-climate change.

BACKGROUND

The migration of *H. erectus* out of Africa includes related hominins who were living at the same time as *H. erectus* such as *H. habilis* who existed 1.8 – 1.6 mya in Africa [6; 7]. The

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

preceding investigators [6; 7] remarked about Dmanisi's (in the country of Georgia) earliest resident, *H. erectus* (from 1.8 mya) possibly radiating from *H. habilis*-like ancestors who already had abandoned Africa. Both groups of hominins are now popularly classified together within the general *H. erectus* species [8] but only and strictly with an association to a specific timeframe of existence (which is 1.8 – 1.6 mya) for the purposes of the present study since *H. erectus* is generally regarded by many as having a very extended, exorbitant, time range of 1.8 mya – 100 kya. Although the topic of this extremely, lengthy, time range is only a very minor part of this study's thesis, the disposition here is that *H. erectus* should fall under a timeline of '1.8 mya to indefinitely while before the time of *H. sapiens*', but only for the sake of not debating against any general consensus that may claim a definitive time interval. Aside from that but even more importantly, the present study focuses only upon *H. erectus*' reason that forced his successful attempt in initially emigrating from Africa to Eurasia.

A description of *H. erectus*' paleo-habitat in east Africa (1.8 – 1.6 mya) is given here. As previously mentioned, his main environment was the grassland biome that originated in Africa due to: a certain combination of high temperatures; a barely adequate amount of rainfall; and a particular time-interval of dryness during the dry season that all together virtually prevented tree growth. The grasslands of Africa also typically happen to envelope a scattering of trees and very small areal forests. Above the northern rim of the grasslands, is the arid, Sahara Desert which experienced very extended time-intervals of a wet climate during the Pleistocene, between 2 – 1.6 mya [9]. Today, the grassland biome factually points to another important element in Africa, which is widespread wildfires, compelling authoritative environmentalists to declare Africa as the most fire-prone continent in the world [10; 11]. That alone, is one criterion towards the perspective of fire in chasing the hominins out of their home environment and ultimately out of Africa.

The usage of fire by *H. erectus* (Fig. 1) was for many reasons such as: cooking food; frightening predators away including insects; warmth and light [12]. Gowlett [13] mentions that hominins most probably learned about the useful aspects

of fire, when previous lightning-strikes ignited vegetative wildfires. But fires caused by hominins and by lightning are difficult to discern in the geological record [13]. He considers that hominins employed fire at a rough estimation of 1.8 mya while Bellomo & Kean [14] and Rowlett [15] both report the earliest evidence of fire used by African hominins (in Kenya), is dated at 1.5 mya based on burnt sediments containing stone tools that were apparently modified by fire. Proof of widespread paleo-fires is a sedimentary layer of charcoal but Scott [16] ascertains that even though grasslands anywhere are vastly and incessantly scorched during the long term, the yield of macroscopic charcoal in the rock record of the Cenozoic Era is scant at best because it is difficult to recognize in the field.

METHODOLOGY

The study's thesis is determined from many established facts derived from the peer-reviewed literature, circumstantial evidence, as well as indirect evidence that collectively add up to the study's conclusion of 'widespread mobile wildfires' that forced *H. erectus*' emigration from Africa and into Eurasia, which is itemized here:

Non-Controlled Fires Accidentally Ignited by Hominins

We've already introduced a presupposition of inevitable lightning-strikes igniting fires compounded by hominins who started using fire around 1.8 – 1.6 mya. But an immediate consequence arises from hominal usage of fire, which is the non-control of fire, that was accidentally and inevitably ignited by them (Fig. 1). The consequential effects were widespread smoke and the vast reduction of food resources, including animals that were killed and the disintegration of the hominins' edible vegetation. A net result of the perspective, widespread wildfires ignited by both lightning and the accidental misuse of fire by *H. erectus* is simply the impossibility of containing and extinguishing paleo-wildfires, especially during long dry seasons in paleo-Africa.

Modern-Day Analogy Reflecting the Potential Magnitude of Paleo-Wildfires

The realization of widespread wildfires as a reason for the hominins' dispersal out of Africa

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

is supported by comparing it to a modern-day analogy that could reflect the magnitude of hypothetical, African paleo-widespread wildfires. The Black Dragon Fire [17] that occurred in both China (Manchuria) and Russia (eastern Siberia) together, in May 1987, was the worse, widespread, wildfire recorded in history, in terms of the amount of seared land surface totaling to 72,843 km², while it took over a month's time for those wildfires to finally die out, mainly due to the tardy arrival of ample rain. Furthermore, that recent fire was accidentally ignited by just one person, which serves as a good example of whenever epitomizing paleo-scenarios of hominins who accidentally started nearly an infinite number of widespread wildfires during the advent and post-advent of fire usage in Africa. This is, of course, keeping in mind, the paleo-scenario's conjuncture of widespread wildfires also ignited by lightning.




Figure 1. Artwork depicting *H. erectus* hominins igniting a bonfire upon the open grasslands and their excitement of it, which implies that this typical paleo-scenario was repeated nearly an infinite amount of times by a hominin and / or his communities. Of course, one can also easily envision how a bonfire could escalate into an uncontrollable wildfire by spreading rapidly and according to wind direction. The artist drew them wearing garb but that actually wasn't the case for *H. erectus* at the time, 1.8 – 1.6 mya in Africa. Drawing by J. Matternes.

Wind as an Agent towards the Advancement of Wildfires

The perspective of a paleo-inferno in east Africa was powered not only by raging, widespread

wildfires but also by associated wind. The force and speed of wind together with the force of fire happen to spread the forward advancement of fire in particular directions. North of the equator, northeasterly trade winds blow toward the southwest, while south of the equator, southeasterly winds blow toward the northwest (Fig. 2). The climate of Africa back then during the Pleistocene Epoch was more or less similar to that of today [18], which implies that the wind regime of Pleistocene Africa may have been more or less similar to today's winds, although climate and wind deviated from that pattern whenever the influential, Pleistocene continental glacier was superimposed over northern Europe. In general, fire is also capable of burning in a backward or upwind direction by simple contact with the adjacent vegetation. Hence, the advancement of widespread wildfires would have been in directions opposite to one another, although its forward direction would have been exponentially faster than in its reverse direction.




Figure 2. Map of trade winds and other wind directions over Africa both north and south of the equator. The blowing direction of these winds would have simply advanced wildfires forward in the same direction. Wildfires may also advance backwards, since it would burn any adjacent vegetation in the upwind direction because of adjacent contact, but at a much, slower rate of speed.

Escape Route of *H. Erectus* during Ca. 1.8 – 1.6 Mya

Africa geo-physically connects to Eurasia where the extreme northeastern end of Egypt borders with the extreme southwestern end of Israel

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

(commonly referred to as the Sinai Peninsula), otherwise known as the 'Northern Dispersal Route (NDR)', which is a possible route traveled by *H. erectus* when he first entered Eurasia [1] (Fig. 3). But a different and more likely dispersal route taken by *H. erectus* was via the Horn of Africa (specifically Djibouti, Africa), and across the Bab el Mandeb Strait and into Eurasia, even though the strait geographically disconnects Africa from Eurasia during the Present, called the Southern Dispersal Route (SDR) [19, p. 38] (Figs. 3, 4). This had to have been accomplished by *H. erectus* when eustatic sealevel was lowered during the Pleistocene Epoch, 1.8 – 1.6 mya, due to

enormous, continental glacial ice growth occurring with tectonic fluctuation that created a temporary land bridge linking the Horn of Africa to neighboring Eurasia which is presently the southwestern end of Yemen on the Arabian Peninsula [20; 21; 19]. Petraglia [22] questioned the validity of a temporary land bridge existing there in the past, but Whalen and Schatte [23] nullified Petraglia [22] by citing a coincidental, archaeological site containing hard evidence of both Olduwan and Acheulean cultures located just past the east end of the Bab el Mandeb Strait on the Arabian Peninsula, 25 – 40 km inland.



Figure 3. Map of general paleo-migrations by *H. erectus* within Africa and Eurasia, ca. 1.8 – 1.6 mya. The map shows routes indicated by a starting point (No. 1) and a blue-dashed arrow up to point A; and a red-dotted line up to point B taken by hominins once they exited Africa and entered Eurasia. The migration within Eurasia also includes an optional route taken by hominins in a direction towards the Far East (blue-dashed line), which either further diverged towards the far north in China (point C), or towards the south within the land areas of the Pacific Ocean (purple-dashed arrows). The map also shows an alternate, northerly route (No. 2 and a yellow-dashed arrow) taken by hominins while within east Africa which was favored by other investigators and is known as the NDR (see text). A detailed look at *H. erectus*' journey up north once he was in Eurasia is shown in Figure 4. No. 1 = somewhere in east Africa; No. 2 = somewhere in east Africa. Point A = Ubeidiya, Israel;

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

point B = Dmanisi, Georgia; point C = Majuangou, north China. Based on Korisettar [24]; Zhu et al. [25] and Lahr [19].

Henceforth, the SDR is embraced in this study which also points to once *H. erectus* hominins were inside Eurasia, travel by them was to the north while hugging the extreme western side of the Arabian Peninsula's coastline, leading to further migration up to Ubeidiya, Israel,

followed by relocating much farther north to Dmanisi, Georgia [19, fig. 3.10] (Fig. 4). This was besides one other optional route chosen by other *H. erectus* hominins once they were on the Arabian Peninsula who headed towards the Far East, ca. 1.8 – 1.6 mya [24] (Fig. 3).



Figure 4. Close-up view of the Middle East route taken by *H. erectus* hominins (1.8 – 1.6 mya) after migrating over the Bab el Mandeb-land bridge at a time of lower sealevel during the Pleistocene Epoch, followed by their northward migration along the western coastline of the Arabian Peninsula, and then up to Ubeidiya, Israel, and further northward to Dmanisi, Georgia. The optional route where hominins elected to take when they were within the southern part of Eurasia is not shown in this figure but is shown in Figure3.

When hominins emigrated through the SDR during hypothetical, widespread wildfires burning in east Africa, their paleo-route up north on the western Arabian Peninsula was protected from those hypothetical fires by the enormity of the juxtaposed, NW-SE oriented, Red Sea (Figs. 3 and 4) which is 2,250 km long with an average width of 280 km. Specifically, it would have acted as either a natural shield or retardant to any easterly-spreading wildfires in east Africa (due to slow, reverse-directional advancement of wildfires) that happened to progress up to the west side of the Red Sea. That particular, geophysical barrier to fire, was most likely the springboard for hominins to initially gravitate towards the north at anytime during their existence within the 1.8 – 1.6 mya timeframe. Corroborating this, is the concentration of *H. erectus* archaeological sites in east Africa which bunch together in a slightly, curving, imaginary line leading up to near Djibouti, Africa where the Bab el Mandeb Strait connects to it (Fig. 5),

collectively pointing to the juncture of the SDR. There are no remains of *H. erectus* found in northern Africa, far away from the Horn of Africa, situated within east Africa. Thus, the present study follows Lahr [19, pp. 42 – 43] who negates an early Pleistocene, northerly, migrational route from the Horn of Africa up to the NDR based on no *H. erectus* bones found along that route or anywhere within northeastern Africa including the Sahara Desert during its extended, wet, climatic paleo-phases between 2 – 1.6 mya [9]. Plus, the oldest artifacts (i.e., tools) ever found in north Africa were dated from only 1 mya, as reported by Raynal et al. [26] and concurred by Lahr [19]. Those reasons leave this study to favor and focus mostly on *H. erectus*' entrance into the extreme southwestern end of the Arabian Peninsula and then blazing a trail up its western coastline in the north direction(Fig 4).

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

Circumstantial evidence serving as additional support for the above, would be a particular pattern of the spatial habitat occupied by pre-*H. erectus* hominins and *H. erectus* of Africa. The earliest of them, which were australopithecines or the “apelike, bipedal hominins”, are found in east Africa, south Africa and central Africa (country of Chad) [3, p. 282]. After them, is the rise of the genus “Homo”, namely *H. habilis* and *H. rudolphensis* who were contemporaneous with each other while they occupied only east Africa and south Africa [3, p. 292]. The early “Homo” are then immediately followed by late “Homo”, or *H. erectus*, and as previously discussed, they only occupied east Africa [3, p. 336]. Obviously, during progressive time within the preceding chain of evolution, one would easily expect an expansion of *H. erectus* into most parts of continental Africa (such as southern, western, central and north Africa), since their ancestors were well-distributed upon the African continent, but instead, we only see a contraction or shrinkage of the restricted, living zones of *H. erectus*, ultimately confined only to east Africa, ca. 1.8 – 1.6 mya. We may blame an environmental reason for that, namely, widespread paleo-wildfires mobilized by wind while not blaming a factor such as a paleo-periglacial climate or an increase in paleo-aridity. For the sake of argument, if there were any hypothetical, *H. erectus* hominins living in south Africa, then one can envision the forward advancement of hypothetical fires forcing them to move to east Africa when correlating that

with the southeasterly trade wind direction (Fig. 2). Likewise, any hypothetical, *H. erectus* hominins living in central Africa would be forced to move to east Africa because of northeasterly trade winds advancing, hypothetical fires (Fig. 2). In both of the preceding, two cases, the interpretation is, that the remaining amount of *H. erectus* hominins surviving in east Africa were paleo-positioned behind the advancing, rapid fires while they only had to worry about the very slow, backward progress of paleo-fires which was previously discussed (superimpose Figure 2 over Figure 5). The study considers that particular area of east Africa as a “pocket” of temporary safe refuge (refugium) since those *H. erectus* hominins were “behind” the hypothetical, rapid, forward advancing, paleo-fires mobilized by wind. Thus, the previously-mentioned “hurried departure” of the hominins from Africa (1.8 – 1.6 mya) was most probably triggered by a very abrupt or “fast” environmental condition such as the widespread, mobilized wildfires because an immediate disintegration of the hominins’ edible vegetative food would have forced constant emergency conditions that prompted an urgent, hominal escape from it. Also, it becomes clear that the “hurried departure” from Africa by hominins was not caused by climate such as a gradual increase in aridity because then only a corresponding, gradual exit by the hominins from Africa would have been induced.



Figure 5. The presence of all *H. erectus*' archaeological sites in east Africa that prove the existence of hominins both within and adjacent to the Horn of Africa (encompassed by Djibouti, Eritrea, Ethiopia and Somalia) not

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidently and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

very far from the Bab el Mandeb Strait (see Fig. 4). Granted, some of those sites are younger than the study's ca. 1.8 – 1.6 mya timeframe, but nevertheless, it's an indication of *H. erectus* preferring east Africa near the equator as their living zone since no other fossilized remains of *H. erectus* are found anywhere else outside of that zone within Africa. This apparently suggests the hominins quickly (relatively in time) funneling or bottlenecking themselves through a narrow corridor such as the Bab el Mandeb Strait (during a time of Pleistocene lower sea level) while literally three quarters of the remaining space within the continent of Africa was supposedly awaiting to accommodate *H. erectus*. That further implies the non-expansion of *H. erectus* far north of equatorial east Africa on the African continent. These biogeographic markers implicate the NDR (see text) as being an unlikely route taken by hominins at that time. The ages of the archaeological sites on the map are as follows: Koobi Fora, Kenya, 1.8 – 1.4 MYBP; Nariokotome, Kenya, 1.6 – 1.5 MYBP; Konso, Ethiopia, ca. 1.4 MYBP; Daka, Ethiopia, ca. 1 MYBP; Buia, Eritrea, ca. 1 MYBP; Olduvai Gorge, Tanzania, 1.2 MYBP – 700 KYBP. Based on Lockwood [2].

ADAPTATION TO LOW PALEO-TEMPERATURES AND A DIFFERENT PALEO-ENVIRONMENT

Relevant to the study's thesis, is the circumstance of "going from out of the fire and into the frying pan" which is a twist of an old European expression being used here whenever movement goes from a hellish situation to a less harsh but still bad situation, such as the different, general, paleo-temperatures and paleo-environment confronted by *H. erectus* during his transition from Africa to Eurasia, 1.8 – 1.6 mya. This not only complements the study's theme but also provides insight into any adaptation by tropically-derived hominins to specific, new challenges during their migration to the mid-latitudes. African *H. erectus*' fossilized remains were found proximal to the equator (Fig. 5) while the fossilized remains of Dmanisi's *H. erectus* (dated at 1.8 mya) were found at the 41° N parallel in Eurasia (Figs. 3 and 4), which simply indicate a very substantial and magnanimous, nomadic climb up north by tropically-derived, hominins. The extremely, large latitudinal difference promulgates the fact of low temperatures and environmental change, since the sun's radiation of the earth always and swiftly diminishes at latitudes above 30° [27]. That change was the transition of grasslands to woodlands [28; 29]. Belmaker [29] also designated a woodlands paleo-environment for the hominal, archaeological site in Ubeidiya, Israel (dated at 1.6 mya) at the 32.6° N parallel. The existence of paleo-woodlands there would also indicate that the hypothetical wildfires may not have stretched far into nor surpassed the northernmost point of east Africa, evidenced by whenever grasslands are protected from fire, they usually convert into woodlands [30].

Warmth was needed by tropically-derived *H. erectus* hominins when migrating to mid-latitudinal locations such as Ubeidiya (32.6° N)

and especially Dmanisi (41° N), 1.8 – 1.6 mya, but evidence of any form of clothing was never archaeologically detected for them (Neanderthal Man was the first to wear clothing). This signifies that small, controlled, campfires were used for warmth in combination with cave-dwelling while the retainment and the growth of more body hair over time, also figure into that equation. This is even whenever any journeys up to the far north experienced a limited time-window of interglacial conditions [31] which still would have seen low temperatures, although not as low as when it was under full glacial conditions. Yet, Dennell [32] claims that hominins such as *H. erectus* of Dmanisi, did not have to endure cold temperatures because he assumed that they only settled into their mid-latitude environment (at latitude 41° N) during the summer months followed by their evacuation from Dmanisi when cold temperatures arrived. However, his assumption is unconvincing because he didn't take into account of what the estimated temperatures were like in the last and first one-third leg of *H. erectus*' journey when they entered and exited Dmanisi, respectively. In other words, there would have been some specific time interval during the linking of the last leg of travel to Dmanisi and their settlement in Dmanisi, along with the first leg of their return trip to the south, that would not have seen the summer months since summer months generally range from May to September. This credible perception was calculated by determining what the temperatures were like when tropically-derived hominins migrated through different temperature zones during their journey in a direction either to or from Dmanisi. Starting from the northernmost point of travel, the coldest temperature at Dmanisi would have reached 5° C [32] while generally south of that, cold temperatures still dominated according to the Koppen-Geiger climate classification [33]. Specifically, the latter authors characterized the climate zone

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidently and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

almost immediately south of Dmanisi as an arid, cold steppe zone (Bsk) located presently in east Turkey bordering with the northwestern edge of Iran. The aspect of aridity in that steppe environment would have still been conducive to living by tropically-derived hominins since they originated from the arid grasslands of east Africa [4]. Directly south of that, is a cold-temperature zone with a dry and hot summer (Dsa), within present-day Iran [33]. Peel et al. [33] then identified, directly south of that, the beginning of the Temperate Zone which has dry and hot summers (Csa), located between present-day north Iraq and west Iran at the 37° N parallel. The northern edge of this last zone is considered here as the margin between the preceding cold zone of the above and a less harsh, temperature zone while this Csa zone may have or may not have been agreeable or hospitable to the tropically-derived hominins. In addition here, we must also factor in any escalatory, cold temperatures experienced by them which always occur with increased elevation or altitude, whenever hominins crossed highly-elevated, topographic, montane barriers.

In an attempt to simply illustrate the above, we may assign numerical values within a theorem that resonate with some degree of accuracy. A sum of approximately 444 km would have been traveled by hominins between Dmanisi (at the 41° N parallel) and the 37° N parallel (the northern edge of the Temperate Zone) whether journeying to or from Dmanisi. From that, we may estimate the total amount of days spent in nomadic, forward-motion, migration. Within reason, it is suggested here that it would have taken roughly, 222 days or > 7 months to cover the distance of 444 km at a rate of speed, let's say, averaging 2 km/day of hominal, nomadic, forward-motion walking when either trekking to or trekking away from Dmanisi. That rate of speed entailing the nomadic daily travel is computed by factoring in occurrences that inevitably happened, such as: stoppages made for rest or while foraging for food/water; slowdowns resulting from climbing over high topographical relief and crossing bodies of water (i.e., swamplands); slowdowns also occurring when eluding carnivores; and punctuated, forward-motion progress from whenever walking in a general zig-zag pattern for whatever reason (i.e., evading general obstacles, etc.). We should also take into consideration any further, inevitable lags such as

whenever a hominal individual/family/tribe settled into a specific, comfortable (convenient) setting or lodging for over a day's or week's time at any physical, geographic point during their trek to/from Dmanisi. Thus, the net result is, that an approximate constant of > 7 months of cold temperatures was experienced by *H. erectus* hominins either during the last leg of the journey to Damanis, Georgia, or the first leg of the journey heading away from there.

In order for *H. erectus* to protect himself against the encounter of low temperatures and environmental change, he had the herculean task of enduring and adapting to those adverse conditions that were forced upon him. Physiological adaptation by a group of hominins means, in the sense, that consequently there would have been a high number of casualties within the group during the short term, while only the strongest (i.e., possessing a very good immune system) and smartest or wisest (i.e., by being adroit) survived. An example of when only the shrewdest hominins were most likely to survive, is when we justifiably conjecture about how hominins may have managed fluctuating cold temperatures. When Dmanisi's coldest temperature would have reached 5° C [32], the combination of a thick layer of body hair and daytime, physical activity would have self-generated sustainable, body heat. But since caves were only intermittently available to *H. erectus* during his leg of the 444 km journey to and/or from Dmanisi, then cold nights may have seen only adept *H. erectus* hominins improvising a homemade sleeping bag by scooping out a shallow pit within dry, vegetated soil, and then piling and hand-compacting the loosened soil upon themselves, horizontally from toe to neck, for the sake of keeping warm during sleep. This is because a vegetative cover acts as an insulator, reducing heat loss from the soil that earlier absorbed and gained heat from the sun during daytime [27]. While asleep, any bothersome insects within the soil would have been buffered by the hominins' veneer of thick, body hair. A nearby lake or stream would have provided wash of the soil from off the body when awake. Further insight into relationships to the soil by nomadic hominins may include: storage of unspoiled food by burial and subsequent refrigeration within 'cold soil' or soil lacking a vegetative cover (such as soil always in the shade) while the food is wrapped in a big, fresh leaf (or leaves) to prevent moisture absorption from any groundwater

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidently and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

infiltration; and a self-plastering of mud upon their naked bodies for the sake of camouflaging or masking their human scent to evade carnivores [34].

Thus, it is realized that *H. erectus* survivors undoubtedly struggled against nature's elements during forced, physiological adaptation. This parallels Palmer's [35] assertion that hominins at that time had a low tolerance to adverse conditions, but at the same time, it is still counter to Dennell's [32] assumption, as previously discussed.

Lastly, since the low latitude (13° N) of the extreme southwestern corner of the Arabian Peninsula was the first point of migration after *H. erectus* hominins left Africa, it was also the pivotal point of where they made the choice of either: traveling up north (as previously discussed); or traveling somewhat within more or less the same low lines of latitudes towards the Far East (Fig. 3). The latter route did however, eventually split up (Fig. 3), since evidence of *H. erectus*' handiwork was found far up north at the 40° N parallel in Majuangou, north China, in the form of stone tools and modified animal bone, dated from 1.66 mya [25]. A possibility was raised by Ao et al. [36] claiming that these hominins initially arrived in Majuangou, north China by originally and laterally migrating from faraway Dmanisi, Georgia, 1.7 – 1.6 mya, based on the similarity of their lines of latitude. However, the claim by Ao et al. [36] is dubious because they contradicted themselves by adopting Dennell's [32] assumption as a reason why *H. erectus* hominins vacated their far northern settlement of Majuangou. This would imply that their hypothetical, very long, lateral paleo-journey directly from the west in Dmanisi to the east in Majuangou, was repeatedly detoured whenever cold temperatures correspondingly forced *H. erectus* hominins to divert deeply to the south, which precluded them from repeatedly returning to the 40° N parallel for the sake of resuming their migration towards Majuangou.

FUTURE RESEARCH

It is worth mentioning here for the purposes of future research, that the very same reason for the hominins' 'Out of Africa Event 1' may also be further extended to the lineages of African *H. erectus* hominins that are much younger than the timeframe of 1.8 – 1.6 mya, a theme which is outside the scope of the present study.

CONCLUSIONS AND SUMMARY

The advent and positive effects of fire-usage by *H. erectus* hominins is well established in the literature, while the negative or consequential effects of it was never reported by authorities. Thus, this study takes the opportunity to present a novel theory or perspective concerning the reason for *H. erectus*' forced, displacement from Africa and their concomitant entrance into Eurasia, commenced by the accidental misuse of fire, leading to the threat of mobilized, widespread wildfires associated with winds. The migrational direction favored by this study regarding *H. erectus*, ca. 1.8 – 1.6 mya, was through the Horn of Africa (which is at present, Djibouti) and into the Arabian Peninsula (the southwestern end of Yemen) where they then traveled north along its western coastline up to Ubeidya, Israel, followed by relocating further north to Dmanisi, Georgia. This was besides the optional route taken by other *H. erectus* hominins towards the Far East, who kept very roughly within the same, low, latitudinal lines as when they dwelled in east Africa. The challenges facing tropically-derived, *H. erectus* hominins at that time were titanic, not only because of the hypothetical, widespread wildfires in Africa but also because they endured a forced adaptation to low temperatures and environmental change during their migration north up to the 41° N parallel. A secondary but exciting conclusion generated by the study, is *H. erectus* possibly utilizing the soil in different ways to his advantage for many reasons such as: when he hypothetically dug a shallow pit in dry, vegetated soil ('warm soil'), and then refilling the pit while he lied in it, for the sake of creating a natural sleeping bag, which allowed him to cope with cold nights of the mid-latitudes in Eurasia.

The study is correct in verbally amplifying and not dramatizing a case for the grim paleo-situation in Africa but only because it is justified to portray it in this light. The central thesis does not replace the popular paleo-climate hypothesis, but instead, provides an alternative comprehension and original interpretation about the past events in Africa related to the theme of the present study.

We may now summarize the main points made by this study in order to justify its determination of its conclusions:

- The past discovery of an archaeological site showing evidence of *H. erectus*' usage of

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidently and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

fire, 1.5 mya, which was very coincidental with his first exit out of Africa, 1.8 – 1.6 mya, especially since other investigators theorize that the first actual usage of fire was more likely 1.8 mya. This correlates to H. erectus' accidental, inevitable misuse of fire combined with the ignition of fires by lightning-strikes in causing paleo-widespread, wildfires that pushed H. erectus out of Africa.

- The archaeological sites of H. erectus' fossilized remains are aligned in a slightly curving, south to north line in east Africa, adjacent to the Bab el Mandeb Strait's temporary, land bridge during the Pleistocene Epoch which is coincidental with H. erectus' point of escape when he fled from the hypothetical, widespread wildfires of east Africa, followed by his initial entrance into Eurasia.
- Modern-day, widespread wildfires occurring in 1987 within Russia and China, that charred > 72,000 km² of land area while raging on for over one month's time, is compared here to the study's advocacy of paleo-widespread wildfires in Africa. That serves as a small glimpse into the time of paleo-Africa because the 1987 widespread wildfire was most probably only a microcosm of paleo-Africa's wildfire regime, especially during droughts and long dry seasons in Africa, ca. 1.8 – 1.6 mya.
- The absence of H. erectus fossils above central Africa within north Africa bracketed by the study's timeframe, reflects some sort of generalized impediment to equatorial H. erectus that prevented him from moving to the northern part of Africa. This happens to be concurrent with the underestimated, geographical, NW-SE oriented, Red Sea acting as a fire-retardant to the hypothetical widespread wildfires of east Africa, which then allowed the hominal refugees to finally migrate safely immediately east of the Red Sea, and up north to the mid-latitudes, eventually routing them up to, at least, Ubeidiya, Israel at the 32.6° N parallel.
- The abstruse, relatively quick exit of H. erectus hominins from Africa is now comprehended when we blame widespread, mobilized paleo-wildfires as the agent in quickly destroying natural resources such as

their food which forced their urgent escape. This is in contrast to a gradual, climatic condition such as increased aridity that would have only caused a corresponding, gradual exit by the hominins, which was not the case.

- The characterization of Africa as being the 'most fire-prone country', is also taken into consideration here.
- Spatially large, paleo-wildfires within any given timeframe of world history, is ultimately proven by correlating them to a sedimentary layer of charcoal, but however, because macroscopic charcoal is practically almost impossible to identify in the Cenozoic rock record [16], then we must rely on other collective data such as the established facts of the literature, circumstantial evidence, and indirect evidence, to help us suspect that the threat of widespread, paleo-wildfires was a feasible, motivating factor in forcing particular paleo-migrations by H. erectus hominins, ca. 1.8 – 1.6 mya.
- Supplementing the thesis of the study, is an inclusion of relevant information identifying the inevitable challenges confronted and resolved by tropically-derived hominins after their evacuation of Africa when they were in the process of relocating to the mid-latitudes such as Dmanisi, Georgia at 41° N, ca. 1.8 – 1.6 mya. They reluctantly and ingeniously had to adapt to different environmental and temperature conditions because their only option was to experience and overcome those two adverse conditions, or else, face a repulsive 'Africa on fire'.

ACKNOWLEDGEMENTS

The author thanks John S. Compton for being informative while providing some references.

REFERENCES

- [1] Carotenuto, F., Tsikaridze, N., Rook, L., Lordipandze, D., Longo, L., et al. (2016). Venturing out safely: the biogeography of Homo Erectus' dispersal out of Africa. *Journal of Human Evolution*, 95, 1 – 12.
- [2] Lockwood, C. (2013). *The Human Story: Where We Come from and How We Evolved*. Natural History Museum: London
- [3] Lewin, R. and Foley, R. (2004). *Principles of Human Evolution*. Blackwell Pub. : Oxford, U.K.

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidently and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

- [4] deMenocal, P.B. (2004). African climate change and faunal evolution during the Pliocene – Pleistocene. *Earth and Planetary Science Letters*, 220, 3 – 24.
- [5] Agusti, J. and Lordkipanidze, D. (2011). How “African” was the early human dispersal out of Africa? *Earth Science Reviews.*, 30, 1338 – 1342.
- [6] Vekua, A., Lordkipanidze, D., Rightmire, G.P., Agusti, J., Ferring, R., et al. (2002). A new skull of early Homo from Dmanisi, Georgia. *Science*, 297, 85 – 89.
- [7] Lordkipanidze, D., Jashshvili, T., Vekua, A., Ponce deLeon, M.S., Zollikofer, P.E., et al. (2007). Postcranial evidence from early homo from Dmanisi, Georgia. *Nature*, 449, 305 – 310.
- [8] Henderson, J.L. (2015). Dmanisi: a taxonomic revolution. *Pure Insights*, v. 4 (4).
- [9] Larrasoana, J.C., Roberts, A.P., Rohling, E.J. (2013). Dynamics of Green Sahara and their role in hominin evolution. *PLOS One*, 8 (10), 1 – 12.
- [10] Bowman, D.M.J.S., Balch, J., Artaxo, P., Bond, W.J., Cochrane, M.A., et al. (2011). The human dimension of fire regimes on earth. *Journal of Biogeography*, 38, 2223 – 2236.
- [11] Daniau, A-L, Goni, M.F.S. , Martinez, P., Urrego, D.H., Bout-Roumazelles, V., et al. (2013). Orbital-scale climate forcing of grassland burning in southern Africa. *PNAS*, 110, (3), 5069 - 5073.
- [12] Ko, K.H. (2016). Origins of human intelligence: the chain of tool-making and brain evolution. *Anthropological Notebooks*, 22 (1), 5 – 22.
- [13] Gowlett, J.A.J. (2016). The discovery of fire by humans: a long and convoluted process. *Philosophical Transactions Royal Society B*, 371 (1696).
- [14] Bellomo, R.V. & Kean, W.F. (1997). Evidence of hominid controlled fire at the FxJj20 site complex, Karari escarpment. Appendix 4A. IN, G.L. Issac and B. Issac (eds.), “Plio-Pleistocene Archaeology”, Koobi Foro Research Project. v. 5 (pp. 224 – 233). Clarendon Press: Oxford, U.K.
- [15] Rowlett, R.M. (2000) Fire control by Homo erectus in East Africa and Asia. *Anthropologica Sinica*, 19, 198 – 208.
- [16] Scott, A.C. (2010). Charcoal recognition, taphonomy and uses in paleoenvironmental analysis. *Paleogeogr. Paleoclim. and Paleocol.*, 291, 11 – 39.
- [17] Salisbury, H.E. (1989). *The Great Black Dragon Fire: A Chinese Inferno*. Little, Brown and Co: New York.
- [18] Maslin, M.A., Brierly, C.M., Milner, A.M., Schultz, S., Trauth, M.H., et al. (2014). East African climate pulses and early human evolution. *Quaternary Science Reviews*, 101, 1 – 17.
- [19] Lahr, M.M. (2010). Saharan corridors and their role in the evolutionary geography of ‘Out of Africa Event 1’. IN, J.G. Fleagle et al. (eds.), “Out of Africa 1: The First Hominin Colonization of Eurasia”, (pp. 27 – 46), *Vertebrae Paleobiology and Paleoanthropology Series*, Springer Science: many locations.
- [20] Haq, B.U., Hardenbol, J., Vail, P.R. (1987). Chronology of fluctuating sealevels since the Triassic. *Science*, 235, 1156 – 1167.
- [21] Turner, A. (1999). Assessing earliest human settlement of Eurasia: late Pliocene diversions from Africa. *Antiquity*, 73, 63 – 570.
- [22] Petraglia, M.D. (2003). The lower Paleolithic of the Arabian Peninsula: occupations, adaptations, and dispersals. *Journal of World Prehistory*, 17 (2), 141 - 179.
- [23] Whalen, N.M. and Schatte, K.E. (1997). Pleistocene sites in southern Yemen. *Arabian Archaeology and Epigraphy*, 8, 1 – 10.
- [24] Korisettar, R. (2016). Out of Africa and into south Asia: the evidence from Paleolithic archaeology. IN, G.R. Schug and S.R. Walimbe (eds.), “A Companion to South Asia in the Past”, (pp. 60 – 71). John Wiley & Sons: many locations.
- [25] Zhu, R.X., Potts, R., Xie, F., Hoffman, K.A., Deng, C.L., et al. (2004). New evidence on the earliest human presence at high northern latitudes in northeast Asia. *Nature*, 431, 559 – 562.
- [26] Raynal, J.P., Sbihi-Alaoui, F.Z., Geraads, D., Magoga, L., Mohib, A. (2001). The earliest occupation of north Africa: the Moroccan perspective. *Quaternary International.*, 75 (1), 65 – 75.
- [27] Jury, W.A. and Horton, R. (2004). *Soil Physics*. J. Wiley & Sons: New Jersey.
- [28] Gabuina, L. Anton, S.C., Lordkipanidze, D., Vekua, A., Justus, A., et al. (2001). Dmanisi and dispersal. *Evolutionary Anthropology* 10, 158 – 170.
- [29] Belmaker, M. (2010). Early Pleistocene faunal connections between Africa and Eurasia: an ecological perspective. IN, J.G. Fleagle, et al. (eds.), “Out of Africa 1: the First Hominid colonization of Eurasia” (pp. 183 – 205), *Vertebrae Paleobiology and Paleoanthropology*, Springer Science: Heidelberg, London and New York.
- [30] Keeley, J.E. and Rundel, P.W. (2005). Fire and the Miocene expansion of C₄ grasslands. *Ecology Letters*, 8, 683 – 690.

Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya

- [31] Leroy, S.A.G., Arpe, K. and Mikolajewicz, U. (2011). Vegetation context and climatic limits of the earlyPleistocene hominin dispersal in Europe. *Quaternary Science Reviews*, 30, 1448 – 1463.
- [32] Dennell, R.W. (2012). The Nihewan Basin of north China in the early Pleistocene: continuous and flourishing, or discontinuous, infrequent and ephemeral occupation. *Quaternary International*, 295, 233 – 236.
- [33] Peel, M.C., Finlayson, B.L., and McMahon, T.A. (2007). Updated world map of the Koppen-Geiger climate classification. *Hydrology and Earth System Sciences*, 11, 1633 – 1644.
- [34] Sorensen, E. (no date). LiveOutdoors: Six Ways to Mask Your Scent [home page on the internet]. [Cited 12 September, 2018]. Available from: <http://www.liveoutdoors.com/hunting/229803-6-ways-to-mask-your-scent/>
- [35] Palmer, D. (2010). *Origins: Human Evolution Revealed*. Mitchell Beazley: London.
- [36] Ao, H., Dekkers, M.J., Qi, W., Qiang, X., Xiao, G. (2013). New evidence for early presence of hominids in north China. *Scientific Reports*, 3, 2403.

Citation: Michael Iannicelli , " Hominins' Hurried Departure from Africa during the out of Africa Event-1 due to Widespread Wildfires, Accidentally and Inevitably Ignited by them, C.A. 1.8 – 1.6 Mya", *Research Journal of Food and Nutrition*, vol.1, no.2, pp. 1-12, 2018.

Copyright:© 2018 Michael Iannicelli, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.